


Quadrilaterals

POINTS TO REMEMBER

- 1. Quadrilateral.** A closed four sided figure is called a quadrilateral.
 - (i) It has four sides, four vertices, four angles and two diagonals.
 - (ii) Sum of its four angles = 360° i.e. $\angle A + \angle B + \angle C + \angle D = 360^\circ$.




2. Types of Quadrilaterals

- 1. Parallelogram.** A quadrilateral in which opposite sides are parallel, is called a parallelogram.

In the given figure, ABCD is a quadrilateral in which $AB \parallel DC$ and $AD \parallel BC$.

\therefore ABCD is a parallelogram.


Its opposite sides are equal i.e. $AB = CD$ and $AD = BC$.



- 2. Rhombus.** A parallelogram having all sides equal, is called a rhombus.

In the given figure, ABCD is a parallelogram in which $AB = BC = CD = DA$.


\therefore ABCD is a rhombus.



- 3. Rectangle.** A parallelogram each of whose angles measures 90° , is called a rectangle.

In the given figure, ABCD is a rectangle.

Its opposite sides are equal.




- 4. Square.** A rectangle having all sides equal, is called a square.

In the given figure, PQRS is a square, in which

$PQ = QR = RS = SP$.

Its each angle is of 90° .



5. **Trapezium.** A quadrilateral having two parallel opposite sides and two non-parallel opposite sides is called a trapezium.

In the given figure, ABCD is a quadrilateral in which $AB \parallel DC$ and AD is not parallel to BC.

\therefore ABCD is a trapezium.

If the non-parallel sides of a trapezium are equal, then it is called an **isosceles trapezium**.

6. **Kite.** A quadrilateral in which two pairs of adjacent sides are equal, is known as a kite.

In the given figure, ABCD is a quadrilateral in which $AB = AD$ and $CB = CD$.

\therefore ABCD is a kite.




Results on Parallelogram

Theorem 1. In a parallelogram :

- the opposite sides are equal ;
- the opposite angles are equal ;
- each diagonal bisects the parallelogram.


Theorem 2. The diagonals of a parallelogram bisect each other.

Theorem 3. If a pair of opposite sides of a quadrilateral are parallel and equal, then it is a parallelogram



EXERCISE 15 (A)

- Q. 1. In the given figure, ABCD is a parallelogram in which $\angle A = 70^\circ$. Calculate $\angle B$, $\angle C$ and $\angle D$.



Sol. \therefore ABCD is a parallelogram.

$$\therefore \angle A = \angle C \quad \text{and} \quad \angle B = \angle D$$

$$\therefore \angle C = \angle A = 70^\circ$$

$$\text{But } \angle A + \angle B = 180^\circ$$

(Co. interior angles)

$$\Rightarrow 70^\circ + \angle B = 180^\circ$$

$$\Rightarrow \angle B = 180^\circ - 70^\circ = 110^\circ$$


$$\text{But } \angle D = \angle B$$

$$\therefore \angle D = 110^\circ$$

$$\text{Hence } \angle B = 110^\circ, \angle C = 70^\circ$$

$$\text{and } \angle D = 110^\circ \text{ Ans.}$$

- Q. 2. In the given figure, ABCD is a parallelogram. Side DC is produced to E and $\angle BCE = 105^\circ$.



Calculate $\angle A$, $\angle B$, $\angle C$ and $\angle D$.

Sol. ABCD is a parallelogram.

Side DC is produced to E

and $\angle BCE = 105^\circ$

But $\angle BCE + \angle BCD = 180^\circ$

(Linear pair)

$$\Rightarrow 105^\circ + \angle BCD = 180^\circ$$

$$\Rightarrow \angle BCD = 180^\circ - 105^\circ = 75^\circ$$

$$\text{or } \angle C = 75^\circ$$

$$\text{But } \angle A = \angle C$$

(Opposite angles of a parallelogram)

$$\therefore \angle A = 75^\circ$$

$$\therefore AB \parallel CD$$

$$\therefore \angle BCE = \angle CBA$$

(Alternate angles)

$$\Rightarrow \angle CBA = 105^\circ \text{ or } \angle B = 105^\circ$$


$$\text{But } \angle D = \angle B$$

(Opposite angles of a parallelogram)

$$\therefore \angle D = 105^\circ$$

$$\text{Hence } \angle A = 75^\circ, \angle B = 105^\circ, \angle C = 75^\circ, \angle D = 105^\circ \text{ Ans.}$$

- Q. 3.** If an angle of a parallelogram is two-third of its adjacent angle, find the angles of the parallelogram.



Sol. In parallelogram ABCD,

$$\text{Let } \angle B = x$$

$$\text{then } \angle A = \frac{2}{3}x$$

$$\text{But } \angle A + \angle B = 180^\circ$$

(Co. interior angles)

$$\Rightarrow \frac{2}{3}x + x = 180^\circ \Rightarrow \frac{5}{3}x = 180^\circ$$

$$\Rightarrow x = 180^\circ \times \frac{3}{5} = 108^\circ$$

$$\therefore \angle B = 108^\circ \text{ and } \angle A = \frac{2}{3} \times 108^\circ = 72^\circ$$

$$\text{But } \angle C = \angle A$$

(Opposite angles of a parallelogram)


$$\therefore \angle C = 72^\circ$$

$$\text{Similarly } \angle D = \angle B = 108^\circ$$

Hence $\angle A = 72^\circ, \angle B = 108^\circ, \angle C = 72^\circ$ and $\angle D = 108^\circ$ Ans.

- Q. 4.** In the adjoining figure, ABCD is a parallelogram in which $\angle BAD = 70^\circ$ and $\angle CBD = 50^\circ$. Calculate :

- (i) $\angle ADB$ (ii) $\angle CDB$.



Sol. ABCD is a parallelogram. BD is joined.

$$\angle BAD = 70^\circ \text{ and } \angle CBD = 50^\circ$$

$$\therefore AD \parallel BC$$

$$\therefore \angle ADB = \angle CBD$$

(Alternate angles)

$$= 50^\circ$$

$$\text{But } \angle BAD + \angle ADC = 180^\circ$$

(Co. interior angles)

$$\Rightarrow \angle BAD + \angle ADB + \angle CDB = 180^\circ$$

$$\Rightarrow 70^\circ + 50^\circ + \angle CDB = 180^\circ$$


$$\Rightarrow 120^\circ + \angle CDB = 180^\circ$$

$$\Rightarrow \angle CDB = 180^\circ - 120^\circ = 60^\circ$$

$$\text{Hence (i) } \angle ADB = 50^\circ \text{ and}$$

$$(ii) \angle CDB = 60^\circ \text{ Ans.}$$

- Q. 5.** In the given figure, ABCD is a rhombus in which $\angle A = 72^\circ$. If $\angle CBD = x^\circ$, find the value of x .



Sol. ABCD is a rhombus.

$$\angle A = 72^\circ \text{ and } \angle CBD = x^\circ$$

\therefore ABCD is a rhombus

\therefore Diagonal BD bisects $\angle B$ and $\angle D$

$\therefore \angle ABD = \angle CBD = x$

$\Rightarrow \angle ABC = x + x = 2x$

But $\angle A + \angle B = 180^\circ$


(Co. interior angles)

$\Rightarrow 72^\circ + 2x = 180^\circ$

$\Rightarrow 2x = 180^\circ - 72^\circ = 108^\circ$

$\therefore x = \frac{108}{2} = 54^\circ$ Ans.

Q. 6. In the adjoining figure, equilateral $\triangle EDC$ surmounts square ABCD. If $\angle DEB = x^\circ$, find the value of x .



Sol. In the figure, ABCD is a square and $\triangle CDE$ is an equilateral. BE is joined.

$\angle DEB = x^\circ$

In $\triangle BCE$, $BC = CE = CD$

$\therefore \angle CBE = \angle CEB$

and $\angle BCE = \angle BCD + \angle DCE$

$= 90^\circ + 60^\circ = 150^\circ$

But $\angle BCE + \angle CBE + \angle CEB = 180^\circ$

(Angles of a triangle)

$\Rightarrow 150^\circ + \angle CEB + \angle CEB = 180^\circ$

$\Rightarrow 150^\circ + 2\angle CEB = 180^\circ$

$\Rightarrow 2\angle CEB = 180^\circ - 150^\circ = 30^\circ$

$\therefore \angle CEB = \frac{30^\circ}{2} = 15^\circ$

But $\angle CED = 60^\circ$


(Angle of an equilateral triangle)

$\Rightarrow x^\circ + \angle CEB = 60^\circ$

$\Rightarrow x^\circ + 15^\circ = 60^\circ \Rightarrow x^\circ = 60^\circ - 15^\circ = 45^\circ$

$\therefore x = 45$ Ans.

Q. 7. In the adjoining figure, ABCD is a rhombus whose diagonals intersect at O. If $\angle OAB : \angle OBA = 2 : 3$, find the angles of $\triangle OAB$.



Sol. ABCD is a rhombus and its diagonals bisect each other at right angles at O.

$\angle OAB : \angle OBA = 2 : 3$

Let $\angle OAB = 2x$ and $\angle OBA = 3x$

But $\angle AOB = 90^\circ$

$\therefore \angle OAB + \angle OBA = 90^\circ$

$\Rightarrow 2x + 3x = 90^\circ \Rightarrow 5x = 90^\circ$


$\therefore x = \frac{90^\circ}{5} = 18^\circ$

$\therefore \angle OAB = 2x = 2 \times 18^\circ = 36^\circ$

$\angle OBA = 3x = 3 \times 18^\circ = 54^\circ$

and $\angle AOB = 90^\circ$ Ans.

Q. 8. In the given figure, ABCD is a rectangle whose diagonals intersect at O. Diagonal AC is produced to E and $\angle ECD = 140^\circ$. Find the angles of $\triangle OAB$.



Sol. ABCD is a rectangle and its diagonals AC and BD bisect each other at O.

Diagonal AC is produced to E such that $\angle ECD = 140^\circ$

$$\angle ECD + \angle DCO = 180^\circ \text{ (Linear pair)}$$

$$\Rightarrow 140^\circ + \angle DCO = 180^\circ$$

$$\Rightarrow \angle DCO = 180^\circ - 140^\circ = 40^\circ$$

$$\text{But } OC = OD$$

(Half of equal diagonals)

$$\therefore \angle CDO = \angle DCO = 40^\circ$$

$$\text{Now } \because AB \parallel CD$$

(Opposite sides of a rectangle)

$$\therefore \angle OAB = \angle DCO$$

(Alternate angles)

$$= 40^\circ$$

$$\text{Similarly } \angle OBA = 40^\circ$$

$$\text{Now in } \triangle AOB$$

$$\angle OBA + \angle OAB + \angle AOB = 180^\circ$$


(Angles of a triangle)

$$\Rightarrow 40^\circ + 40^\circ + \angle AOB = 180^\circ$$

$$\Rightarrow 80^\circ + \angle AOB = 180^\circ$$

$$\Rightarrow \angle AOB = 180^\circ - 80^\circ = 100^\circ \text{ Ans.}$$

- Q. 9.** In the given figure, ABCD is a kite whose diagonals intersect at O. If $\angle DAB = 54^\circ$ and $\angle BCD = 76^\circ$, calculate : (i) $\angle ODA$ (ii) $\angle OBC$.



Sol. In the figure,

ABCD is a kite

$$\therefore AB = AD, BC = DC$$

Its diagonals AC and BD intersect at O.

$$\angle DAB = 54^\circ \text{ and } \angle BCD = 76^\circ$$

In $\triangle BCD$,

$$\angle CDB = \angle CBD (\because BC = DC)$$

$$\text{But } \angle BCD + \angle CDB + \angle CBD = 180^\circ$$

$$\Rightarrow 76^\circ + \angle CBD + \angle CDB = 180^\circ$$

$$\Rightarrow 76^\circ + 2 \angle CBD = 180^\circ$$

$$\Rightarrow 2 \angle CBD = 180^\circ - 76^\circ = 104^\circ$$

$$\therefore \angle CBD = \frac{104^\circ}{2} = 52^\circ$$

$$\text{or } \angle OBC = 52^\circ$$

Similarly in $\triangle ABD$,

$$\angle DAB = 54^\circ \text{ and } \angle ABD = \angle ADB$$

($\because AB = AD$)

$$\text{But } \angle DAB + \angle ABD + \angle ADB = 180^\circ$$

$$\Rightarrow 54^\circ + \angle ADB + \angle ADB = 180^\circ$$

$$\Rightarrow 54^\circ + 2 \angle ADB = 180^\circ$$

$$\Rightarrow 2 \angle ADB = 180^\circ - 54^\circ = 126^\circ$$


$$\therefore \angle ADB = \frac{126^\circ}{2} = 63^\circ$$

$$\text{or } \angle ODA = 63^\circ$$

$$\text{Hence } \angle ODA = 63^\circ$$

$$\text{and } \angle OBC = 52^\circ \text{ Ans.}$$

- Q. 10.** In the given figure, ABCD is an isosceles trapezium in which $\angle CDA = 2x^\circ$ and $\angle BAD = 3x^\circ$.



Find all the angles of the trapezium.

Sol. ABCD is an isosceles trapezium in which $AD = BC$ and $AB \parallel CD$.

$$\angle BAD + \angle CDA = 180^\circ$$

(Co. interior angles)

$$\Rightarrow 3x + 2x = 180^\circ \Rightarrow 5x = 180^\circ$$

$$\therefore x = \frac{180^\circ}{5} = 36^\circ$$

$$\therefore \angle A = 3x = 3 \times 36^\circ = 108^\circ$$

$$\angle D = 2x = 2 \times 36^\circ = 72^\circ$$

\therefore ABCD is an isosceles trapezium.

$$\therefore \angle A = \angle B \text{ and } \angle C = \angle D$$

$$\therefore \angle B = 108^\circ$$

and $\angle C = 72^\circ$


Hence $\angle A = 108^\circ$, $\angle B = 108^\circ$,

$\angle C = 72^\circ$, $\angle D = 72^\circ$ Ans.

Q. 11. In the given figure, ABCD is a trapezium in which

$\angle A = (x + 25)^\circ$, $\angle B = y^\circ$, $\angle C = 95^\circ$

and $\angle D = (2x + 5)^\circ$. Find the values of x and y .



Sol. In trapezium ABCD

$\angle A = (x + 25)^\circ$, $\angle B = y^\circ$, $\angle C = 95^\circ$ and

$\angle D = (2x + 5)^\circ$

$\angle A + \angle D = 180^\circ$ (Co. interior angles)

$$\Rightarrow (x + 25)^\circ + (2x + 5)^\circ = 180^\circ$$

$$\Rightarrow x + 25^\circ + 2x + 5^\circ = 180^\circ$$

$$\Rightarrow 3x + 30^\circ = 180^\circ$$

$$\Rightarrow 3x = 180^\circ - 30^\circ = 150^\circ$$

$$\therefore x = \frac{150^\circ}{3} = 50^\circ$$


Similarly, $\angle B + \angle C = 180^\circ$

$$\Rightarrow y + 95^\circ = 180^\circ$$

$$\Rightarrow y = 180^\circ - 95^\circ = 85^\circ$$

Hence $x = 50^\circ$, $y = 85^\circ$ Ans.

Q. 12. In the given figure, ABCD is a rhombus and $\triangle EDC$ is equilateral. If $\angle BAD = 78^\circ$, calculate : (i) $\angle CBE$ (ii) $\angle DBE$.



Sol. (i) ABCD is a rhombus and $\triangle EDC$ is an equilateral triangle,

$$\angle BAD = 78^\circ$$

$$\therefore \angle BCD = \angle A = 78^\circ$$

(Opposite angles of a rhombus)

$$\therefore \angle BCE = \angle BCD + \angle DCE$$

$$= 78^\circ + 60^\circ = 138^\circ$$

$$\therefore BC = CE$$

$$\therefore \angle CBE = \angle CEB$$

$$\text{But } \angle CBE + \angle CEB + \angle BCE = 180^\circ$$

(Sum of angles of a triangle)

$$\Rightarrow \angle CBE + \angle CBE + 138^\circ = 180^\circ$$

$$\Rightarrow 2 \angle CBE = 180^\circ - 138^\circ = 42^\circ$$

$$\therefore \angle CBE = \frac{42^\circ}{2} = 21^\circ$$

Now in $\triangle ABD$,

(ii) $AB = AD$ (Sides of a rhombus)

$$\therefore \angle ABD = \angle ADB$$

$$\text{But } \angle ABD + \angle ADB + \angle BAD = 180^\circ$$

$$\Rightarrow \angle ABD + \angle ABD + 78^\circ = 180^\circ$$

$$\Rightarrow 2 \angle ABD + 78^\circ = 180^\circ$$

$$\Rightarrow 2 \angle ABD = 180^\circ - 78^\circ = 102^\circ$$

$$\therefore \angle ABD = \frac{102^\circ}{2} = 51^\circ$$

$$\text{But } \angle BAD + \angle ABC = 180^\circ$$

(Co. interior angles)

$$\Rightarrow 78^\circ + \angle ABC = 180^\circ$$

$$\Rightarrow \angle ABC = 180^\circ - 78^\circ = 102^\circ$$

$$\Rightarrow \angle ABD + \angle DBE + \angle CBE = 102^\circ$$

$$\Rightarrow 51^\circ + \angle DBE + 21^\circ = 102^\circ$$

$$\Rightarrow 72^\circ + \angle DBE = 102^\circ$$


$$\Rightarrow \angle DBE = 102^\circ - 72^\circ$$

$$= 30^\circ$$

Hence $\angle CBE = 21^\circ$

and $\angle DBE = 30^\circ$ Ans.

Q. 13. DEC is an equilateral triangle in a square ABCD. If BD and CE intersect at O and $\angle COD = x^\circ$, find the value of x .



Sol. ABCD is a square and $\triangle ECD$ is an equilateral triangle. Diagonal BD and CE intersect each other at O, $\angle COD = x^\circ$.

\therefore BD is the diagonal of square ABCD

$$\therefore \angle BDC = \frac{90^\circ}{2} = 45^\circ \Rightarrow \angle ODC = 45^\circ$$

$$\angle ECD = 60^\circ$$

(Angle of equilateral triangle)

or $\angle OCD = 60^\circ$

Now in $\triangle OCD$,

$$\angle OCD + \angle ODC + \angle COD = 180^\circ$$

(Angles of a triangle)


$$\Rightarrow 45^\circ + 60^\circ + x^\circ = 180^\circ$$

$$\Rightarrow 105^\circ + x^\circ = 180^\circ$$

$$\therefore x^\circ = 180^\circ - 105^\circ = 75^\circ$$

Hence $x = 75$ **Ans.**

Q. 14. If one angle of a parallelogram is 90° , show that each of its angles measures 90° .



Sol. Given : ABCD is a parallelogram and $\angle A = 90^\circ$.

To prove : Each angle of the parallelogram ABCD is 90° .

Proof : $\therefore \angle A = \angle C$

(Opposite angles of a parallelogram)

$$\therefore \angle C = 90^\circ \quad (\because \angle A = 90^\circ)$$

$$\text{But } \angle A + \angle D = 180^\circ$$

(Co. interior angles)

$$\Rightarrow \angle D = 180^\circ - 90^\circ = 90^\circ$$

$$\text{and } \angle B = \angle D$$

(Opposite angles of a parallelogram)


$$\therefore \angle B = 90^\circ$$

$$\text{Hence } \angle B = \angle C = \angle D = 90^\circ$$

Q.E.D.

Q. 15. In the adjoining figure, ABCD and PQBA are two parallelograms. Prove that :

- (i) DPQC is a parallelogram.
- (ii) $DP = CQ$.
- (iii) $\triangle DAP \cong \triangle CBQ$.



Sol. Given : ABCD and PQBA are two parallelogram PD and QC are joined.

To prove : (i) DPQC is a parallelogram.

- (ii) $DP = CQ$
- (iii) $\triangle DAP \cong \triangle CBQ$.

Proof : (i) $DC \parallel AB$ and $AB \parallel PQ$

(Given)

$$\therefore DC \parallel PQ$$

$$\text{Again } DC = AB \text{ and } AB = PQ$$

(Opposite sides of parallelograms)

$$\therefore DC = PQ$$

$$\therefore DC = PQ \text{ and } DC \parallel PQ$$

$$\therefore DPQC \text{ is a parallelogram.}$$

- (ii) $\therefore DA = CQ$

(Opposite sides of parallelogram)

- (iii) In $\triangle DAP$ and $\triangle CBQ$

$$DA = CB$$

{Opposite sides of a parallelogram}

$$AP = BQ$$

(Opposite sides of parallelogram)

$$PD = CQ \quad (\text{Proved})$$

$$\therefore \triangle DAP \cong \triangle CBQ$$


(S.S.S. axiom of congruency)

Hence proved.

Q. 16. In the adjoining figure, ABCD is a parallelogram. $BM \perp AC$ and $DN \perp AC$.

Prove that : (i) $\triangle BMC \cong \triangle DNA$.

(ii) $BM = DN$.



Sol. **Given :** ABCD is a parallelogram.

$BM \perp AC$ and $DN \perp AC$.

To prove :

(i) $\triangle BMC \cong \triangle DNA$

(ii) $BM = DN$.

Proof : In $\triangle BMC$ and $\triangle DNA$

$$BC = AD$$

(Opposite sides of a parallelogram)

$$\angle M = \angle N \quad (\text{Each } 90^\circ)$$

$$\angle BCM = \angle DAN$$

(Alternate angles)

$$(i) \therefore \triangle BMC \cong \triangle DNA$$

(AAS axiom of congruency)

$$(ii) \therefore BM = DN \quad (\text{C.P.C.T.})$$


Hence proved.

Q. 17. In the adjoining figure, ABCD is a parallelogram and X is the mid-point of BC. The line AX produced meets DC produced at Q. The parallelogram AQP B is completed.

Prove that :

(i) $\triangle ABX \cong \triangle QCX$.

(ii) $DC = CQ = QP$.



Sol. **Given :** ABCD is a parallelogram. X is mid-point of BC.

AX is joined and produced to meet DC produced at Q. From B, BP is drawn parallel to AQ so that AQP B is a parallelogram.

To prove : (i) $\triangle ABX \cong \triangle QCX$.

(ii) $DC = CQ = QP$.

Proof : (i) In $\triangle ABX$ and $\triangle QCX$,

$$XB = XC \quad (\because X \text{ is mid-point of } BC)$$

$$\angle AXB = \angle CXQ$$

(Vertically opposite angles)

$$\angle BAX = \angle XQC \quad (\text{Alternate angles})$$

$$\therefore \triangle ABX \cong \triangle QCX$$

(AAS axiom of congruency)

Hence proved.

(ii) In parallelogram ABCD,

$$AB = DC \quad \dots(i)$$

(Opposite sides of a parallelogram)

Similarly, in parallelogram AQP B

$$AB = QP \quad \dots(ii)$$

\therefore From (i) and (ii)

$$DC = QP \quad \dots(iii)$$

In $\triangle BCP$,

X is mid-point of BC and $AQ \parallel BP$.

\therefore Q is mid-point of CP.

$$\Rightarrow CQ = QP \quad \dots(ii)$$

From (iii) and (iv)

$$DC = QP = CQ$$

$$\text{or } DC = CQ = QP$$


Hence proved.

Q. 18. In the adjoining figure, ABCD is a parallelogram.

Line segments AX and CY bisect $\angle A$ and $\angle C$ respectively.

Prove that :

- (i) $\triangle ADX \cong \triangle CBY$
- (ii) $AX = CY$
- (iii) $AX \parallel CY$
- (iv) AYCX is a parallelogram.



Sol. **Given :** ABCD is a parallelogram.

Line segments AX and CY bisect $\angle A$ and $\angle C$ respectively.

To prove : (i) $\triangle ADX \cong \triangle CBY$

(ii) $AX = CY$ (iii) $AX \parallel CY$

(iv) AYCX is a parallelogram.

Proof : (i) In $\triangle ADX$ and $\triangle CBY$,

$$AD = BC$$

(Opposite sides of a parallelogram)

$$\angle D = \angle B$$

(Opposite angles of the parallelogram)

$$\angle DAX = \angle BCY$$

{half of equal angles A and C}

$$\therefore \triangle ADX \cong \triangle CBY$$

(AAS axiom of congruency)

$$(ii) \therefore AX = CY \quad (\text{C.P.C.T.})$$

$$(iii) \angle 1 = \angle 2 \quad (\text{Half of equal angles})$$

$$\text{But } \angle 2 = \angle 3 \quad (\text{Alternate angles})$$

$$\therefore \angle 1 = \angle 3$$

But these are corresponding angles.


$$\therefore AX \parallel CY$$

$$(iv) \therefore AX = CY \text{ and } AX \parallel CY$$

\therefore AYCX is a parallelogram.

Hence proved.

Q. 19. In the given figure, ABCD is a parallelogram and X, Y are points on diagonal BD such that $DX = BY$. Prove that CXAY is a parallelogram.



Sol. **Given :** ABCD is a parallelogram. X and Y are points on diagonal BD such that $DX = BY$.

To prove : CXAY is a parallelogram.

Const : Join AC meeting BD at O.

Proof : \because AC and BD are the diagonals of the parallelogram ABCD.

\therefore AC and BD bisect each other at O.

$$\therefore AO = OC \text{ and } BO = OD$$

But $DX = BY$ (Given)

$$\therefore DO - DX = OB - BY$$


$$\Rightarrow OX = OY$$

Now in quadrilateral CXAY, diagonals AC and XY bisect each other at O.

\therefore CXAY is a parallelogram.

Hence proved.

Q. 20. Show that the bisectors of the angles of a parallelogram enclose a rectangle.



Sol. **Given :** ABCD is a parallelogram.

Bisectors of $\angle A$ and $\angle B$ meet at S and bisectors of $\angle C$ and $\angle D$ meet at Q.

To prove : PQRS is a rectangle.

Proof : $\because \angle A + \angle B = 180^\circ$

$$\therefore \frac{1}{2} \angle A + \frac{1}{2} \angle B = 90^\circ$$

$$\Rightarrow \angle SAB = \angle SBA = 90^\circ$$

\therefore In $\triangle ASB$,

$$\angle ASB = 90^\circ$$

Similarly we can prove that

$$\angle CQD = 90^\circ$$

$$\text{Again } \angle A + \angle D = 180^\circ$$

(Co. interior angles)

$$\therefore \frac{1}{2} \angle A + \frac{1}{2} \angle D = 90^\circ$$

$$\Rightarrow \angle PAD = \angle PDA = 90^\circ$$

$$\therefore \angle APD = 90^\circ$$

$$\text{But } \angle SPQ = \angle APD$$

(Vertically opposite angles)

$$\therefore \angle SPQ = 90^\circ$$

\therefore Similarly we can prove that


$$\angle SRQ = 90^\circ$$

\therefore In quadrilateral PQRS, its each angle is of 90°

\therefore PQRS is a rectangle.

Hence proved.

- Q. 21.** If a diagonal of a parallelogram bisects one of the angles of the parallelogram, prove that it also bisects the second angle and then the two diagonals are perpendicular to each other.



Sol. Given : In parallelogram ABCD, diagonal AC bisects $\angle A$. BD is joined meeting AC at O.

To prove : (i) AC bisects $\angle C$.

(ii) Diagonal AC and BD are perpendicular to each other.

Proof : $\therefore AB \parallel DC$

$$\therefore \angle 1 = \angle 4 \quad \text{and} \quad \angle 2 = \angle 3$$

(Alternate angles)

$$\text{But } \angle 1 = \angle 2$$

(Given)

$$\therefore \angle 3 = \angle 4$$

Hence AC bisects $\angle C$ also.

Similarly we can prove that diagonal BD will also bisect the $\angle B$ and $\angle D$.


\therefore ABCD is a rhombus.

But diagonals of a rhombus bisect each other at right angles.

\therefore AC and BD are perpendicular to each other.

Hence proved.

- Q. 22.** In the given figure, ABCD is a parallelogram and E is the mid-point of BC. If DE and AB produced meet at F, prove that $AF = 2 AB$.



Sol. Given : ABCD is a parallelogram. E is mid-point of BC. DE and AB are produced to meet at F.

To prove : $AF = 2 AB$.

Proof : In $\triangle DEC$ and $\triangle FEB$

$$CE = EB \quad (\because E \text{ is mid-point of } BC)$$

$$\angle DEC = \angle BEF$$

(Vertically opposite angles)

$$\angle DCE = \angle EBF \quad (\text{Alternate angles})$$

$$\therefore \triangle DEC \cong \triangle FEB$$

(AAS axiom of congruency)

$$\therefore CD = BF \quad (\text{C.P.C.T.})$$

$$\text{But } AB = CD$$

(Opposite sides of a parallelogram)

$$\begin{aligned} \therefore AB &= BF \\ AF &= AB + BF \\ &= AB + AB = 2 AB \end{aligned}$$


Hence proved.

EXERCISE 15 (B)

Q. 1. Construct a quadrilateral PQRS in which $PQ = 4$ cm, $\angle P = 90^\circ$, $QR = 4.3$ cm, $RS = 3.6$ cm and $SP = 3.2$ cm.

Sol. Steps of Construction :

(i) Draw a line segment $PQ = 4$ cm.



(ii) At P, draw a ray AX making an angle of 90° .

(iii) From PX cut off $PS = 3.2$ cm.

(iv) With centre Q and radius 4.3 cm draw an arc.

(v) With centre S and radius 3.6 cm draw another arc which intersects the first arc at R.

(vi) Join QR and SR.

Then PQRS is the required quadrilateral.

Q. 2. Construct a quadrilateral ABCD in which $AB = 4.5$ cm, $BC = 5.2$ cm, $CD = 5$ cm, $DA = 4.7$ cm and $\angle ABC = 75^\circ$.


Sol. Steps of Construction :

(i) Draw a line segment $AB = 4.5$ cm.

(ii) At B, draw an arc BX making an angle of 75° .

(iii) From BX, cut off $BC = 5.2$ cm.

(iv) With centre C and radius 5 cm draw an arc.



(v) With centre A and radius 4.7 cm draw another arc which intersects the first arc at D.


(vi) Join AD and CD.

Then ABCD is the required quadrilateral.

Q. 3. Construct a quadrilateral ABCD in which $AB = CD = 5.1$ cm, $BC = 4.7$ cm, $DA = 4.2$ cm and $\angle BCD = 60^\circ$.

Sol. Steps of Construction :

(i) Draw a line segment $BC = 4.7$ cm.



(ii) At C, draw an arc CX making an angle of 60° and cut off $CD = 5.1$ cm.

(iii) With centre D and radius 4.2 cm, draw an arc.

(iv) With centre B and radius 5.1 cm, draw another arc intersecting the first arc at A.

(v) Join AB and AD.


Then ABCD is the required quadrilateral.

Q. 4. Construct a quadrilateral UVWZ in which $UV = 6$ cm, $VW = 5.3$ cm, $UZ = 5$ cm, $\angle U = 60^\circ$ and $\angle V = 75^\circ$.

Sol. Steps of Construction :

(i) Draw a line segment $UV = 6$ cm.

(ii) At U, draw a ray UX making an angle of 60° .



(iii) At V, draw another ray VY making an angle of 75° .

(iv) From UX, cut $UZ = 5$ cm and from VY, cut off $VW = 5.3$ cm.

(v) Join WZ.

Then UVWZ is the required quadrilateral.


Q. 5. Construct a quadrilateral PQRS in which $PQ = 4.5$ cm, $QR = 5.6$ cm, $RS = 4.1$ cm, $\angle Q = 60^\circ$ and $\angle R = 120^\circ$.

Sol. Steps of Construction :

(i) Draw a line $QR = 5.6$ cm.

(ii) At Q, draw a ray QX making an angle of 60° .

(iii) At R, draw another ray RY making an angle of 120° .



(iv) Cut off from QX, $QP = 4.5$ cm and from RY, $RS = 4.1$ cm.


(v) Join PS.

Then PQRS is the required quadrilateral.

Q. 6. Draw a quadrilateral ABCD in which $AB = AD = 5.3$ cm, $BC = CD = 5$ cm and diagonal $BD = 6.4$ cm.

Sol. Steps of Construction :

(i) Draw a line segment $BD = 6.4$ cm.



(ii) With B and D as centres and radius 5.3 cm each draw two arcs intersecting each other at A above BD.

(iii) Join AB and AD.

(iv) Again with centre B and D and radius 5 cm, draw two arcs intersecting each other below the line segment BD.


(v) Join BC and CD.

Then ABCD is the required quadrilateral.

Q. 7. Construct a quadrilateral XYZT in which $XY = XZ = 5.7$ cm, $YZ = 4.5$ cm, $ZX = 4.6$ cm and $XT = 5$ cm.

Sol. Steps of Construction :

(i) Draw a line segment $XZ = 5.7$ cm.



(ii) With centre X and radius 5 cm and with centre Z and radius 4.6 cm, draw two arcs which intersect each other at T above the line segment XZ.

(iii) Join TX and TZ.

(iv) Again with centre X and radius 5.7 cm and with centre Z and radius 4.5 cm, draw two arcs intersecting each other at Y below the line segment XZ.


(v) Join XY and ZY.

Then XYZT is the required quadrilateral.

Q. 8. Construct a parallelogram ABCD in which $BC = 6$ cm, $CD = 4$ cm and $\angle C = 60^\circ$.

Sol. Steps of Construction :

(i) Draw a line segment $BC = 6$ cm.



(ii) At C, draw a ray CX making an angle of 60° and cut off $CD = 4$ cm.

(iii) With centre B and radius 4 cm and with centre D and radius 5 cm, draw arc intersecting each other at A.


(iv) Join AB and AD.

Then ABCD is the required parallelogram.

Q. 9. Construct a parallelogram PQRS in which $QR = 4.7$ cm, $\angle Q = 120^\circ$ and $PQ = 2.8$ cm.

Sol. Steps of Construction :

(i) Draw a line segment $PQ = 2.8$ cm.



(ii) At Q, draw a ray QX making an angle of 120° and cut off $QR = 4.7$ cm.

(iii) With centre P and radius 4.7 cm and with centre R and radius 2.8 cm draw arcs intersecting each other at S.


(iv) Join SP and SR.

Then PQRS is the required parallelogram.

Q. 10. Construct a parallelogram ABCD in which $BC = 5$ cm, $CD = 3$ cm and diagonal $BD = 6$ cm.

Sol. Steps of Construction :

(i) Draw a line segment $BC = 5$ cm.



(ii) With B as centre and radius 6 cm, draw an arc.

(iii) With C as centre and radius 3 cm, draw another arc intersecting the first arc at D.

(iv) Join BD and CD.

(v) Again with centre B and radius 3 cm draw an arc and with centre D and radius 5 cm draw another arc intersecting each other at A.

(vi) Join AB and AD.

Then ABCD is the required parallelogram.

Q. 11. Construct a parallelogram ABCD in which $AB = 4.8$ cm, $BC = 3.5$ cm and diagonal $AC = 5.4$ cm.


Sol. Steps of Construction :

(i) Draw a line segment $AB = 4.5$ cm.

(ii) With centre A and radius 5.4 cm draw an arc and with centre B and radius 3.5 cm, draw another arc intersecting the first arc at C.

(iii) Join AC and BC.

(iv) Again with centre A and radius 3.5 cm and with centre C and radius 4.8 cm, draw two arcs intersecting each other at D.



(v) Join CD and AD.


Then ABCD is the required parallelogram.

Q. 12. Construct a parallelogram ABCD in which diagonal $AC = 5.6$ cm, diagonal $BD = 6.2$ cm and angle between them is 60° .

Sol. Steps of Construction :

(i) Draw a line segment $AC = 5.6$ cm.

(ii) Bisect AC at O.



(iii) At O, draw a line XY making an angle of 60° and cut off $OB = OD$

$$= \frac{6.2}{2} = 3.1 \text{ cm.}$$


(iv) Join AB, BC, CD and DA.

Then ABCD is the required parallelogram.

Q. 13. Construct a rectangle whose adjacent sides are 4.7 cm and 3.2 cm.

Sol. Steps of Construction :

(i) Draw a line segment AB = 4.7 cm.



(ii) At A and B, draw rays AX and BY making an angle of 90° each and cut off $AD = BC = 3.2$ cm.


(iii) Join CD.

Then ABCD is the required rectangle.

Q. 14. Construct a rectangle ABCD in which $AB = 4.5$ cm and diagonal $AC = 6$ cm.

Sol. Steps of Construction :

(i) Draw a line segment AB = 4.5 cm.



(ii) At B, draw a ray BX making an angle of 90° .

(iii) With centre A and radius 6 cm draw an arc which intersects the ray BX at C.

(iv) Join AC.

(v) With centre A and radius BC and with centre C and radius 4.5 cm, draw arcs intersecting each other at D.


(vi) Join AD and CD.

Then ABCD is the required rectangle.

Q. 15. Construct a rectangle ABCD in which diagonal $AC = 6.3$ cm and the angle between the two diagonals is 50° .

Sol. Steps of Construction :

(i) Draw a line segment AC = 6.3 cm.



(ii) Bisect AC at O.

(iii) At O, draw a line XY making an angle of 50° and produce it both sides.

(iv) Cut off $OB = OD = \frac{6.3}{2}$ cm = 3.15 cm

(\because Diagonal of a rectangle are equal)


(v) Join AB, BC, CD and DA.

Then ABCD is the required rectangle.

Q. 16. Construct a square one of whose diagonals measures 6 cm.

Sol. Steps of Construction :

(i) Draw a line segment $AC = 6$ cm.



(ii) Draw the perpendicular bisector XY of AC cutting AC at O .

(iii) From O , cut off $OB = OD = \frac{6}{2} = 3$ cm

(\because Diagonals of a square are equal and bisect each other at right angles)


(iv) Join AB, BC, CD and DA .

Then $ABCD$ is the required square.

Q. 17. Construct a rhombus $ABCD$ in which $AB = 4.3$ cm and $\angle A = 45^\circ$.

Sol. Steps of Construction :

(i) Draw a line segment $AB = 4.3$ cm.



(ii) At A , draw a ray AX making an angle of 45° .

(iii) From AX cut off $AD = 4.3$ cm.

(iv) With centre B and D and radius equal to 4.3 cm, draw arcs intersecting each other at C .


(v) Join BC and DC .

Then $ABCD$ is a rhombus.

Q. 18. Construct a rhombus $ABCD$ in which $AB = 5$ cm and diagonal $AC = 6.8$ cm.

Sol. Steps of Construction :

(i) Draw a line segment $AC = 6.8$ cm.



(ii) With centre A and C and radius 5 cm, draw arcs which intersect each other at B and D i.e. on both sides of AC .

(iii) Join AB, BC, CD and DA .

Then $ABCD$ is the required rhombus.

Q. 19. Construct a rhombus $ABCD$ in which diagonal $AC = 5.8$ cm and diagonal $BD = 6.4$ cm.

Sol. Steps of Construction :


(i) Draw a line segment $AC = 5.8$ cm.

(ii) Draw the perpendicular bisector of AC intersecting it at O .

(iii) From O , cut off $OB = OD$

$$= \frac{6.4}{2} = 3.2 \text{ cm.}$$

(iv) Join AB, BC, CD and DA .




Then ABCD is the required rhombus.

Q. 20. Construct a square one of whose diagonals is 5.6 cm.

Sol. Steps of Construction :

- (i) Draw a line segment $AC = 5.6$ cm.
- (ii) Draw its right bisector XY intersecting it at O .



- (iii) From O , cut off $OB = OD$

$$= \frac{5.6}{2} = 2.8 \text{ cm.}$$

- (iv) Join AB, BC, CD and DA .

Then ABCD is the required square.