





Mathematics & Statistics

Arts & Science Part 1

STANDARD XII



The Coordination Committee formed by GR No.Abhyas - 2116/(Pra.Kra 43/16) SD - 4
Dated 25.4.2016 has given approved to prescribe this textbook in its meeting held on
30.01.2020 and it has been decided to implement it from the educational year 2020 - 21.

Mathematics and Statistics (Arts and Science)

Part - I

STANDARD - XII



Maharashtra State Bureau of Textbook Production and Curriculum Research, Pune. - 411 004



Z4R8V9

Download DIKSHA App on your smartphone. If you scan the Q.R.Code on this page of your textbook, you will be able to access full text and the audio-visual study material relevant to each lesson provided as teaching and learning aids.

First Edition : 2020 © Maharashtra State Bureau of Textbook Production and Curriculum Research, Pune- 411 004.

Maharashtra State Bureau of Textbook Production and Curriculum Research reserves all rights relating to the book. No part of this book should be reproduced without the written permission of the Director, Maharashtra State Bureau of Textbook Production and Curriculum Research, Pune.

**Mathematics and Statistics
(Arts and Science)
Committee Members**

Dr. Mangla Narlikar	(Chairman)
Dr. Sharad Gore	Member
Shri. Prahallad Chippalagatti	Member
Shri. Prasad Kunte	Member
Shri. Sujit Shinde	Member
Smt. Prajakti Gokhale	Member
Shri. Ramakant Sarode	Member
Smt. Ujjwala Godbole	(Member -Secretary)

Co-ordinator

Ujjwala Shrikant Godbole
I/C Special Officer for Mathematics

**Cover
&
Computer Drawings**

Shri. Sandip Koli, Artist, Mumbai

**Mathematics and Statistics
(Arts and Science)
Study Group Members**

Dr. Ishwor Patil	Dr. Pradeep Mugale
Dr. Pradnyankumar Bhojekar	Shri. Uday Mahindrakar
Shri. Prafullachandra Pawar	Shri. Balkrishna Mapari
Shri. Sachin Batwal	Shri. Devanand Bagul
Shri. Swapnil Shinde	Smt. Swati Pawar
Shri. Pradipkumar Bhavsar	Smt. Mahadevi Mane
Shri. Amit Totade	Shri. Dilip Panchamia
Smt. Gouri Prachand	Shri. Pramod Deshpande
Shri. Sharadchandra Walagade	Shri. Milind Patil
Shri. Dhananjay Panhalkar	Smt. Supriya Abhyankar
Smt. Deepti Kharas	Smt. Nileema Khaladkor

Production

Sachchitanand Aphale
Chief Production Officer

Sanjay Kamble
Production Officer

Prashant Harne
Asst. Production Officer

Paper

70 GSM Cream wove

Print Order No.

N/PB/2020-21/50000

Printer


DIGITAL PRINT SYSTEM & SOLUTIONS PVT. LTD.,
AHMEDNAGAR

Typesetter

M/s. Shree Graphic Centre,
Mumbai

Publisher

Vivek Uttam Gosavi, Controller
Maharashtra State Textbook Bureau,
Prabhadevi Mumbai- 400 025



The Constitution of India

Preamble

WE, THE PEOPLE OF INDIA, having solemnly resolved to constitute India into a SOVEREIGN SOCIALIST SECULAR DEMOCRATIC REPUBLIC and to secure to all its citizens:

JUSTICE, social, economic and political;

LIBERTY of thought, expression, belief, faith and worship;

EQUALITY of status and of opportunity; and to promote among them all

FRATERNITY assuring the dignity of the individual and the unity and integrity of the Nation;

IN OUR CONSTITUENT ASSEMBLY this twenty-sixth day of November, 1949, do HEREBY ADOPT, ENACT AND GIVE TO OURSELVES THIS CONSTITUTION.

NATIONAL ANTHEM

Jana-gana-mana-adhināyaka jaya hē
Bhārata-bhāgya-vidhātā,

Panjāba-Sindhu-Gujarāta-Marāthā
Drāvida-Utkala-Banga

Vindhya-Himāchala-Yamunā-Gangā
uchchala-jaladhi-taranga

Tava subha nāmē jāgē, tava subha āsisa māgē,
gāhē tava jaya-gāthā,

Jana-gana-mangala-dāyaka jaya hē
Bhārata-bhāgya-vidhātā,

Jaya hē, Jaya hē, Jaya hē,
Jaya jaya jaya, jaya hē.

PLEDGE

India is my country. All Indians
are my brothers and sisters.

I love my country, and I am proud
of its rich and varied heritage. I shall
always strive to be worthy of it.

I shall give my parents, teachers
and all elders respect, and treat
everyone with courtesy.

To my country and my people,
I pledge my devotion. In their
well-being and prosperity alone lies
my happiness.

PREFACE

Dear Students,

Welcome to Standard XII, an important milestone in your life.

Standard XII or Higher Secondary School Certificate opens the doors of higher education. Alternatively, you can pursue other career paths like joining the workforce. Either way, you will find that mathematics education helps you considerably. Learning mathematics enables you to think logically, consistently, and rationally. The curriculum for Standard XII Mathematics and Statistics for Science and Arts students has been designed and developed keeping both of these possibilities in mind.

The curriculum of Mathematics and Statistics for Standard XII for Science and Arts students is divided in two parts. Part I deals with topics like Mathematical Logic, Matrices, Vectors and Introduction to three dimensional geometry. Part II deals with Differentiation, Integration and their applications, Introduction to random variables and statistical methods.

The new text books have three types of exercises for focussed and comprehensive practice. First, there are exercises on every important topic. Second, there are comprehensive exercises at the end of all chapters. Third, every chapter includes activities that students must attempt after discussion with classmates and teachers. Additional information has been provided on the E-balbharati website (www.ebalbharati.in).

We are living in the age of Internet. You can make use of modern technology with the help of the Q.R. code given on the title page. The Q.R. code will take you to links that provide additional useful information. Your learning will be fruitful if you balance between reading the text books and solving exercises. Solving more problems will make you more confident and efficient.

The text books are prepared by a subject committee and a study group. The books (Paper I and Paper II) are reviewed by experienced teachers and eminent scholars. The Bureau would like to thank all of them for their valuable contribution in the form of creative writing, constructive and useful suggestions for making the text books valuable. The Bureau hopes and wishes that the text books are very useful and well received by students, teachers and parents.

Students, you are now ready to study. All the best wishes for a happy learning experience and a well deserved success. Enjoy learning and be successful.



(Vivek Gosavi)
Director

Pune

Date: 21 February 2020

Bharatiya Saur: 2 Phalguna 1941

Maharashtra State Bureau of Textbook
Production and Curriculum Research, Pune.

Mathematics and Statistics Std. XII Part - I
Competency Statement

Sr.No.	Area	Topic	Competency Statement
1.	Mathematical Logic	Mathematical Logic	<p>The student will be able to</p> <ul style="list-style-type: none"> • Identify statement in logic and truth value of it. • Combine two or more statements • Construct the truth table and examine logical equivalence of statement patterns • Find dual and negation of statement pattern • Study the applications of logic to switching circuits.
2.	Matrices	Matrices	<ul style="list-style-type: none"> • Identify orders and types of matrices • Perform basic algebraic operations on matrices. • Find the inverse of a matrix using elementary transformation and adjoint method • Solve the system of linear equations using matrices.
3.	Trigonometric Equations	Trigonometric Equations Solution of a triangle Invers trigonometric function	<ul style="list-style-type: none"> • Understand and write trigonometric equation • Find the principal and general solution of a trigonometric equation. • Solve triangle by using sine rule, cosine rule and projection rule and find area of a triangle. • Understand inverse trigonometric functions with domain and range.
4.	Pair of straight lines	Pair of straight lines	<ul style="list-style-type: none"> • Write and interpret the combined equation of two straight lines in plane. • Find the point of intersection of two lines and calculate the acute angle between them • Study the general second degree equation in x and y with reference to homogeneous part of it

5.	Vectors	Vectors	<ul style="list-style-type: none"> • Understand scalars and vectors and algebra of vectors. • Write vectors of 2 or 3 dimensions, understand the scalar and vector products • Study applications of vectors to area of triangle, work done by a force, moment of a force. • Interpret scalar triple product and its applications.
6.	Line and Plane	Line and Plane	<ul style="list-style-type: none"> • Find different forms of equation of line • Find angle between two intersecting planes • Find the angle between a line and a plane • Find condition for perpendicularity and parallelness of planes • Calculate distance of a point from a plane • Find equation of a plane in different forms • Find angle between two intersecting planes • Find the angle between a line and a plane
7.	Linear programming Problem	Linear programming Problem	<ul style="list-style-type: none"> • Understand linear equations in one and two variables. • Find graphical solution of linear inequation. • Understand meaning and formulation of L.P.P. • Find solution of L.P.P. by graphical methods.

INDEX

Sr.No.	Chapter	Page No
1.	Mathematical Logic	1
2.	Matrics	36
3.	Trigonometric Functions	65
4.	Pair of Straight Lines	112
5.	Vectors	133
6.	Line and Plane	194
7.	Linear Programming	227
	Answers	246

1

Mathematical Logic



Let's study.

1. Statement and its truth value.
2. Logical connective, compound statements.
3. Truth tables negation of statements and compound statements.
4. Statement pattern, logical equivalence.
5. Tautology, contradiction and contingency.
6. Quantifiers and quantified statements, Duality.
7. Application of logic to switching circuits, switching table.



Let's recall.

1.1.1 Introduction :

Mathematics is a logical subject and tries to be exact. For exactness, it requires proofs which depend upon proper reasoning. Reasoning requires logic. The word Logic is derived from the Greek word "LOGOS" which means reason. Therefore logic deals with the method of reasoning. In ancient Greece the great philosopher and thinker Aristotle started study of Logic systematically. In mathematics Logic has been developed by English Philosopher and mathematician George Boole (2 November 1815 - 8 December 1864)

Language is the medium of communication of our thoughts. For communication we use sentences. In logic, we use the statements which are special sentences.

1.1.2 Statement :

A statement is a declarative (assertive) sentence which is either true or false, but not both simultaneously. Statements are denoted by p, q, r, \dots

1.1.3 Truth value of a statement :

Each statement is either true or false. If a statement is true then its truth value is 'T' and if the statement is false then its truth value is F .

Illustrations :**1) Following sentences are statements.**

- i) Sun rises in the East.
- ii) $5 \times 2 = 11$
- iii) Every triangle has three sides.

- iv) Mumbai is the capital of Maharashtra.
- v) Every equilateral triangle is an equiangular triangle.
- vi) A natural number is an integer.

2) Following sentences are not statements.

- i) Please, give your Pen.
- ii) What is your name ?
- iii) What a beautiful place it is !
- iv) How are you ?
- v) Do you like to play tennis ?
- vi) Open the window.
- vii) Let us go for tea
- viii) Sit dows.

Note : Interrogative, exclamatory, command, order, request, suggestion are not statements.

3) Consider the following.

- i) $\frac{3x}{2} - 9 = 0$
- ii) He is tall.
- iii) Mathematics is an interesting subject.
- iv) It is black in colour.

Let us analyse these statements.

- i) For $x = 6$ it is true but for other than 6 it is not true.
- ii) Here, we cannot determine the truth value.

For iii) & iv) the truth value varies from person to person. In all the above sentences, the truth value depends upon the situation. Such sentences are called as open sentences. Open sentence is not a statement.



Solved examples

Q.1. Which of the following sentences are statements in logic ? Write down the truth values of the statements.

- i) $6 \times 4 = 25$
- ii) $x + 6 = 9$
- iii) What are you doing ?
- iv) The quadratic equation $x^2 - 5x + 6 = 0$ has 2 real roots.
- v) Please, sit down
- vi) The Moon revolves around the earth.
- vii) Every real number is a complex number.
- viii) He is honest.
- ix) The square of a prime number is a prime number.

Solution :

- i) It is a statement which is false, hence its truth value is F .
- ii) It is an open sentence hence it is not a statement.
- iii) It is an interrogative hence it is not a statement.
- iv) It is a statement which is true hence its truth value is T .
- v) It is a request hence it is not a statement.
- vi) It is a statement which is true, hence its truth value is T .
- vii) It is a statement which is true, hence its truth value is T .
- viii) It is open sentence, hence it is not a statement.
- ix) It is a statement which is false, hence its truth value is F .

1.1.4 Logical connectives, simple and compound statements :

The words or phrases which are used to connect two statements are called logical connectives. We will study the connectives 'and', 'or', 'if then', 'if and only if', 'not'.

Simple and Compound Statements : A statement which cannot be split further into two or more statements is called a simple statement. If a statement is the combination of two or more simple statements, then it is called a compound statement.

"3 is a prime and 4 is an even number", is a compound statement.

"3 and 5 are twin primes", is a simple statement.

We describe some connectives.

1) Conjunction : If two statements are combined using the connective 'and' then it is called as a conjunction. In other words if p , q are two statements then ' p and q ' is called as conjunction. It is denoted by ' $p \wedge q$ ' and it is read as ' p conjunction q ' or ' p and q '. The conjunction $p \wedge q$ is said to be true if and only if both p and q are true.

Truth table for conjunction

p	q	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

Table 1.1

2) Disjunction : If two statements are combined by using the logical connective 'or' then it is called as a disjunction. In other words if p , q are two statements then ' p or q ' is called as disjunction. It is denoted by ' $p \vee q$ ' and it is read as ' p or q ' or ' p disjunction q '.

Truth table for disjunction

p	q	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

The disjunction $p \vee q$ is false if and only if both p and q are false.

Table 1.2

- 3) **Conditional (Implication) :** If two statements are combined by using the connective. 'if ... then', then it is called as conditional or implication. In other words if p, q are two statements then 'if p then q ' is called as conditional. It is denoted by $p \rightarrow q$ or $p \Rightarrow q$ and it is read as 'p implies q' or 'if p then q '.

Truth table for conditional.

p	q	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

The conditional statement $p \rightarrow q$ is False only if p is true and q is false. Otherwise it is true. Here p is called hypothesis or antecedent and q is called conclusion or consequence.

Table 1.3

Note : The following are also conditional statement $p \rightarrow q$

- i) p is sufficient for q
- ii) q is necessary for p
- iii) p implies q
- iv) q follows from p
- v) p only if q .

- 4) **Biconditional (Double implication) :**

If two statements are combined using the logical connective 'if and only if' then it is called as biconditional. In other words if p, q are two statements then ' p if and only if q ' is called as biconditional. It is denoted by ' $p \leftrightarrow q$ ' or $p \Leftrightarrow q$. It is read as ' p biconditional q ' or ' p if and only if q '.

Truth table for biconditional.

p	q	$p \leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

Biconditional statement $p \leftrightarrow q$ is true if p and q have same truth values. Otherwise it is False.

Table 1.4

- 5) **Negation of a statement :** For any given statement p , there is another statement which is defined to be true when p is false, and false when p is true, is called the negation of p and is denoted by $\sim p$.

Truth table for negation.

p	$\sim p$
T	F
F	T

Table 1.5

Note : Negation of negation of a statement is the statement itself. That is, $\sim(\sim p) = p$.



Solved examples

Ex.1: Express the following compound statements symbolically without examining the truth values.

- i) 2 is an even number and 25 is a perfect square.
- ii) A school is open or there is a holiday.
- iii) Delhi is in India but Dhaka is not in Srilanks.
- iv) $3 + 8 \geq 12$ if and only if $5 \times 4 \leq 25$.

Solution :

- i) Let p : 2 is an even number
 q : 25 is a perfect square.
 The symbolic form is $p \wedge q$.
- ii) Let p : The school is open
 q : There is a holiday
 The symbolic form is $p \vee q$
- iii) Let p : Delhi is in India
 q : Dhaka is in Srilanka
 The symbolic form is $p \wedge \sim q$.
- iv) Let p : $3 + 8 \geq 12$; q : $5 \times 4 \leq 25$
 The symbolic form is $p \leftrightarrow q$

Ex.2. Write the truth values of the following statements.

- i) 3 is a prime number and 4 is a rational number.
- ii) All flowers are red or all cows are black.
- iii) If Mumbai is in Maharashtra then Delhi is the capital of India.
- iv) Milk is white if and only if the Sun rises in the West.

Solution :

- i) Let p : 3 is a prime number
 q : 4 is a rational number.
 Truth values of p and q are T and T respectively.
 The given statement in symbolic form is $p \wedge q$.
 The truth value of given statement is T.
- ii) Let p : All flowers are red ; q : All cows are black.
 Truth values of p and q are F and F respectively.
 The given statement in the symbolic form is $p \vee q$
 $\therefore p \vee q \equiv F \vee F$ is F
 \therefore Truth value of given statement is F.
- iii) Let p : Mumbai is in Maharashtra
 q : Delhi is capital of India
 Truth values of p and q are T and T respectively.
 The given statement in symbolic form is $p \rightarrow q$

$$\therefore p \rightarrow q \equiv T \rightarrow T \text{ is } T$$

\therefore Truth value of given statement is T

iv) Let p : Milk is white; q : Sun rises in the West.

Truth values of p and q are T and F respectively.

The given statement in symbolic form is $p \leftrightarrow q$

$$\therefore p \leftrightarrow q \equiv T \leftrightarrow F \text{ is } F$$

\therefore Truth value of given statement is F

Ex.3 : If statements p, q are true and r, s are false, determine the truth values of the following.

i) $\sim p \wedge (q \vee \sim r)$ ii) $(p \wedge \sim r) \wedge (\sim q \vee s)$

iii) $\sim(p \rightarrow q) \leftrightarrow (r \wedge s)$ iv) $(\sim p \rightarrow q) \wedge (r \leftrightarrow s)$

Solution :

i) $\sim p \wedge (q \vee \sim r) \equiv \sim T \wedge (T \vee \sim F) \equiv F \wedge (T \vee T) \equiv F \wedge T \equiv F$

Hence truth value is F .

ii) $(p \wedge \sim r) \wedge (\sim q \vee s) \equiv (T \wedge \sim F) \wedge (\sim T \vee F) \equiv (T \wedge T) \wedge (F \vee F) \equiv T \wedge F \equiv F$.

Hence truth value is F .

iii) $[\sim(p \rightarrow q)] \leftrightarrow (r \wedge s) \equiv [\sim(T \rightarrow T)] \leftrightarrow (F \wedge F) \equiv (\sim T) \leftrightarrow (F) \equiv F \leftrightarrow F \equiv T$.

Hence truth value is T

iv) $(\sim p \rightarrow q) \wedge (r \leftrightarrow s) \equiv (\sim T \rightarrow T) \wedge (F \leftrightarrow F) \equiv (F \rightarrow T) \wedge T \equiv T \wedge T \equiv T$.

Hence truth value is T .

Ex.4. Write the negations of the following.

i) Price increases

ii) $0! \neq 1$

iii) $5 + 4 = 9$

Solution :

i) Price does not increase

ii) $0! = 1$

iii) $5 + 4 \neq 9$



Exercise 1.1

Q.1. State which of the following are statements. Justify. In case of statement, state its truth value.

i) $5 + 4 = 13$.

ii) $x - 3 = 14$.

iii) Close the door.

iv) Zero is a complex number.

v) Please get me breakfast.

vi) Congruent triangles are similar.

vii) $x^2 = x$.

- viii) A quadratic equation cannot have more than two roots.
- ix) Do you like Mathematics ?
- x) The sun sets in the west
- xi) All real numbers are whole numbers
- xii) Can you speak in Marathi ?
- xiii) $x^2 - 6x - 7 = 0$, when $x = 7$
- xiv) The sum of cuberoots of unity is zero.
- xv) It rains heavily.

Q.2. Write the following compound statements symbolically.

- i) Nagpur is in Maharashtra and Chennai is in Tamilnadu
- ii) Triangle is equilateral or isosceles.
- iii) The angle is right angle if and only if it is of measure 90° .
- iv) Angle is neither acute nor obtuse.
- v) If ΔABC is right angled at B, then $m \angle A + m \angle C = 90^\circ$
- iv) Hima Das wins gold medal if and only if she runs fast.
- vii) x is not irrational number but is a square of an integer.

Q.3. Write the truth values of the following.

- i) 4 is odd or 1 is prime.
- ii) 64 is a perfect square and 46 is a prime number.
- iii) 5 is a prime number and 7 divides 94.
- iv) It is not true that $5-3i$ is a real number.
- v) If $3 \times 5 = 8$ then $3 + 5 = 15$.
- vi) Milk is white if and only if sky is blue.
- vii) 24 is a composite number or 17 is a prime number.

Q.4. If the statements p, q are true statements and r, s are false statements then determine the truth values of the following.

- | | |
|---|--|
| i) $p \vee (q \wedge r)$ | ii) $(p \rightarrow q) \vee (r \rightarrow s)$ |
| iii) $(q \wedge r) \vee (\sim p \wedge s)$ | iv) $(p \rightarrow q) \wedge \sim r$ |
| v) $(\sim r \leftrightarrow p) \rightarrow \sim q$ | vi) $[\sim p \wedge (\sim q \wedge r)] \vee [(q \wedge r) \vee (p \wedge r)]$ |
| vii) $[(\sim p \wedge q) \wedge \sim r] \vee [(q \rightarrow p) \rightarrow (\sim s \vee r)]$ | viii) $\sim[(\sim p \wedge r) \vee (s \rightarrow \sim q)] \leftrightarrow (p \wedge r)$ |

Q.5. Write the negations of the following.

- i) Tirupati is in Andhra Pradesh
- ii) 3 is not a root of the equation $x^2 + 3x - 18 = 0$
- iii) $\sqrt{2}$ is a rational number.
- iv) Polygon ABCDE is a pentagon.
- v) $7 + 3 > 5$

1.2 STATEMENT PATTERN, LOGICAL EQUIVALENCE, TAUTOLOGY, CONTRADICTION, CONTINGENCY.

1.2.1 Statement Pattern :

Letters used to denote statements are called statement letters. Proper combination of statement letters and connectives is called a statement pattern. Statement pattern is also called as a proposition. $p \rightarrow q, p \wedge q, \sim p \vee q$ are statement patterns. p and q are their prime components.

A table which shows the possible truth values of a statement pattern obtained by considering all possible combinations of truth values of its prime components is called the truth table of the statement pattern.

1.2.2. Logical Equivalence :

Two statement patterns are said to be equivalent if their truth tables are identical. If statement patterns A and B are equivalent, we write it as $A \equiv B$.

1.2.3 Tautology, Contradiction and Contingency :

Tautology : A statement pattern whose truth value is true for all possible combinations of truth values of its prime components is called a tautology. We denote tautology by t .

Statement pattern $p \vee \sim p$ is a tautology.

Contradiction : A statement pattern whose truth value is false for all possible combinations of truth values of its prime components is called a contradiction. We denote contradiction by c .

Statement pattern $p \wedge \sim p$ is a contradiction.

Contingency : A statement pattern which is neither a tautology nor a contradiction is called a contingency. $p \wedge q$ is a contingency.

Important table for all connectives :

p	q	$\sim p$	$p \wedge q$	$p \vee q$	$p \rightarrow q$	$p \leftrightarrow q$
T	T	F	T	T	T	T
T	F	F	F	T	F	F
F	T	T	F	T	T	F
F	F	T	F	F	T	T

★ In a statement pattern, different symbols are considered in the following priority

$\sim, \vee, \wedge, \rightarrow, \leftrightarrow$



Solved Examples

Ex.1.: Construct the truth table for each of the following statement patterns.

- $p \rightarrow (q \rightarrow p)$
- $(\sim p \vee q) \leftrightarrow \sim (p \wedge q)$
- $\sim (\sim p \wedge \sim q) \vee q$
- $[(p \wedge q) \vee r] \wedge [\sim r \vee (p \wedge q)]$
- $[(\sim p \vee q) \wedge (q \rightarrow r)] \rightarrow (p \rightarrow r)$

Solution :

i) $p \rightarrow (q \rightarrow p)$

p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$
T	T	T	T
T	F	T	T
F	T	F	T
F	F	T	T

Table 1.7

ii) $(\sim p \vee q) \leftrightarrow \sim(p \wedge q)$

p	q	$\sim p$	$\sim p \vee q$	$p \wedge q$	$\sim(p \wedge q)$	$(\sim p \vee q) \leftrightarrow \sim(p \wedge q)$
T	T	F	T	T	F	F
T	F	F	F	F	T	F
F	T	T	T	F	T	T
F	F	T	T	F	T	T

Table 1.8

iii) $\sim(\sim p \wedge \sim q) \vee q$

p	q	$\sim p$	$\sim q$	$\sim(p \wedge \sim q)$	$\sim(\sim p \wedge \sim q)$	$\sim(\sim p \wedge \sim q)$	$\sim(\sim p \wedge \sim q) \vee q$
T	T	F	F	F	T	T	T
T	F	F	T	F	T	T	T
F	T	T	F	F	T	T	T
F	F	T	T	T	F	F	F

Table 1.9

iv) $[(p \wedge q) \vee r] \wedge [\sim r \vee (p \wedge q)]$

p	q	r	$\sim r$	$p \wedge q$	$(p \wedge q) \vee r$	$\sim r \vee (p \wedge q)$	$[(p \wedge q) \vee r] \wedge [\sim r \vee (p \wedge q)]$
T	T	T	F	T	T	T	T
T	T	F	T	T	T	T	T
T	F	T	F	F	T	F	F
T	F	F	T	F	F	T	F
F	T	T	F	F	T	F	F
F	T	F	T	F	F	T	F
F	F	T	F	F	T	F	F
F	F	F	T	F	F	T	F

Table 1.10

v) $[(\sim p \vee q) \wedge (q \rightarrow r)] \rightarrow (p \rightarrow r)$

p	q	r	$\sim p$	$\sim p \vee q$	$q \rightarrow r$	$p \rightarrow r$	$(\sim p \vee q) \wedge (q \rightarrow r)$	$[(\sim p \vee q) \wedge (q \rightarrow r)] \rightarrow (p \rightarrow r)$
T	T	T	F	T	T	T	T	T
T	T	F	F	T	F	F	F	T
T	F	T	F	F	T	T	F	T
T	F	F	F	F	T	F	F	T
F	T	T	T	T	T	T	T	T
F	T	F	T	T	F	T	F	T
F	F	T	T	T	T	T	T	T
F	F	F	T	T	T	T	T	T

Table 1.11

Ex.2: Using truth tables, prove the following logical equivalences

- i) $(p \wedge q) \equiv \sim(p \rightarrow \sim q)$
- ii) $(p \leftrightarrow q) \equiv (p \wedge q) \vee (\sim p \wedge \sim q)$
- iii) $(p \wedge q) \rightarrow r \equiv p \rightarrow (q \rightarrow r)$
- iv) $p \rightarrow (q \vee r) \equiv (p \rightarrow q) \vee (p \rightarrow r)$

i) **Solution :** (1) $(p \wedge q) \equiv \sim(p \rightarrow \sim q)$

I	II	III	IV	V	VI
p	q	$\sim q$	$p \wedge q$	$p \rightarrow \sim q$	$\sim(p \rightarrow \sim q)$
T	T	F	T	F	T
T	F	T	F	T	F
F	T	F	F	T	F
F	F	T	F	T	F

Table 1.12

Columns (IV) and (VI) are identical $\therefore (p \wedge q) \equiv \sim(p \rightarrow \sim q)$

ii) $(p \leftrightarrow q) \equiv (p \wedge q) \vee (\sim p \wedge \sim q)$

I	II	III	IV	V	VI	VII	VIII
p	q	$\sim p$	$\sim q$	$p \leftrightarrow q$	$(p \wedge q)$	$\sim p \wedge \sim q$	$(p \wedge q) \vee (\sim p \wedge \sim q)$
T	T	F	F	T	T	F	T
T	F	F	T	F	F	F	F
F	T	T	F	F	F	F	F
F	F	T	T	T	F	T	T

Table 1.13

Columns V and VIII are identical

$\therefore (p \leftrightarrow q) \equiv (p \wedge q) \vee (\sim p \wedge \sim q)$

(iii) $(p \wedge q) \rightarrow r \equiv p \rightarrow (q \rightarrow r)$

I	II	III	IV	V	VI	VII
p	q	r	$p \wedge q$	$(p \wedge q) \rightarrow r$	$q \rightarrow r$	$p \rightarrow (q \rightarrow r)$
T	T	T	T	T	T	T
T	T	F	T	F	F	F
T	F	T	F	T	T	T
T	F	F	F	T	T	T
F	T	T	F	T	T	T
F	T	F	F	T	F	T
F	F	T	F	T	T	T
F	F	F	F	T	T	T

Table 1.14

Column (V) and (VII) are identical

$\therefore (p \wedge q) \rightarrow r \equiv p \rightarrow (q \rightarrow r)$

(iv) $p \rightarrow (q \vee r) \equiv (p \rightarrow q) \vee (p \rightarrow r)$

I	II	III	IV	V	VI	VII	VIII
p	q	r	$q \vee r$	$p \rightarrow (q \vee r)$	$p \rightarrow q$	$p \rightarrow r$	$(p \rightarrow q) \vee (p \rightarrow r)$
T	T	T	T	T	T	T	T
T	T	F	T	T	T	F	T
T	F	T	T	T	F	T	T
T	F	F	F	F	F	F	F
F	T	T	T	T	T	T	T
F	T	F	T	T	T	T	T
F	F	T	T	T	T	T	T
F	F	F	F	T	T	T	T

Table 1.15

Columns V and VIII are identical

$\therefore p \rightarrow (q \vee r) \equiv (p \rightarrow q) \vee (p \rightarrow r)$

Ex.3. Using truth tables, examine whether each of the following statements is a tautology or a contradiction or contingency.

- i) $(p \wedge q) \wedge (\sim p \vee \sim q)$
- ii) $[p \wedge (p \rightarrow \sim q)] \rightarrow p$
- iii) $(p \rightarrow q) \wedge [(q \rightarrow r) \rightarrow (p \rightarrow r)]$
- iv) $[(p \vee q) \vee r] \leftrightarrow [p \vee \leftrightarrow q \vee r]$

Solution:

i) $(p \wedge q) \wedge (\sim p \vee \sim q)$

p	q	$\sim p$	$\sim q$	$p \wedge q$	$\sim p \vee \sim q$	$(p \wedge q) \wedge (\sim p \wedge \sim q)$
T	T	F	F	T	F	F
T	F	F	T	F	T	F
F	T	T	F	F	T	F
F	F	T	T	F	T	F

Table 1.16

All the truth values in the last column are F. Hence it is contradiction.

ii) $[p \wedge (p \rightarrow \sim q)] \rightarrow q$

p	q	$\sim q$	$p \rightarrow \sim q$	$p \wedge (p \rightarrow \sim q)$	$p \wedge (p \rightarrow \sim q) \rightarrow q$
T	T	F	F	F	T
T	F	T	T	T	F
F	T	F	T	F	T
F	F	T	T	F	T

Table 1.17

Truth values in the last column are not identical. Hence it is contingency.

iii) $(p \rightarrow q) \wedge [(q \rightarrow r) \rightarrow (p \rightarrow r)]$

p	q	r	$p \rightarrow q$	$q \rightarrow r$	$p \rightarrow r$	$(q \rightarrow r) \rightarrow (p \rightarrow r)$	$(p \rightarrow q) \wedge (q \rightarrow r) \rightarrow (p \rightarrow r)$
T	T	T	T	T	T	T	T
T	T	F	T	F	F	T	T
T	F	T	F	T	T	T	F
T	F	F	F	T	F	F	F
F	T	T	T	T	T	T	T
F	T	F	T	F	T	T	T
F	F	T	T	T	T	T	T
F	F	F	T	T	T	T	T

Table 1.18

Truth values in the last column are not same, hence it is contingency.

iv) $[(p \vee q) \vee r] \leftrightarrow (p \vee (q \vee r))$

p	q	r	$p \vee q$	$(p \vee q) \vee r$	$q \vee r$	$p \vee (q \vee r)$	$[(p \vee q) \vee r] \leftrightarrow [p \vee (q \vee r)]$
T	T	T	T	T	T	T	T
T	T	F	T	T	T	T	T
T	F	T	T	T	T	T	T
T	F	F	T	T	F	T	T
F	T	T	T	T	T	T	T
F	T	F	T	T	T	T	T
F	F	T	F	T	T	T	T
F	F	F	F	F	F	F	T

Table 1.19

All the truth values in the last column are T, hence it is tautology.


Exercie.1.2

Q.1. Construct the truth table for each of the following statement patterns.

- i) $[(p \rightarrow q) \wedge q] \rightarrow p$
- ii) $(p \wedge \sim q) \leftrightarrow (p \rightarrow q)$
- iii) $(p \wedge q) \leftrightarrow (q \vee r)$
- iv) $p \rightarrow [\sim (q \wedge r)]$
- v) $\sim p \wedge [(p \vee \sim q) \wedge q]$
- vi) $(\sim p \rightarrow \sim q) \wedge (\sim q \rightarrow \sim p)$
- vii) $(q \rightarrow p) \vee (\sim p \leftrightarrow q)$
- viii) $[p \rightarrow (q \rightarrow r)] \leftrightarrow [(p \wedge q) \rightarrow r]$
- ix) $p \rightarrow [\sim (q \wedge r)]$
- x) $(p \vee \sim q) \rightarrow (r \wedge p)$

Q.2. Using truth tables prove the following logical equivalences.

- i) $\sim p \wedge q \equiv (p \vee q) \wedge \sim p$
- ii) $\sim (p \vee q) \vee (\sim p \wedge q) \equiv \sim p$
- iii) $p \leftrightarrow q \equiv \sim [(p \vee q) \wedge \sim (p \wedge q)]$
- iv) $p \rightarrow (q \rightarrow p) \equiv \sim p \rightarrow (p \rightarrow q)$
- v) $(p \vee q) \rightarrow r \equiv (p \rightarrow r) \wedge (q \rightarrow r)$
- vi) $p \rightarrow (q \wedge r) \equiv (p \rightarrow q) \wedge (p \rightarrow r)$
- vii) $p \rightarrow (q \wedge r) \equiv (p \wedge q) \rightarrow r$
- viii) $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$
- ix) $[\sim (p \vee q) \vee (p \vee q)] \wedge r \equiv r$
- x) $\sim (p \leftrightarrow q) \equiv (p \wedge \sim q) \vee (q \wedge \sim p)$

Q.3. Examine whether each of the following statement patterns is a tautology or a contradiction or a contingency.

- i) $(p \wedge q) \rightarrow (q \vee p)$
- ii) $(p \rightarrow q) \leftrightarrow (\sim p \vee q)$
- iii) $[\sim (\sim p \wedge \sim q)] \vee q$
- iv) $[(p \rightarrow q) \wedge q] \rightarrow p$
- v) $[(p \rightarrow q) \wedge \sim q] \rightarrow \sim p$
- vi) $(p \leftrightarrow q) \wedge (p \rightarrow \sim q)$
- vii) $\sim (\sim q \wedge p) \wedge q$
- viii) $(p \wedge \sim q) \leftrightarrow (p \rightarrow q)$
- ix) $(\sim p \rightarrow q) \wedge (p \wedge r)$
- x) $[p \rightarrow (\sim q \vee r)] \leftrightarrow \sim [p \rightarrow (q \rightarrow r)]$

1.3 QUANTIFIERS, QUANTIFIED STATEMENTS, DUALS, NEGATION OF COMPOUND STATEMENTS, CONVERSE, INVERSE AND CONTRAPOSITIVE OF IMPLICATION.

1.3.1 Quantifiers and quantified statements.

Look at the following statements :

p : "There exists an even prime number in the set of natural numbers".

q : "All natural numbers are positive".

Each of them asserts a *condition* for some or all objects in a *collection*. Words "there exists" and "for all" are called quantifiers. "There exists" is called existential quantifier and is denoted by symbol \exists . "For all" is called universal quantifier and is denoted by \forall . Statements involving quantifiers are called quantified statements. Every quantified statement corresponds to a *collection* and a *condition*. In statement p the collection is 'the set of natural numbers' and the condition is 'being even prime'. What is the condition in the statement q ?

A statement quantified by universal quantifier \forall is true if all objects in the collection satisfy the condition. And it is false if at least one object in the collection does not satisfy the condition.

A statement quantified by existential quantifier \exists is true if at least one object in the collection satisfies the condition. And it is false if no object in the collection satisfies the condition.

Ex.1. If $A = \{1, 2, 3, 4, 5, 6, 7\}$, determine the truth value of the following.

- i) $\exists x \in A$ such that $x - 4 = 3$
- ii) $\forall x \in A, x + 1 \geq 3$
- iii) $\forall x \in A, 8 - x \leq 7$
- iv) $\exists x \in A$, such that $x + 8 = 16$

Solution :

- i) For $x = 7, x - 4 = 7 - 4 = 3$
 $\therefore x = 7$ satisfies the equation $x - 4 = 3$
 \therefore The given statement is true and its truth value is T.
- ii) For $x = 1, x + 1 = 1 + 1 = 2$ which is not greater than or equal to 3
 \therefore For $x = 1, x + 1 \geq 3$ is not true.
 \therefore The truth value of given statement is F.
- iii) For each $x \in A, 8 - x \leq 7$
 \therefore The given statement is true.
 \therefore Its truth value is T.
- iv) There is no x in A which satisfies $x + 8 = 16$.
 \therefore The given statement is false. \therefore Its truth value is F.

1.3.2 Dual : We use letters t and c to denote tautology and contradiction respectively.

If two statements contain logical connectives like \vee, \wedge and letters t and c then they are said to be duals of each other if one of them is obtained from the other by interchanging \vee with \wedge and t with c .

The dual of i) $p \vee q$ is $p \wedge q$ ii) $t \vee p$ is $c \wedge p$ iii) $t \wedge p$ is $c \vee p$

Ex.1. Write the duals of each of the following :

i) $(p \wedge q) \vee r$

ii) $t \vee (p \vee q)$

iii) $p \wedge [\sim q \vee (p \wedge q) \vee \sim r]$

iv) $(p \vee q) \wedge t$

v) $(p \vee q) \vee r \equiv p \vee (q \vee r)$

vi) $p \wedge q \wedge r$

vii) $(p \wedge t) \vee (c \wedge \sim q)$

Solution :

i) $(p \vee q) \wedge r$

ii) $c \wedge (p \wedge q)$

iii) $p \vee [(\sim q \wedge (p \vee q) \wedge \sim r)]$

iv) $(p \wedge q) \vee c$

v) $(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$

vi) $p \vee q \vee r$

vii) $(p \vee c) \wedge (t \vee \sim q)$

1.3.3 Negations of compound statements :

Negation of conjunction : When is the statement "6 is even and perfect number" is false? It is so, if 6 is not even or 6 is not perfect number. The negation of $p \wedge q$ is $\sim p \vee \sim q$. The negation of "6 is even and perfect number" is "6 not even or not perfect number".

Activity : Using truth table verify that $\sim (p \wedge q) \equiv \sim p \vee \sim q$

Negation of disjunction : When is the statement "x is prime or y is even" is false? It is so, if x is not prime and y is not even. The negation of $p \vee q$ is $\sim p \wedge \sim q$. The negation of "x is prime or y is even" is "x is not prime and y is not even".

Activity : Using truth table verify that $\sim (p \vee q) \equiv \sim p \wedge \sim q$

Note : ' $\sim (p \wedge q) \equiv \sim p \vee \sim q$ ' and ' $\sim (p \vee q) \equiv \sim p \wedge \sim q$ ' are called **De'Morgan's Laws**

Negation of implication : Implication $p \rightarrow q$ asserts that "if p is true statement then q is true statement". When is an implication a true statement and when is it false? Consider the statement "If bakery is open then I will buy a cake for you." Clearly statement is false only when the bakery was open and I did not buy a cake for you. The conditional statement "If p then q" is false only in the case "p is true and q is false". In all other cases it is true. The negation of the statement "If p then q" is the statement "p and not q". i.e. p does not imply q

Activity : Using truth table verify that $\sim (p \rightarrow q) \equiv p \wedge \sim q$

Negation of biconditional : The biconditional $p \leftrightarrow q$ is the conjunction of statement $p \rightarrow q$ and $q \rightarrow p$.

$$\therefore p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$$

\therefore The conditional statement $p \leftrightarrow q$ is false if $p \rightarrow q$ is false or $q \rightarrow p$ false.

The negation of the statement "p if and only if q" is the statement "p and not q, or q and not p".

$$\therefore \sim (p \leftrightarrow q) \equiv (p \wedge \sim q) \vee (q \wedge \sim p)$$

Activity : Using truth table verify that $\sim (p \leftrightarrow q) \equiv (p \wedge \sim q) \vee (q \wedge \sim p)$

1.3.4 Converse, inverse and contrapositive

From implication $p \rightarrow q$ we can obtain three implications, called converse, inverse and contrapositive.

$q \rightarrow p$ is called the converse of $p \rightarrow q$

$\sim p \rightarrow \sim q$ is called the inverse of $p \rightarrow q$.

$\sim q \rightarrow \sim p$ is called the contrapositive of $p \rightarrow q$.

Activity :

Prepare the truth table for $p \rightarrow q$, $q \rightarrow p$, $\sim p \rightarrow \sim q$ and $\sim q \rightarrow \sim p$. What is your conclusion from the truth table ?

- i) \equiv
- ii) \equiv

Ex.1) Write the negations of the following.

- i) $3 + 3 < 5$ or $5 + 5 = 9$
- ii) $7 > 3$ and $4 > 11$
- iii) The number is neither odd nor perfect square.
- iv) The number is an even number if and only if it is divisible by 2.

Solution :

- i) Let $p : 3 + 3 < 5$; $q : 5 + 5 = 9$
 Given statement is $p \vee q$ and its negation is $\sim(p \vee q)$ and $\sim(p \vee q) \equiv \sim p \wedge \sim q$
 \therefore The negation of given statement is $3 + 3 \geq 5$ and $5 + 5 \neq 9$
- ii) Let $p : 7 > 3$; $q : 4 > 11$
 The given statement is $p \wedge q$
 Its negation is $\sim(p \wedge q)$ and $\sim(p \wedge q) \equiv \sim p \vee \sim q$
 \therefore The negation of given statement is $7 \leq 3$ or $4 \leq 11$
- iii) Let p : The number is odd
 q : The number is perfect square
 Given statement can be written as 'the number is not odd and not perfect square'
 Given statement is $\sim p \wedge \sim q$
 Its negation is $\sim(\sim p \wedge \sim q) \equiv p \vee q$
 The negation of given statement is 'The number is odd or perfect square'.
- iv) Let p : The number is an even number.
 q : The number is divisible by 2
 Given statement is $p \leftrightarrow q$
 Its negation is $\sim(p \leftrightarrow q)$
 But $\sim(p \leftrightarrow q) \equiv (p \wedge \sim q) \vee (q \wedge \sim p)$
 \therefore The negation of given statement is 'A number is even but not divisible by 2 or a number is divisible by 2 but not even'.

Ex.2. Write the negations of the following statements.

- i) All natural numbers are rational.
- ii) Some students of class X are sixteen year old.
- iii) $\exists n \in \mathbb{N}$ such that $n + 8 > 11$
- iv) $\forall x \in \mathbb{N}$, $2x + 1$ is odd

Solution :

- i) Some natural numbers are not rationals.
- ii) No student of class X is sixteen year old.
- iii) $\forall n \in \mathbb{N}, n + 8 \leq 11$
- iv) $\exists x \in \mathbb{N}$ such that $2x + 1$ is not odd

Ex.3. Write the converse, inverse and contrapositive of the following statements.

- i) If a function is differentiable then it is continuous.
- ii) If it rains then the match will be cancelled.

Solution :

- (1) Let p : A function is differentiable
 q : A function is continuous.
 \therefore Given statement is $p \rightarrow q$
- i) Its converse is $q \rightarrow p$
 If a function is continuous then it is differentiable.
 - ii) Its inverse is $\sim p \rightarrow \sim q$.
 If a function not differentiable then it is not continuous.
 - iii) Its contrapositive is $\sim q \rightarrow \sim p$
 If a function is not continuous then it is not differentiable.
- (2) Let p : It rains, q : The match gets cancelled.
 \therefore Given statement is $p \rightarrow q$
- i) Its converse is $q \rightarrow p$
 If the match gets cancelled then it rains.
 - ii) Inverse is $\sim p \rightarrow \sim q$
 If it does not rain then the match will not be cancelled.
 - iii) Its contrapositive is $\sim q \rightarrow \sim p$.
 If the match is not cancelled then it does not rain.



Exercise 1.3

Q.1. If $A = \{3, 5, 7, 9, 11, 12\}$, determine the truth value of each of the following.

- i) $\exists x \in A$ such that $x - 8 = 1$
- ii) $\forall x \in A, x^2 + x$ is an even number
- iii) $\exists x \in A$ such that $x^2 < 0$
- iv) $\forall x \in A, x$ is an even number
- v) $\exists x \in A$ such that $3x + 8 > 40$
- vi) $\forall x \in A, 2x + 9 > 14$

Q.2. Write the duals of each of the following.

- i) $p \vee (q \wedge r)$
- ii) $p \wedge (q \wedge r)$
- iii) $(p \vee q) \wedge (r \vee s)$
- iv) $p \wedge \sim q$
- v) $(\sim p \vee q) \wedge (\sim r \wedge s)$
- vi) $\sim p \wedge (\sim q \wedge (p \vee q) \wedge \sim r)$
- vii) $[\sim (p \vee q)] \wedge [p \vee \sim (q \wedge \sim s)]$
- viii) $c \vee \{p \wedge (q \vee r)\}$
- ix) $\sim p \vee (q \wedge r) \wedge t$
- x) $(p \vee q) \vee c$

Q.3. Write the negations of the following.

- i) $x + 8 > 11$ or $y - 3 = 6$
- ii) $11 < 15$ and $25 > 20$
- iii) Quadrilateral is a square if and only if it is a rhombus.
- iv) It is cold and raining.
- v) If it is raining then we will go and play football.
- vi) $\sqrt{2}$ is a rational number.
- vii) All natural numbers are whole numbers.
- viii) $\forall n \in \mathbb{N}, n^2 + n + 2$ is divisible by 4.
- ix) $\exists x \in \mathbb{N}$ such that $x - 17 < 20$

Q.4. Write converse, inverse and contrapositive of the following statements.

- i) If $x < y$ then $x^2 < y^2$ ($x, y \in \mathbb{R}$)
- ii) A family becomes literate if the woman in it is literate.
- iii) If surface area decreases then pressure increases.
- iv) If voltage increases then current decreases.

1.4 SOME IMPORTANT RESULTS :

1.4.1.

- i) $p \rightarrow q \equiv \sim p \vee q$
- ii) $p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$
- iii) $p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$
- iv) $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$

I	II	III	IV	V	VI	VII	VIII
p	q	$\sim p$	$p \rightarrow q$	$q \rightarrow p$	$p \leftrightarrow q$	$\sim p \vee q$	$p \rightarrow q \wedge q \rightarrow p$
T	T	F	T	T	T	T	T
T	F	F	F	T	F	F	F
F	T	T	T	F	F	T	F
F	F	T	T	T	T	T	T

Table 1.20

Columns (IV, VII) and (VI, VIII) are identical.

$\therefore p \rightarrow q \equiv \sim p \vee q$ and $p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$ are proved.

Activity :

Prove the results (iii) and (iv) by using truth table.

1.4.2. Algebra of statements.

Idempotent Law	$p \wedge p \equiv p, \quad p \vee p \equiv p$
Commutative Law	$p \vee q \equiv q \vee p \quad p \wedge q \equiv q \wedge p$
Associative Law	$p \wedge (q \wedge r) \equiv (p \wedge q) \wedge r \equiv p \wedge q \wedge r$
	$p \vee (q \vee r) \equiv (p \vee q) \vee r \equiv p \vee q \vee r$
Distributive Law	$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$
	$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$
De Morgan's Law	$\sim(p \wedge q) \equiv \sim p \vee \sim q, \quad \sim(p \vee q) \equiv \sim p \wedge \sim q$
Identity Law	$p \wedge T \equiv p, \quad p \wedge F \equiv F, \quad p \vee F \equiv p, \quad p \vee T \equiv T$
Complement Law	$p \wedge \sim p \equiv F, \quad p \vee \sim p \equiv T$
Absorption Law	$p \vee (p \wedge q) \equiv p, \quad p \wedge (p \vee q) \equiv p$
Conditional Law	$p \rightarrow q \equiv \sim p \vee q$
Biconditional Law	$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p) \equiv (\sim p \vee q) \wedge (\sim q \vee p)$

Table 1.21



Solved Examples

Ex.1. Write the negations of the following stating the rules used.

- i) $(p \vee q) \wedge (q \vee \sim r)$
- ii) $(p \rightarrow q) \vee r$
- iii) $p \wedge (q \vee r)$
- iv) $(\sim p \wedge q) \vee (p \wedge \sim q)$
- v) $(p \wedge q) \rightarrow (\sim p \vee r)$

Solution :

- i) $\sim [(p \vee q) \wedge (q \vee \sim r)] \equiv \sim (p \vee q) \vee \sim (q \vee \sim r)$ [DeMorgan's law]
 $\equiv (\sim p \wedge \sim q) \vee (\sim q \wedge r)$ [DeMorgan's law]
 $\equiv (\sim q \wedge \sim p) \vee (\sim q \wedge r)$ [Commutative law]
 $\equiv \sim q \wedge (\sim p \vee r)$ [Distributive law]
- ii) $\sim [(p \rightarrow q) \vee r] \equiv \sim (p \rightarrow q) \wedge \sim r$ [DeMorgan's law]
 $\equiv (p \wedge \sim q) \wedge \sim r$ [$\sim (p \rightarrow q) \equiv p \wedge \sim q$]
- iii) $\sim [p \wedge (q \vee r)] \equiv \sim p \vee \sim (q \vee r)$ [DeMorgan's law]
 $\equiv \sim p \vee (\sim q \wedge \sim r)$ [DeMorgan's law]
- iv) $\sim [(\sim p \wedge q) \vee (p \wedge \sim q)] \equiv \sim (\sim p \wedge q) \wedge \sim (p \wedge \sim q)$ [DeMorgan's law]
 $\equiv (p \vee \sim q) \wedge (\sim p \vee q)$ [DeMorgan's law]
- v) $\sim [(p \wedge q) \rightarrow (\sim p \vee r)] \equiv (p \wedge q) \wedge \sim (\sim p \vee r)$ [$\sim (p \rightarrow q) \equiv p \wedge \sim q$]
 $\equiv (p \wedge q) \wedge [p \wedge \sim r]$ [DeMorgan's law]
 $\equiv q \wedge p \wedge p \wedge \sim r$ [Associative law]
 $\equiv q \wedge p \wedge \sim r$ [Idempotent law]

Ex.2. Rewrite the following statements without using if then.

- i) If prices increase then the wages rise.
- ii) If it is cold, then we wear woollen clothes.

Solution :

- i) Let p : Prices increase
 q : The wages rise.
 The given statement is $p \rightarrow q$
 but $p \rightarrow q \equiv \sim p \vee q$
 The given statement can be written as
 'Prices do not increase or the wages rise'.
- ii) Let p : It is cold, q : We wear woollen clothes.
 The given statement is $p \rightarrow q$
 but $p \rightarrow q \equiv \sim p \vee q$
 The given statement can be written as
 It is not cold or we wear woollen clothes.

Ex.3. Without using truth table prove that :

- i) $p \leftrightarrow q \equiv \sim (p \wedge \sim q) \wedge \sim (q \wedge \sim p)$
- ii) $\sim (p \vee q) \vee (\sim p \wedge q) \equiv \sim p$
- iii) $\sim p \wedge q \equiv (p \vee q) \wedge \sim p$

Solution :

- i) We know that

$$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$$

$$\equiv (\sim p \vee q) \wedge (\sim q \vee p)$$

$$\equiv \sim (p \wedge \sim q) \wedge \sim (q \wedge \sim p)$$

[Conditional law]
 [Demorgan's law]
- ii) $\sim (p \vee q) \vee (\sim p \wedge q) \equiv (\sim p \wedge \sim q) \vee (\sim p \wedge q)$

$$\equiv \sim p \wedge (\sim q \vee q)$$

$$\equiv \sim p \wedge T$$

$$\equiv \sim p$$

[Demorgan's law]
 [Distributive law]
 [Complement law]
 [Identity law]
- iii) $(p \wedge q) \wedge \sim p \equiv \sim p \wedge (p \vee q)$

$$\equiv (\sim p \wedge p) \vee (\sim p \wedge q)$$

$$\equiv F \vee (\sim p \wedge q)$$

$$\equiv \sim p \wedge q$$

[Commutative law]
 [Distributive law]
 [Complement law]
 [Identity law]



Exercise 1.4

Q.1. Using rules of negation write the negations of the following with justification.

- | | |
|--|---|
| i) $\sim q \rightarrow p$ | ii) $p \wedge \sim q$ |
| iii) $p \vee \sim q$ | iv) $(p \vee \sim q) \wedge r$ |
| v) $p \rightarrow (p \vee \sim q)$ | vi) $\sim (p \wedge q) \vee (p \vee \sim q)$ |
| vii) $(p \vee \sim q) \rightarrow (p \wedge \sim q)$ | viii) $(\sim p \vee \sim q) \vee (p \wedge \sim q)$ |

Q.2. Rewrite the following statements without using if .. then.

- i) If a man is a judge then he is honest.
- ii) If 2 is a rational number then $\sqrt{2}$ is irrational number.
- iii) If $f(2) = 0$ then $f(x)$ is divisible by $(x - 2)$.

Q.3. Without using truth table prove that :

- i) $p \leftrightarrow q \equiv (p \wedge q) \vee (\sim p \wedge \sim q)$
- ii) $(p \vee q) \wedge (p \vee \sim q) \equiv p$
- iii) $(p \wedge q) \vee (\sim p \wedge q) \vee (p \wedge \sim q) \equiv p \vee q$
- iv) $\sim [(p \vee \sim q) \rightarrow (p \wedge \sim q)] \equiv (p \vee \sim q) \wedge (\sim p \vee q)$

Application of Logic to switching circuits :

We shall study how the theory of Logic can be applied in switching network. We have seen that a logical statement can be either true or false i.e. it can have truth value either T or F.

A similar situation exists in various electrical devices. For example, an electric switch can be on or off. In 1930 Claude Shannan noticed an analogy between operation of switching circuits and operation of logical connectives.

In an electric circuit, switches are connected by wires. If the switch is 'on', it allows the electric current to pass through, it. If the switch is 'off', it does not allow the electric current to pass through it. We now define the term 'switch' as follows.

Switch : A switch is a two state device used to control the flow of current in a circuit.

We shall denote the switches by letters S, S₁, S₂, S₃ etc.




Fig. 1.1




Fig. 1.2

In figure 1.2, we consider a circuit containing an electric lamp L, controlled by a switch S.

When the switch S is closed (i.e. on), then current flows in the circuit and hence the lamp glows. When the switch S is open (i.e. off), then current does not flow in the circuit and subsequently the lamp does not glow.

The theory of symbolic logic can be used to represent a circuit by a statement pattern. Conversely for given statement pattern a circuit can be constructed. Corresponding to each switch in the circuit we take a statement letter in statement pattern. Switches having the same state will be denoted by the same letter and called equivalent switches. Switches having opposite states are denoted by S and S'. They are called complementary switches. In circuit we don't show whether switch is open or closed. In figure 1.3 switch S₁ corresponds to statement letter p in the corresponding statement pattern.

We write it as p : switch S₁ and $\sim p$: switch S'₁

The correspondence between switch S₂ and statement letter q is shown as q : switch S₂ and $\sim q$: switch S'₂.

We don't know the actual states of switches in the circuit. We consider all possible combinations of states of all switches in the circuit and prepare a table, called "Input Output table", which is similar to truth table of the corresponding statement pattern.

★ In an Input-output table we represent '1' when the state of the switch is 'on' and '0' when the state of the switch is 'off'.

1.5.1. Two switches in series.

Two switches S₁ and S₂ connected in series and electric lamp 'L' as shown in fig 1.3.




Fig. 1.3

Let p : The switch S₁

q : The switch S₂

L : The lamp L

Input output table (switching table) for $p \wedge q$.

p	q	$p \wedge q$
1	1	1
1	0	0
0	1	0
0	0	0

Table 1.22

1.5.2 Two switches in parallel :

Two switches S_1 and S_2 are connected in parallel and electric lamp L is as shown in fig. 1.4

Let p : The switch S_1
 q : The switch S_2
 L : The lamp L

Input - output table. for $p \vee q$.

p	q	$p \vee q$
1	1	1
1	0	1
0	1	1
0	0	0

Table 1.23




Fig. 1.4

Q.1. Express the following circuits in the symbolic form of logic and write the input-output table.

i)




Fig. 1.5

ii)




Fig. 1.6

iii)




Fig. 1.7

Solution :

- i) Let p : The switch S_1 q : The switch S_2 L : The lamp L
 Given circuit is expressed as $(p \vee q) \vee (\sim p \wedge \sim q)$

Solution :

p	q	$\sim p$	$\sim q$	$p \vee q$	$\sim p \wedge \sim q$	$(p \vee q) \vee (\sim p \wedge \sim q)$
1	1	0	0	1	0	1
1	0	0	1	1	0	1
0	1	1	0	1	0	1
0	0	1	1	0	1	1

Table 1.24

- ii) Let p : The switch S_1 is closed
 q : The switch S_2 r : The switch S_3 L : The lamp L

The symbolic form is $[(p \wedge q) \vee (\sim p \wedge \sim q)] \wedge r$

p	q	r	$\sim p$	$\sim q$	$p \wedge q$	$(p \wedge q) \vee (\sim p \wedge \sim q)$	$(p \wedge q) \vee (\sim p \wedge \sim q)$	$[(p \wedge q) \vee (\sim p \wedge \sim q)] \wedge r$
1	1	1	0	0	1	0	1	1
1	1	0	0	0	1	0	1	0
1	0	1	0	1	0	0	0	0
1	0	0	0	1	0	0	0	0
0	1	1	1	0	0	0	0	0
0	1	0	1	0	0	0	0	0
0	0	1	1	1	0	1	1	1
0	0	0	1	1	0	1	1	0

Table 1.25

- iii) Let p : The switch S_1 q : The switch S_2
 r : The switch S_3 L : The lamp L

The symbolic form of given circuit is $(p \vee q) \wedge q \wedge (r \vee \sim p)$

p	q	r	$\sim p$	$p \vee q$	$r \vee \sim p$	$(p \vee q) \wedge \sim q$	$(p \vee q) \wedge q \wedge (r \vee \sim p)$
1	1	1	0	1	1	1	1
1	1	0	0	1	0	1	0
1	0	1	0	1	1	0	0
1	0	0	0	1	0	0	0
0	1	1	1	1	1	1	1
0	1	0	1	1	1	1	1
0	0	1	1	0	1	0	0
0	0	0	1	0	1	0	0

Table 1.26

Ex.2. Construct switching circuits of the following.

- i) $[(p \vee (\sim p \wedge q)) \vee [(\sim q \wedge r) \vee \sim p]$
- ii) $(p \wedge q \wedge r) \vee [p \vee (q \wedge \sim r)]$
- iii) $[(p \wedge r) \vee (\sim q \wedge \sim r)] \vee (\sim p \wedge \sim r)$

Solution :

Let p : The Switch S_1

q : The switch S_2

r : The switch S_3

The circuits are as follows.




Fig. 1.8




Fig. 1.9




Fig. 1.10

Ex.3. Give an alternative arrangement for the following circuit, so that the new circuit has minimum switches.




Fig. 1.11

Let p : The switch S_1

q : The switch S_2

The symbolic form is $(p \wedge \sim q) \vee (\sim p \wedge q) \vee (\sim p \wedge \sim q)$

Consider $(p \wedge \sim q) \vee (\sim p \wedge q) \vee (\sim p \wedge \sim q)$

$$\equiv (p \wedge \sim q) \vee [\sim p \wedge (q \vee \sim q)]$$

$$\equiv (p \wedge \sim q) \vee [\sim p \wedge T]$$

$$\equiv (p \wedge \sim q) \vee \sim p$$

$$\equiv \sim p \vee (p \wedge \sim q)$$

$$\equiv (\sim p \vee p) \wedge (\sim p \wedge \sim q)$$

$$\equiv t \wedge (\sim p \vee \sim q)$$

$$\equiv \sim p \vee \sim q$$

[Distributive Law]

[Complement Law]

[Identity Law]

[Commutative Law]

[Distributive Law]

[Law of Complement]

[Identity Law]

The alternative arrangement for the given circuit is as follows :




Fig. 1.12

Ex.4. Express the following switching circuit in the symbolic form of Logic. Construct the switching table and interpret it.




Fig. 1.13

Solution :

Let p : The switch S_1

q : The switch S_2

The symbolic form of the given switching circuit is $(p \vee q) \wedge (\sim p) \wedge (\sim q)$

The switching table.

p	q	$\sim p$	$\sim q$	$p \vee q$	$(p \vee q) \wedge (\sim p)$	$(p \vee q) \wedge (\sim p) \wedge (\sim q)$
1	1	0	0	1	0	0
1	0	0	1	1	0	0
0	1	1	0	1	1	0
0	0	1	1	0	0	0

Table 1.27

Last column contains all 0, lamp will not glow irrespective of the status of the switches.

Ex.5. Simplify the given circuit by writing its logical expression. Also,write your conclusion.




Fig. 1.14

Let p : The switch S_1

q : The The switch S_2

The logical expression for the given circuit is $p \wedge (\sim p \vee \sim q) \wedge q$

Consider

$$\begin{aligned}
 & p \wedge (\sim p \vee \sim q) \wedge q \\
 \equiv & [p \wedge (\sim p \vee \sim q)] \wedge q && \text{[Associative Law]} \\
 \equiv & [(p \wedge \sim p) \vee (p \wedge \sim q)] \wedge q && \text{[Distributive Law]} \\
 \equiv & [F \vee (p \wedge \sim q)] \wedge q && \text{[Complement Law]} \\
 \equiv & (p \wedge \sim q) \wedge q && \text{[Identity Law]} \\
 \equiv & p \wedge (\sim q \wedge q) && \text{[Associative Law]} \\
 \equiv & p \wedge F && \text{[Complement Law]} \\
 \equiv & F && \text{[Identity Law]}
 \end{aligned}$$

Conclusion : The lamp will not glow irrespective of the status of the switches.

Ex. 6 : In the following switching circuit,

- i) Write symbolic form ii) Construct switching table iii) Simplify the circuit




Fig. 1.15

Solution : Let p : The switch S_1 . q : The switch S_2 . r : The switch S_3

1) The symbolic form of given circuit is $(p \wedge q) \vee (\sim p \wedge q) \vee (r \wedge \sim q)$.

ii) Switching Table :

P	q	r	$\sim p$	$\sim q$	$\sim p \wedge q$	$\sim p \wedge \sim q$	$r \wedge \sim q$	$(p \wedge q) \vee (\sim p \wedge q) \vee (r \wedge \sim q)$
1	1	1	0	0	1	0	0	1
1	1	0	0	0	1	0	0	1
1	0	1	0	1	0	0	1	1
1	0	0	0	1	0	0	0	0
0	1	1	1	0	0	1	0	1
0	1	0	1	0	0	1	0	1
0	0	1	1	1	0	0	1	1
0	0	0	1	1	0	0	0	0

Table 1.28

iii) Consider $= (p \wedge q) \vee (\sim p \wedge q) \vee (r \wedge \sim q)$
 $= [(p \vee \sim p) \wedge q] \vee [(r \wedge \sim q)]$ [Distributive Law]
 $= (T \wedge q) \vee (r \wedge \sim q)$ [Complement Law]
 $= q \vee (r \wedge \sim q)$ [Identity Law]
 $= (q \vee r) \wedge (q \vee \sim q)$ [Distributive Law]
 $= (q \vee r) \wedge T$ [Complement Law]
 $= (q \vee r)$ [Identify Law]

Simplified circuit is :




Fig. 1.16

Q.1. Express the following circuits in the symbolic form of logic and write the input-output table.




Fig. 1.17




Fig. 1.18




Fig. 1.19




Fig. 1.20




Fig. 1.21




Fig. 1.22

Q.2. Construct the switching circuit of the following :

- i) $(\sim p \wedge q) \vee (p \wedge \sim r)$
- ii) $(p \wedge q) \vee [\sim p \wedge (\sim q \vee p \vee r)]$
- iii) $(p \wedge r) \vee (\sim q \wedge \sim r) \wedge (\sim p \wedge \sim r)$
- iv) $(p \wedge \sim q \wedge r) \vee [p \wedge (\sim q \vee \sim r)]$
- v) $p \vee (\sim p) \vee (\sim q) \vee (p \wedge q)$
- vi) $(p \wedge q) \vee (\sim p) \vee (p \wedge \sim q)$

Q.3. Give an alternative equivalent simple circuits for the following circuits :




Fig. 1.23




Fig. 1.24

Q.4. Write the symbolic form of the following switching circuits construct its switching table and interpret it.




Fig. 1.25




Fig. 1.26




Fig. 1.27

Q.5. Obtain the simple logical expression of the following. Draw the corresponding switching circuit.

- i) $p \vee (q \wedge \sim q)$
- ii) $(\sim p \wedge q) \vee (\sim p \wedge \sim q) \vee (p \wedge \sim q)$
- iii) $[p \vee (\sim q) \vee \sim r] \wedge (p \vee (q \wedge r))$
- iv) $(p \wedge q \wedge \sim p) \vee (\sim p \wedge q \wedge r) \vee (p \wedge \sim q \wedge r)$



Let's remember!

- 1) A declarative sentence which is either true or false, but not both simultaneously is called a statement.

Sr. No.	Connective	Symbolic Form	Name of Compound Statement	Hint in truth table	Negation
i)	and	$p \wedge q$	Conjunction	$T \wedge T \equiv T$	$\sim p \vee \sim q$
ii)	or	$p \vee q$	Disjunction	$F \vee F \equiv F$	$\sim p \wedge \sim q$
iii)	if... then	$p \rightarrow q$	Conditional	$T \rightarrow F \equiv F$	$p \wedge \sim q$
iv)	if and only if	$p \leftrightarrow q$	Biconditional	$T \leftrightarrow T \equiv T$ $F \leftrightarrow F \equiv T$	$(p \wedge \sim q) \vee (q \wedge \sim p)$

Table 1.29

- 3) In the truth table of the statement pattern if all truth values in the last column
 a) are 'T' then it is tautology.
 b) are 'F' then it is contradiction.
- 4) In the truth table of the statement pattern if some entries are 'T' and some are 'F' then it is called as contingency.
- 5) The symbol \forall stands for 'for all' or 'for every'. It is universal quantifier. The symbol \exists stands for 'for some' or 'for one' or 'there exists at least one'. It is called as existential quantifier.
- 6) Algebra of statements.

Idempotent Law	$p \wedge p \equiv p, \quad p \vee p \equiv p$
Commutative Law	$p \vee q \equiv q \vee p \quad p \wedge q \equiv q \wedge p$
Associative Law	$p \wedge (q \wedge r) \equiv (p \wedge q) \wedge r \equiv p \wedge q \wedge r$ $p \vee (q \vee r) \equiv (p \vee q) \vee r \equiv p \vee q \vee r$
Distributive Law	$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$ $p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$
De Morgan's Law	$\sim(p \wedge q) \equiv \sim p \vee \sim q, \quad \sim(p \vee q) \equiv \sim p \wedge \sim q$
Identity Law	$p \wedge T \equiv p, \quad p \wedge F \equiv F, \quad p \vee F \equiv p, \quad p \vee T \equiv T$
Complement Law	$p \wedge \sim p \equiv F, \quad p \vee \sim p \equiv T$
Absorption Law	$p \vee (p \wedge q) \equiv p, \quad p \wedge (p \vee q) \equiv p$
Conditional Law	$p \rightarrow q \equiv \sim p \vee q$
Biconditional Law	$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p) \equiv (\sim p \vee q) \wedge (\sim q \vee p)$

- 7) If $p \rightarrow q$ is conditional then its converse is $q \rightarrow p$, inverse is $\sim p \rightarrow \sim q$ and contrapositive is $\sim q \rightarrow \sim p$.

8) Switching circuits :

i) Switches in series




Fig. 1.28

ii) Switches in parallel.




Fig. 1.29

Input-output table

p	q	$p \wedge q$	$p \vee q$
1	1	1	1
1	0	0	1
0	1	0	1
0	0	0	0

Table 1.30

Miscellaneous Exercise 1

I] Select and write the correct answer from the given alternatives in each of the following questions :

- i) If $p \wedge q$ is false and $p \vee q$ is true, the _____ is not true.
 A) $p \vee q$ B) $p \leftrightarrow q$ C) $\sim p \vee \sim q$ D) $q \vee \sim p$
- ii) $(p \wedge q) \rightarrow r$ is logically equivalent to _____.
 A) $p \rightarrow (q \rightarrow r)$ B) $(p \wedge q) \rightarrow \sim r$ C) $(\sim p \vee \sim q) \rightarrow \sim r$ D) $(p \vee q) \rightarrow r$
- iii) Inverse of statement pattern $(p \vee q) \rightarrow (p \wedge q)$ is _____.
 A) $(p \wedge q) \rightarrow (p \vee q)$ B) $\sim(p \vee q) \rightarrow (p \wedge q)$
 C) $(\sim p \wedge \sim q) \rightarrow (\sim p \vee \sim q)$ D) $(\sim p \vee \sim q) \rightarrow (\sim p \wedge \sim q)$
- iv) If $p \wedge q$ is F, $p \rightarrow q$ is F then the truth values of p and q are _____.
 A) T, T B) T, F C) F, T D) F, F
- v) The negation of inverse of $\sim p \rightarrow q$ is _____.
 A) $q \wedge p$ B) $\sim p \wedge \sim q$ C) $p \wedge q$ D) $\sim q \rightarrow \sim p$
- vi) The negation of $p \wedge (q \rightarrow r)$ is _____.
 A) $\sim p \wedge (\sim q \rightarrow \sim r)$ B) $p \vee (\sim q \vee r)$
 C) $\sim p \wedge (\sim q \rightarrow \sim r)$ D) $\sim p \vee (\sim q \wedge \sim r)$
- vii) If $A = \{1, 2, 3, 4, 5\}$ then which of the following is not true?
 A) $\exists x \in A$ such that $x + 3 = 8$ B) $\exists x \in A$ such that $x + 2 < 9$
 C) $\forall x \in A, x + 6 \geq 9$ D) $\exists x \in A$ such that $x + 6 < 10$

Q.2. Which of the following sentences are statements in logic? Justify. Write down the truth value of the statements :

- i) $4! = 24$.
- ii) π is an irrational number.
- iii) India is a country and Himalayas is a river.
- iv) Please get me a glass of water.
- v) $\cos^2\theta - \sin^2\theta = \cos 2\theta$ for all $\theta \in \mathbb{R}$.
- vi) If x is a whole number the $x + 6 = 0$.

Q.3. Write the truth values of the following statements :

- i) $\sqrt{5}$ is an irrational but $3\sqrt{5}$ is a complex number.
- ii) $\forall n \in \mathbb{N}$, $n^2 + n$ is even number while $n^2 - n$ is an odd number.
- iii) $\exists n \in \mathbb{N}$ such that $n + 5 > 10$.
- iv) The square of any even number is odd or the cube of any odd number is odd.
- v) In $\triangle ABC$ if all sides are equal then its all angles are equal.
- vi) $\forall n \in \mathbb{N}$, $n + 6 > 8$.

Q.4. If $A = \{1, 2, 3, 4, 5, 6, 7, 8, 9\}$, determine the truth value of each of the following statement :

- i) $\exists x \in A$ such that $x + 8 = 15$.
- ii) $\forall x \in A$, $x + 5 < 12$.
- iii) $\exists x \in A$, such that $x + 7 \geq 11$.
- iv) $\forall x \in A$, $3x \leq 25$.

Q.5. Write the negations of the following :

- i) $\forall n \in A$, $n + 7 > 6$.
- ii) $\exists x \in A$, such that $x + 9 \leq 15$.
- iii) Some triangles are equilateral triangle.

Q.6. Construct the truth table for each of the following :

- i) $p \rightarrow (q \rightarrow p)$
- ii) $(\sim p \vee \sim q) \leftrightarrow [\sim(p \wedge q)]$
- iii) $\sim(\sim p \wedge \sim q) \vee q$
- iv) $[(p \wedge q) \vee r] \wedge [\sim r \vee (p \wedge q)]$
- v) $[(\sim p \vee q) \wedge (q \rightarrow r)] \rightarrow (p \rightarrow r)$

Q.7. Determine whether the following statement patterns are tautologies contradictions or contingencies :

- i) $[(p \rightarrow q) \wedge \sim q] \rightarrow \sim p$
- ii) $[(p \vee q) \wedge \sim p] \wedge \sim q$
- iii) $(p \rightarrow q) \wedge (p \wedge \sim q)$
- iv) $[p \rightarrow (q \rightarrow r)] \leftrightarrow [(p \wedge q) \rightarrow r]$
- v) $[(p \wedge (p \rightarrow q))] \rightarrow q$
- vi) $(p \wedge q) \vee (\sim p \wedge q) \vee (p \vee \sim q) \vee (\sim p \wedge \sim q)$
- vii) $[(p \vee \sim q) \vee (\sim p \wedge q)] \wedge r$
- viii) $(p \rightarrow q) \vee (q \rightarrow p)$

Q.8. Determine the truth values of p and q in the following cases :

- i) $(p \vee q)$ is T and $(p \wedge q)$ is T
- ii) $(p \vee q)$ is T and $(p \vee q) \rightarrow q$ is F
- iii) $(p \wedge q)$ is F and $(p \wedge q) \rightarrow q$ is T

Q.9. Using truth tables prove the following logical equivalences :

- i) $p \leftrightarrow q \equiv (p \wedge q) \vee (\sim p \wedge \sim q)$
- ii) $(p \wedge q) \rightarrow r \equiv p \rightarrow (q \rightarrow r)$

Q.10. Using rules in logic, prove the following :

- i) $p \leftrightarrow q \equiv \sim(p \wedge \sim q) \vee \sim(q \wedge \sim p)$
- ii) $\sim p \wedge q \equiv (p \vee q) \wedge \sim p$
- iii) $\sim(p \vee q) \vee (\sim p \wedge q) \equiv \sim p$

Q.11. Using the rules in logic, write the negations of the following :

- i) $(p \vee q) \wedge (q \vee \sim r)$
- ii) $p \wedge (q \vee r)$
- iii) $(p \rightarrow q) \wedge r$
- iv) $(\sim p \wedge q) \vee (p \wedge \sim q)$

Q.12. Express the following circuits in the symbolic form. Prepare the switching table :




Fig. 1.30




Fig. 1.31

Q.13. Simplify the following so that the new circuit has minimum number of switches. Also, draw the simplified circuit.




Fig. 1.32




Fig. 1.33

Q.14. Check whether the following switching circuits are logically equivalent - Justify.




Fig. 1.34

Fig. 1.35




Fig. 1.36

Fig. 1.37

Q.15. Give alternative arrangement of the switching following circuit, has minimum switches.




Fig. 1.38

Q.16 Simplify the following so that the new circuit circuit.




Fig. 1.39

Q.17. Represent the following switching circuit in symbolic form and construct its switching table. Write your conclusion from the switching table.




Fig. 1.40



2

Matrices



Let's Study.

- 2.1 Elementary transformations.
- 2.2 Inverse of a matrix
 - 2.2.1 Elementary transformation Method
 - 2.2.2 Adjoint method
- 2.3 Application of matrices .
 - Solution of a system of linear equations
 - 2.3.1 Method of Inversion
 - 2.3.2 Method of Reduction



Let's learn.

A matrix of order $m \times m$ is a square arrangement of m^2 elements. The corresponding determinant of the same elements, after expansion is seen to be a value which is an element itself.

In standard XI, we have studied the types of matrices and algebra of matrices namely addition, subtraction, multiplication of two matrices.

The matrices are useful in almost every branch of science. Many problems in Statistics are expressed in terms of matrices. Matrices are also useful in Economics, Operation Research. It would not be an exaggeration to say that the matrices are the language of atomic Physics.

Hence, it is necessary to learn the uses of matrices with the help of **elementary transformations** and the **inverse of a matrix**.

2.1 Elementary Transformation :

Let us first understand the meaning and applications of elementary transformations.

The elementary transformation of a matrix are the six operations, three of which are due to row and three are due to column.

They are as follows :

(a) **Interchange of any two rows or any two columns.** If we interchange the i^{th} row and the j^{th} row of a matrix then after this interchange the original matrix is transformed to a new matrix.

This transformation is symbolically denoted as $R_i \leftrightarrow R_j$ or R_{ij} .

The similar transformation can be due to two columns say $C_k \leftrightarrow C_i$ or C_{ki} .

(Recall that R and C symbolically represent the rows and columns of a matrix.)

For example, if $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ then $R_1 \leftrightarrow R_2$ gives the new matrix $\begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}$ and $C_1 \leftrightarrow C_2$ gives the new matrix $\begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}$.

Note that $A \neq \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}$ and $\neq \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}$ but we write $A \sim \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix}$ and $A \sim \begin{bmatrix} 2 & 1 \\ 4 & 3 \end{bmatrix}$

Note : The symbol \sim is read as equivalent to.

(b) Multiplication of the elements of any row or column by a non-zero scalar :

If k is a non-zero scalar and the row R_i is to be multiplied by constant k then we multiply every element of R_i by the constant k and symbolically the transformation is denoted by kR_i or $R_i \rightarrow kR_i$.

For example, if $A = \begin{bmatrix} 0 & 2 \\ 3 & 4 \end{bmatrix}$ then $R_2 \rightarrow 4R_2$ gives $A \sim \begin{bmatrix} 0 & 2 \\ 12 & 16 \end{bmatrix}$

Similarly, if any column of a matrix is to be multiplied by a constant then we multiply every element of the column by the constant. It is denoted as kC_i or $C_i \rightarrow kC_i$.

For example, if $A = \begin{bmatrix} 0 & 2 \\ 3 & 4 \end{bmatrix}$ then $C_1 \rightarrow -3 C_1$ gives $A \sim \begin{bmatrix} 0 & 2 \\ -9 & 4 \end{bmatrix}$

Can you say that $A = \begin{bmatrix} 0 & 2 \\ 12 & 16 \end{bmatrix}$ or $A = \begin{bmatrix} 0 & 2 \\ -9 & 4 \end{bmatrix}$

(c) Adding the scalar multiples of all the elements of any row (column) to corresponding elements of any other row (column).

If k is a non-zero scalar and the k -multiples of the elements of $R_i (C_j)$ are to be added to the elements of $R_j (C_i)$ then the transformation is symbolically denoted as $R_j \rightarrow R_j + kR_i, C_j \rightarrow C_j + kC_i$

For example, if $A = \begin{bmatrix} -1 & 4 \\ 2 & 5 \end{bmatrix}$ and $k = 2$ then $R_1 \rightarrow R_1 + 2R_2$ gives

$$A \sim \begin{bmatrix} -1+2(2) & 4+2(5) \\ 2 & 5 \end{bmatrix}$$

i.e. $A \sim \begin{bmatrix} 3 & 14 \\ 2 & 5 \end{bmatrix}$

(Can you find the transformation of A using $C_2 \rightarrow C_2 + (-3) C_1$?)

Note (1) : After the transformation, $R_j \rightarrow R_j + kR_i$, R_i remains the same as in the original matrix. Similarly, with the transformation, $C_j \rightarrow C_j + kC_i$, C_i remains the same as in the original matrix.

Note (2) : After the elementary transformation, the matrix obtained is said to be equivalent to the original matrix.

Ex. 1 : If $A = \begin{bmatrix} 1 & 0 \\ -1 & 3 \end{bmatrix}$, apply the transformation $R_1 \leftrightarrow R_2$ on A.

Solution :

$$\text{As } A = \begin{bmatrix} 1 & 0 \\ -1 & 3 \end{bmatrix}$$

$R_1 \leftrightarrow R_2$ gives

$$A \sim \begin{bmatrix} -1 & 3 \\ 1 & 0 \end{bmatrix}$$

Ex. 2 : If $A = \begin{bmatrix} 1 & 0 & 2 \\ 2 & 3 & 4 \end{bmatrix}$, apply the transformation $C_1 \rightarrow C_1 + 2C_3$.

Solution :

$$A = \begin{bmatrix} 1 & 0 & 2 \\ 2 & 3 & 4 \end{bmatrix}$$

$C_1 \rightarrow C_1 + 2C_3$ gives

$$A \sim \begin{bmatrix} 1+2(2) & 0 & 2 \\ 2+2(4) & 3 & 4 \end{bmatrix}$$

$$A \sim \begin{bmatrix} 5 & 0 & 2 \\ 10 & 3 & 4 \end{bmatrix}$$

Ex. 3 : If $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -2 & 5 \end{bmatrix}$, apply $R_1 \leftrightarrow R_2$ and then $C_1 \rightarrow C_1 + 2C_3$ on A.

Solution :

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & -2 & 5 \end{bmatrix}$$

$R_1 \leftrightarrow R_2$ gives

$$A \sim \begin{bmatrix} 3 & -2 & 5 \\ 1 & 2 & -1 \end{bmatrix}$$

Now $C_1 \rightarrow C_1 + 2C_3$ gives

$$A \sim \begin{bmatrix} 3+2(5) & -2 & 5 \\ 1+2(-1) & 2 & -1 \end{bmatrix} \quad \therefore A \sim \begin{bmatrix} 13 & -2 & 5 \\ -1 & 2 & -1 \end{bmatrix}$$



Exercise

Apply the given elementary transformation on each of the following matrices.

1. $A = \begin{bmatrix} 1 & 0 \\ -1 & 3 \end{bmatrix}$, $R_1 \leftrightarrow R_2$. 2. $B = \begin{bmatrix} 1 & -1 & 3 \\ 2 & 5 & 4 \end{bmatrix}$, $R_1 \rightarrow R_1 - R_2$.

3. $A = \begin{bmatrix} 5 & 4 \\ 1 & 3 \end{bmatrix}$, $C_1 \leftrightarrow C_2$; $B = \begin{bmatrix} 3 & 1 \\ 4 & 5 \end{bmatrix}$, $R_1 \leftrightarrow R_2$.

What do you observe?

4. $A = \begin{bmatrix} 1 & 2 & -1 \\ 0 & 1 & 3 \end{bmatrix}$, $2C_2$ $B = \begin{bmatrix} 1 & 0 & 2 \\ 2 & 4 & 5 \end{bmatrix}$, $-3R_1$.

Find the addition of the two new matrices.

5. $A = \begin{bmatrix} 1 & -1 & 3 \\ 2 & 1 & 0 \\ 3 & 3 & 1 \end{bmatrix}$, $3R_3$ and then $C_3 + 2C_2$.

6. $A = \begin{bmatrix} 1 & -1 & 3 \\ 2 & 1 & 0 \\ 3 & 3 & 1 \end{bmatrix}$, $C_3 + 2C_2$ and then $3R_3$.

What do you conclude from ex. 5 and ex. 6?

7. Use suitable transformation on $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ to convert it into an upper triangular matrix.

8. Convert $\begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$ into an identity matrix by suitable row transformations.

9. Transform $\begin{bmatrix} 1 & -1 & 2 \\ 2 & 1 & 3 \\ 3 & 2 & 4 \end{bmatrix}$ into an upper triangular matrix by suitable column transformations.

2.2 Inverse of a matrix :

Definition : If A is a square matrix of order m and if there exists another square matrix B of the same order such that $AB = BA = I$, where I is the identity matrix of order m, then B is called as the inverse of A and is denoted by A^{-1} .

Using the notation A^{-1} for B we get the above equation as $AA^{-1} = A^{-1}A = I$. Hence, using the same definition we can say that A is also the inverse of B.

$$\therefore B^{-1} = A$$

For example, if $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix}$ then $AB = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix}$

$$\therefore AB = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2 = I_2$$

$$\text{and } BA = \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2$$

$$\therefore B = A^{-1} \quad \text{and} \quad A = B^{-1}$$

If $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$, can you find a matrix X such that $AX = I$? Justify the answer.

This example illustrates that for the existence of such a matrix X, the necessary condition is $|A| \neq 0$, i.e. A is a non-singular matrix.

Note that -

- (1) Every square matrix A of order $m \times m$ has its corresponding determinant; $\det A = |A|$
- (2) A matrix is said to be invertible if its inverse exists.
- (3) A square matrix A has inverse if and only if $|A| \neq 0$

Uniqueness of inverse of a matrix

It can be proved that if A is a square matrix where $|A| \neq 0$ then its inverse, say A^{-1} , is unique.

Theorem : Prove that if A is a square matrix and its inverse exists then it is unique.

Proof : Let, 'A' be a square matrix of order 'm' and let its inverse exist.

Let, if possible, B and C be the two inverses of A.

Therefore, by definition of inverse $AB = BA = I$ and $AC = CA = I$.

$$\begin{aligned} \text{Now consider} \quad & B = BI = B(AC) \\ & \therefore B = (BA)C = IC \\ & \therefore B = C \end{aligned}$$

Hence $B = C$ i.e. the inverse is unique.

The inverse of a matrix (if it exists) can be obtained by using two methods.

- (i) Elementary row or column transformation
- (ii) Adjoint method

We now study these methods.

2.2.1 Inverse of a nonsingular matrix by elementary transformation :

By definition of inverse of A, if A^{-1} exists then $AA^{-1} = A^{-1}A = I$.

Let us consider the equation $AA^{-1} = I$. Here A is the given matrix of order m and I is the identity matrix of order 'm'. Hence the only unknown matrix is A^{-1} . Therefore, to find A^{-1} , we have to first convert A into I. This can be done by using elementary transformations.

Here we note that whenever any elementary row transformation is to be applied on the product $AB = C$ of two matrices A and B, it is enough to apply it only on the prefactor, A. B remains unchanged. And apply the same row transformation to C.

$$\text{For example, if } A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad \text{and } B = \begin{bmatrix} -1 & 0 \\ 1 & 5 \end{bmatrix} \quad \text{then } AB = \begin{bmatrix} 1 & 10 \\ 1 & 20 \end{bmatrix} = C \text{ (say)}$$

$$\text{Now if we require C to be transformed to a new matrix by } R_1 \leftrightarrow R_2 \quad \text{then } C \sim \begin{bmatrix} 1 & 20 \\ 1 & 10 \end{bmatrix}$$

$$\text{If the same transformation is used for A then } A \sim \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix} \quad \text{and B remains unchanged,}$$

$$\text{then the product } AB = \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 20 \\ 1 & 10 \end{bmatrix} = \text{as required. (Verify the product.)}$$

Hence, the equation $AA^{-1} = I$ can be transformed into an equation of the type $IA^{-1} = B$, by applying same series of row transformations on both the sides of the equation.

However, if we start with the equation $A^{-1}A = I$ (which is also true by the definition of inverse) then the transformation of A should be due to the column transformation. Apply column transformation to post factor and other side, where as prefactor remains unchanged.

Thus, starting with the equation $AA^{-1} = I$, we perform a series of row transformations on both sides of the equation, so that 'A' gets transformed to I. Thus,

$$\begin{array}{ccc} A & A^{-1} & = & I \\ \downarrow \text{Row} & & & \downarrow \text{Row} \\ \downarrow \text{Transformations} & & & \downarrow \text{Transformations} \\ I & A^{-1} & = & B \\ \therefore & A^{-1} & = & B \end{array}$$

and for the equation $A^{-1}A = I$, we use a series of column transformations. Thus

$$\begin{array}{rcc} A^{-1} A & = & I \\ \downarrow \text{Column} & & \downarrow \text{Column} \\ \downarrow \text{Transformation} & & \downarrow \text{Transformation} \\ A^{-1} I & = & B \\ \therefore A^{-1} & = & B \end{array}$$

Now if A is a given matrix of order '3' and it is nonsingular then we consider

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

For reducing the above matrix to

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ the suitable row transformations are as follows :}$$

- (1) Reduce a_{11} to '1'.
- (2) Then, reduce a_{21} and a_{31} to '0'.
- (3) Reduce a_{22} to '1'.
- (4) Then, reduce a_{12} and a_{32} to '0'.
- (5) Reduce a_{33} to '1'.
- (6) Then, reduce a_{13} and a_{23} to '0'.

Remember that a similar working rule (but not the same) can be used if you are using column transformations.



Solved Examples

Ex. 1 : Find which of the following matrices are invertible

$$(i) \quad A = \begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} \quad (ii) \quad B = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \quad (iii) \quad C = \begin{bmatrix} 1 & 3 & 2 \\ 3 & 1 & 2 \\ 1 & 2 & 3 \end{bmatrix}$$

Solution :

$$(i) \quad \text{As } |A| = \begin{vmatrix} 2 & 1 \\ 4 & 2 \end{vmatrix} = 0$$

\therefore A is singular and hence
A is not invertible.

$$(ii) \quad \text{As } |B| = \begin{vmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{vmatrix}$$

$$= \cos^2 \theta + \sin^2 \theta = 1 \neq 0$$

\therefore B is nonsingular,

\therefore B is invertible.

$$(iii) \quad C = \begin{bmatrix} 1 & 3 & 2 \\ 3 & 1 & 2 \\ 1 & 2 & 3 \end{bmatrix}$$

$$\therefore |C| = \begin{vmatrix} 1 & 3 & 2 \\ 3 & 1 & 2 \\ 1 & 2 & 3 \end{vmatrix} = -12 \neq 0$$

\therefore C is nonsingular and hence C is invertible.

Ex. 2 : Find the inverse of $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$.

Solution :

$$\text{As } |A| = \begin{vmatrix} 1 & 2 \\ 3 & 4 \end{vmatrix} = -2$$

$\therefore |A| \neq 0 \quad \therefore A^{-1}$ exists.

Let $AA^{-1} = I$ (Here we can use only row transformation)

Using $R_2 \rightarrow R_2 - 3R_1$

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ becomes}$$

$$\begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix}$$

Using $-\frac{1}{2}R_2$ we get

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix}$$

Using $R_1 \rightarrow R_1 - 2R_2$

$$\text{We get } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix} \text{ (Verify the answer.)}$$

Ex. 3 : Find the inverse of $A = \begin{bmatrix} 3 & 2 & 6 \\ 1 & 1 & 2 \\ 2 & 2 & 5 \end{bmatrix}$ by using elementary row transformations.

Solution : Consider $|A| = 1 \neq 0 \therefore A^{-1}$ exists.

Now as row transformations are to be used we have to consider the equation $AA^{-1} = I$ and have to perform row transformations on A.

Consider $AA^{-1} = I$

i.e. $\begin{bmatrix} 3 & 2 & 6 \\ 1 & 1 & 2 \\ 2 & 2 & 5 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Use $R_1 \leftrightarrow R_2$

i.e. $\begin{bmatrix} 1 & 1 & 2 \\ 3 & 2 & 6 \\ 2 & 2 & 5 \end{bmatrix} A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Use $R_2 \rightarrow R_2 - 3R_1$ and $R_3 \rightarrow R_3 - 2R_1$

$\therefore \begin{bmatrix} 1 & 1 & 2 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & -3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$

Now use $R_2 \rightarrow -R_2$

$\therefore \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$

Use $R_1 \rightarrow R_1 - R_2$

$\therefore \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & -2 & 0 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$

Use $R_1 \rightarrow R_1 - 2R_3$

$\therefore \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} A^{-1} = \begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$

$\therefore IA^{-1} = \begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix} \therefore A^{-1} = \begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$

(You can verify that $AA^{-1} = I$)

Ex. 4 : Find the inverse of $A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$ by elementary column transformation.

Solution :

As A^{-1} is required by column transformations therefore we have to consider $A^{-1}A = I$ and have to perform column transformations on A .

Consider

$$\begin{aligned} A^{-1}A &= I \\ \therefore A^{-1} \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

Using $C_2 \rightarrow C_2 - 3C_1$ and $C_3 \rightarrow C_3 - 3C_1$

$$\therefore A^{-1} \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -3 & -3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Use $C_1 \rightarrow C_1 - C_2$

$$\therefore A^{-1} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 4 & -3 & -3 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Use $C_1 \rightarrow C_1 - C_3$

$$\therefore A^{-1} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

$$\therefore A^{-1}I = \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

2.2.2 Inverse of a square matrix by adjoint method :

From the previous discussion about finding the inverse of a square matrix by elementary transformation it is clear that the method is elaborate and requires a series of transformations.

There is another method for finding the inverse and it is called as the inverse by the adjoint method. This method can be directly used for finding the inverse. However, for understanding this method you should know the definition of a minor, a co-factor and adjoint of the given matrix.

Let us first recall the definition of minor and co-factor of an element of a determinant.

Definition : Minor of an element a_{ij} of a determinant is the determinant obtained by deleting i^{th} row and j^{th} column in which the element a_{ij} lies. Minor of an element a_{ij} is denoted by M_{ij} .

(Can you find the order of the minor of any element of a determinant of order ‘n’?)

Definition : Co-factor of an element a_{ij} of a determinant is given by $(-1)^{i+j} M_{ij}$, where M_{ij} is minor of the element a_{ij} . Co-factor of an element a_{ij} is denoted by A_{ij} .

Now for defining the adjoint of a matrix, we require the co-factors of the elements of the matrix.

Consider a matrix $A = \begin{bmatrix} 1 & -2 & 3 \\ 4 & 5 & -6 \\ 7 & -8 & 9 \end{bmatrix}$. Its corresponding determinant is $|A| = \begin{vmatrix} 1 & -2 & 3 \\ 4 & 5 & -6 \\ 7 & -8 & 9 \end{vmatrix}$

Here if we require the minor of the element ‘4’, then it is $\begin{vmatrix} -2 & 3 \\ -8 & 9 \end{vmatrix} = -18+24 = 6$

Now as the element ‘4’ belongs to 2nd row and 1st column,

using the notation we get $M_{21} = 6$

If further we require the co-factor of ‘4’ then it is

$$= (-1)^{2+1} M_{21}$$

$$= (-1)(6)$$

$$= -6$$

Hence using notation, $A_{21} = -6$

Thus for any given matrix A, which is a square matrix, we can find the co-factors of all of its elements.

Definition :

The adjoint of a square matrix $A = [a_{ij}]_{m \times m}$ is defined as the transpose of the matrix $[A_{ij}]_{m \times m}$ where A_{ij} is the co-factor of the element a_{ij} of A, for all i and j , where $i, j = 1, 2, \dots, m$.

The adjoint of the matrix A is denoted by $\text{adj } A$.

For example, if A is a square matrix of order 3×3 then the matrix of its co-factors is

$$\begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix}$$

and the required adjoint of A is the transpose of the above matrix. Hence

$$\text{adj } A = \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix}$$

Ex. 1 : Find the co-factors of the elements of $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

Solution :

$$\begin{array}{llll} \text{Here } a_{11} = 1 & \therefore M_{11} = 4 & \text{and } A_{11} = (-1)^{1+1} (4) & = 4 \\ a_{12} = 2 & \therefore M_{12} = 3 & \text{and } A_{12} = (-1)^{1+2} (3) & = -3 \\ a_{21} = 3 & \therefore M_{21} = 2 & \text{and } A_{21} = (-1)^{2+1} (2) & = -2 \\ a_{22} = 4 & \therefore M_{22} = 1 & \text{and } A_{22} = (-1)^{2+2} (1) & = 1 \end{array}$$

\therefore the required co-factors are 4, -3, -2, 1.

Ex. 2 : Find the adjoint of matrix $A = \begin{bmatrix} 2 & -3 \\ 4 & 1 \end{bmatrix}$

Solution :

$$\begin{array}{llll} \text{Here } a_{11} = 2 & \therefore M_{11} = 1 & \therefore A_{11} = (-1)^{1+1} (1) = 1 \\ a_{12} = -3 & \therefore M_{12} = 4 & \therefore A_{12} = (-1)^{1+2} (4) = -4 \\ a_{21} = 4 & \therefore M_{21} = -3 & \therefore A_{21} = (-1)^{2+1} (-3) = 3 \\ a_{22} = 1 & \therefore M_{22} = 2 & \therefore A_{22} = (-1)^{2+2} (2) = 2 \end{array}$$

$$\therefore \text{ the matrix } [A_{ij}]_{2 \times 2} = \begin{bmatrix} 1 & -4 \\ 3 & 2 \end{bmatrix}$$

$$\therefore [A_{ij}]^T_{2 \times 2} = \begin{bmatrix} 1 & 3 \\ -4 & 2 \end{bmatrix}$$

$$\therefore \text{adj } A = \begin{bmatrix} 1 & 3 \\ -4 & 2 \end{bmatrix}$$

Ex. 3 : Find the adjoint of matrix $A = \begin{bmatrix} 2 & 0 & -1 \\ 3 & 1 & 2 \\ -1 & 1 & 2 \end{bmatrix}$

Solution :

$$\begin{array}{llll} \text{Here } a_{11} = 2 & \therefore M_{11} = 0 & \therefore A_{11} = (-1)^{1+1} M_{11} = 0 \\ a_{12} = 0 & \therefore M_{12} = 8 & \therefore A_{12} = (-1)^{1+2} M_{12} = -8 \end{array}$$

$$\begin{aligned}
 a_{13} = -1 & \quad \therefore M_{13} = 4 \\
 & \quad \therefore A_{13} = (-1)^{1+3} M_{13} = 4 \\
 a_{21} = 3 & \quad \therefore M_{21} = 1 \\
 & \quad \therefore A_{21} = (-1)^{2+1} M_{21} = -1 \\
 a_{22} = 1 & \quad \therefore M_{22} = 3 \\
 & \quad \therefore A_{22} = (-1)^{2+2} M_{22} = 3 \\
 a_{23} = 2 & \quad \therefore M_{23} = 2 \\
 & \quad \therefore A_{23} = (-1)^{2+3} M_{23} = -2 \\
 a_{31} = -1 & \quad \therefore M_{31} = 1 \\
 & \quad \therefore A_{31} = (-1)^{3+1} M_{31} = 1 \\
 a_{32} = 1 & \quad \therefore M_{32} = 7 \\
 & \quad \therefore A_{32} = (-1)^{3+2} M_{32} = -7 \\
 a_{33} = 2 & \quad \therefore M_{33} = 2 \\
 & \quad \therefore A_{33} = (-1)^{3+3} M_{33} = 2
 \end{aligned}$$

$$\therefore \text{the matrix of co-factors is } \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix} = \begin{bmatrix} 0 & -8 & 4 \\ -1 & 3 & -2 \\ 1 & -7 & 2 \end{bmatrix}$$

$$\therefore \text{adj } A = [A_{ij}]^T_{3 \times 3} = \begin{bmatrix} 0 & -1 & 1 \\ -8 & 3 & -7 \\ 4 & -2 & 2 \end{bmatrix}$$

We know that a determinant can be expanded with the help of any row. For example, expansion by 2nd row $a_{21} A_{21} + a_{22} A_{22} + \dots + a_{2n} A_{2n} = |A|$.

But if we multiply the row by a different row of cofactors, then the sum is zero.

For example, $a_{21} A_{31} + a_{22} A_{32} + \dots + a_{2n} A_{3n} = 0$

This helps us to prove that $A^{-1} = \frac{\text{adj } A}{|A|}$

$$\begin{aligned}
 \therefore A \cdot \text{adj } A &= \begin{vmatrix} |A| & 0 & 0 & \dots & 0 \\ 0 & |A| & 0 & \dots & 0 \\ \vdots & & & & \\ 0 & 0 & 0 & \dots & |A| \end{vmatrix} = |A| \cdot I \\
 \therefore A^{-1} &= \frac{\text{adj } A}{|A|}
 \end{aligned}$$

Thus, if $A = [a_{ij}]_{m \times m}$ is a non-singular square matrix then its inverse exists and it

is given by $A^{-1} = \frac{1}{|A|} (\text{adj } A)$

Think why A^{-1} does not exist if A is singular.

Ex. 1 : If $A = \begin{bmatrix} 2 & -2 \\ 4 & 3 \end{bmatrix}$, then find A^{-1} by the adjoint method.

Solution : For given matrix A, we get,

$$\begin{aligned} M_{11} &= 3, & A_{11} &= (-1)^{1+1} (3) = 3 \\ M_{12} &= 4, & A_{12} &= (-1)^{1+2} (4) = -4 \\ M_{21} &= -2, & A_{21} &= (-1)^{2+1} (-2) = 2 \\ M_{22} &= 2, & A_{22} &= (-1)^{2+2} (2) = 2 \end{aligned}$$

$$\therefore \text{adj } A = \begin{bmatrix} 3 & 2 \\ -4 & 2 \end{bmatrix}$$

$$\text{and } |A| = \begin{vmatrix} 2 & -2 \\ 4 & 3 \end{vmatrix} = 6 + 8 = 14 \neq 0$$

\therefore using $A^{-1} = \frac{1}{|A|} (\text{adj } A)$

$$A^{-1} = \frac{1}{14} \begin{bmatrix} 3 & 2 \\ -4 & 2 \end{bmatrix}$$

Ex. 2 : If $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$, find A^{-1} by the adjoint method.

Solution : For the given matrix A

$$A_{11} = (-1)^{1+1} \begin{vmatrix} 2 & -1 \\ -1 & 2 \end{vmatrix} = 3$$

$$A_{12} = (-1)^{1+2} \begin{vmatrix} -1 & -1 \\ 1 & 2 \end{vmatrix} = 1$$

$$A_{13} = (-1)^{1+3} \begin{vmatrix} -1 & 2 \\ 1 & -1 \end{vmatrix} = -1$$

$$A_{21} = (-1)^{2+1} \begin{vmatrix} -1 & 1 \\ -1 & 2 \end{vmatrix} = 1$$

$$A_{22} = (-1)^{2+2} \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} = 3$$

$$A_{23} = (-1)^{2+3} \begin{vmatrix} 2 & -1 \\ 1 & -1 \end{vmatrix} = 1$$

$$A_{31} = (-1)^{3+1} \begin{vmatrix} -1 & 1 \\ 2 & -1 \end{vmatrix} = -1$$

$$A_{32} = (-1)^{3+2} \begin{vmatrix} 2 & 1 \\ -1 & -1 \end{vmatrix} = 1$$

$$A_{33} = (-1)^{3+3} \begin{vmatrix} 2 & -1 \\ -1 & 2 \end{vmatrix} = 3$$

$$\therefore \text{adj } A = \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$$

$$\begin{aligned} \text{Now } |A| &= \begin{vmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{vmatrix} \\ &= 2(4-1) + 1(-2+1) + 1(1-2) \\ &= 6-1-1 \\ &= 4 \end{aligned}$$

Therefore by using the formula for A^{-1}

$$A^{-1} = \frac{1}{|A|} (\text{adj } A)$$

$$\therefore A^{-1} = \frac{1}{4} \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$$

Ex. 3 : If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, verify that $A(\text{adj } A) = (\text{adj } A)A = |A|I$.

Solution : For $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

$$A_{11} = (-1)^{1+1} (4) = 4$$

$$A_{12} = (-1)^{1+2} (3) = -3$$

$$A_{21} = (-1)^{2+1} (2) = -2$$

$$A_{22} = (-1)^{2+2} (1) = 1$$

$$\text{adj } A = \begin{bmatrix} A_{11} & A_{21} \\ A_{12} & A_{22} \end{bmatrix} = \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix}$$

$$\therefore A(\text{adj } A) = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix} = \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} \quad \dots \text{(i)}$$

$$\begin{aligned} (\text{adj } A) \cdot A &= \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 4-6 & 8-8 \\ -3+3 & -6+4 \end{bmatrix} \\ &= \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} \quad \dots \text{(ii)} \end{aligned}$$

$$\begin{aligned} \text{and } |A| I &= \begin{vmatrix} 1 & 2 \\ 3 & 4 \end{vmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &= (-2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} \quad \dots \text{(iii)} \end{aligned}$$

From (i), (ii) and (iii) we get, $A(\text{adj } A) = (\text{adj } A) A = |A| I$

(Note that this equation is valid for every nonsingular square matrix A)



Exercise

1. Find the co-factors of the elements of the following matrices

$$\text{(i)} \begin{bmatrix} -1 & 2 \\ -3 & 4 \end{bmatrix} \quad \text{(ii)} \begin{bmatrix} 1 & -1 & 2 \\ -2 & 3 & 5 \\ -2 & 0 & -1 \end{bmatrix}$$

2. Find the matrix of co-factors for the following matrices

$$\text{(i)} \begin{bmatrix} 1 & 3 \\ 4 & -1 \end{bmatrix} \quad \text{(ii)} \begin{bmatrix} 1 & 0 & 2 \\ -2 & 1 & 3 \\ 0 & 3 & -5 \end{bmatrix}$$

3. Find the adjoint of the following matrices.

$$\text{(i)} \begin{bmatrix} 2 & -3 \\ 3 & 5 \end{bmatrix} \quad \text{(ii)} \begin{bmatrix} 1 & -1 & 2 \\ -2 & 3 & 5 \\ -2 & 0 & -1 \end{bmatrix}$$

4. If $A = \begin{bmatrix} 1 & -1 & 2 \\ 3 & 0 & -2 \\ 1 & 0 & 3 \end{bmatrix}$

verify that $A(\text{adj } A) = (\text{adj } A) A = |A| I$

5. Find the inverse of the following matrices by the adjoint method.

(i) $\begin{bmatrix} -1 & 5 \\ -3 & 2 \end{bmatrix}$

(ii) $\begin{bmatrix} 2 & -2 \\ 4 & 3 \end{bmatrix}$

(iii) $\begin{bmatrix} 1 & 0 & 0 \\ 3 & 3 & 0 \\ 5 & 2 & -1 \end{bmatrix}$

(iv) $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & 5 \end{bmatrix}$

6. Find the inverse of the following matrices

(i) $\begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix}$

(ii) $\begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix}$

(iii) $\begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$

(iv) $\begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$

Miscellaneous exercise 2 (A)

1. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 3 & 1 \end{bmatrix}$ then reduce it to I_3 by using column transformations.

2. If $A = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ then reduce it to I_3 by using row transformations.

3. Check whether the following matrices are invertible or not

(i) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

(ii) $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

(iii) $\begin{bmatrix} 1 & 2 \\ 3 & 3 \end{bmatrix}$

(iv) $\begin{bmatrix} 2 & 3 \\ 10 & 15 \end{bmatrix}$

(v) $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$

(vi) $\begin{bmatrix} \sec \theta & \tan \theta \\ \tan \theta & \sec \theta \end{bmatrix}$

(vii) $\begin{bmatrix} 3 & 4 & 3 \\ 1 & 1 & 0 \\ 1 & 4 & 5 \end{bmatrix}$

(viii) $\begin{bmatrix} 1 & 2 & 3 \\ 2 & -1 & 3 \\ 1 & 2 & 3 \end{bmatrix}$

(ix) $\begin{bmatrix} 1 & 2 & 3 \\ 3 & 4 & 5 \\ 4 & 6 & 8 \end{bmatrix}$

4. Find AB , if $A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & -2 & -3 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -1 \\ 1 & 2 \\ 1 & -2 \end{bmatrix}$ Examine whether AB has inverse or not.

5. If $A = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix}$ is a nonsingular matrix then find A^{-1} by elementary row transformations.

Hence, find the inverse of $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$

6. If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and X is a 2×2 matrix such that $AX = I$, then find X.

7. Find the inverse of each of the following matrices (if they exist).

(i) $\begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}$ (ii) $\begin{bmatrix} 2 & 1 \\ 1 & -1 \end{bmatrix}$ (iii) $\begin{bmatrix} 1 & 3 \\ 2 & 7 \end{bmatrix}$ (iv) $\begin{bmatrix} 2 & -3 \\ 5 & 7 \end{bmatrix}$

(v) $\begin{bmatrix} 2 & 1 \\ 7 & 4 \end{bmatrix}$ (vi) $\begin{bmatrix} 3 & -10 \\ 2 & -7 \end{bmatrix}$ (vii) $\begin{bmatrix} 2 & -3 & 3 \\ 2 & 2 & 3 \\ 3 & -2 & 2 \end{bmatrix}$ (viii) $\begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix}$

(ix) $\begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$ (x) $\begin{bmatrix} 1 & 2 & -2 \\ 0 & -2 & 1 \\ -1 & 3 & 0 \end{bmatrix}$

8. Find the inverse of $A = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$

- by (i) elementary row transformations
(ii) elementary column transformations

9. If $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}$ find AB and $(AB)^{-1}$. Verify that $(AB)^{-1} = B^{-1}A^{-1}$

10. If $A = \begin{bmatrix} 4 & 5 \\ 2 & 1 \end{bmatrix}$, then show that $A^{-1} = \frac{1}{6} (A - 5I)$

11. Find matrix X such that $AX = B$, where

$A = \begin{bmatrix} 1 & 2 \\ -1 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ 2 & 4 \end{bmatrix}$

12. Find X, if $AX = B$ where

$A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$

13. If $A = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 1 \\ 3 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} 24 & 7 \\ 31 & 9 \end{bmatrix}$ then find matrix X such that $AXB = C$.

14. Find the inverse of $\begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$ by adjoint method.

15. Find the inverse of $\begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 3 \\ 1 & 2 & 1 \end{bmatrix}$ by adjoint method.

16. Find A^{-1} by adjoint method and by elementary transformations if $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 1 & 2 & 4 \end{bmatrix}$.

17. Find the inverse of $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 3 \\ 1 & 2 & 1 \end{bmatrix}$ by elementary column transformations.

18. Find the inverse of $\begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$ by elementary row transformations.

19. Show with usual notations that for any matrix $A = [a_{ij}]_{3 \times 3}$

(i) $a_{11}A_{21} + a_{12}A_{22} + a_{13}A_{23} = 0$

(ii) $a_{11}A_{11} + a_{12}A_{12} + a_{13}A_{13} = |A|$

20. If $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 3 \\ 1 & 2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 1 & 5 \\ 2 & 4 & 7 \end{bmatrix}$ then, find a matrix X such that $XA = B$.

2.3 Application of matrices :

In the previous discussion you have learnt the concept of inverse of a matrix. Now we intend to discuss the application of matrices for solving a system of linear equations.

For this we first learn to convert the given system of equations in the form of a matrix equation.

Consider the two linear equations, $2x + 3y = 5$ and $x - 4y = 9$. These equations can be written as shown below

$$\begin{bmatrix} 2x+3y \\ x-4y \end{bmatrix} = \begin{bmatrix} 5 \\ 9 \end{bmatrix}$$

(Recall the meaning of equality of two matrices.)

Now using the definition of multiplication of matrices we can consider the above equation as

$$\begin{bmatrix} 2 & 3 \\ 1 & -4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 9 \end{bmatrix}$$

Now if we denote $\begin{bmatrix} 2 & 3 \\ 1 & -4 \end{bmatrix} = A$, $\begin{bmatrix} x \\ y \end{bmatrix} = X$ and $\begin{bmatrix} 5 \\ 9 \end{bmatrix} = B$

then the above equation can be written as $AX = B$

In the equation $AX = B$, X is the column matrix of variables, A is the matrix of coefficients of variables and B is the column matrix of constants.

Note that if A is of order 2×2 , X is of order 2×1 , then B is of order 2×1 .

Similarly, if there are three linear equations in three variables then as shown above they can be expressed as $AX = B$.

Find the respective orders of the matrices A , X and B in case of three equations in three variables.

This matrix equation $AX = B$ (in both the cases) can be used to find the values of the variables x and y or x , y and z as the case may be. There are two methods for this application which are namely

- (i) method of inversion
- (ii) method of reduction

2.3.1 Method of inversion :

From the name of this method you can guess that here we are going to use the inverse of a matrix.

This can be done as follows :

Consider the three equations as

$$\begin{aligned} a_1x + b_1y + c_1z &= d_1 \\ a_2x + b_2y + c_2z &= d_2 \\ a_3x + b_3y + c_3z &= d_3 \end{aligned}$$

As explained in the beginning, they can be expressed as

$$\begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix} \quad \text{i.e. } AX = B.$$

Observe that the respective orders of A , X and B are 3×3 , 3×1 and 3×1 .

Now, if the solution of the three equations exists, then the matrix A must be nonsingular. Hence, A^{-1} exists. Therefore, A^{-1} can be found out either by transformation method or by adjoint method.

After finding A^{-1} , pre-multiply the matrix equation $AX = B$ by A^{-1}

Thus we get,

$$\begin{aligned} A^{-1}(AX) &= A^{-1}(B) \\ \text{i.e. } (A^{-1}A)X &= A^{-1}B \\ \text{i.e. } IX &= A^{-1}B \\ \text{i.e. } X &= A^{-1}B \quad \text{which gives the required solution.} \end{aligned}$$



Solved Examples

Ex. 1 : Solve the equations $2x + 5y = 1$ and $3x + 2y = 7$ by the method of inversion.

Solution : Using the given equations we get the corresponding matrix equation as

$$\begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 7 \end{bmatrix}$$

i.e. $AX = B$, where $A = \begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \end{bmatrix}$ and $B = \begin{bmatrix} 1 \\ 7 \end{bmatrix}$

Hence, premultiplying the above matrix equation by A^{-1} , we get

$$(A^{-1}A)X = A^{-1}B$$

i.e. $IX = A^{-1}B$

i.e. $X = A^{-1}B$ (i)

Now as $A = \begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix}$, $|A| = -11$ and $\text{adj } A = \begin{bmatrix} 2 & -5 \\ -3 & 2 \end{bmatrix}$

$$\therefore A^{-1} = \frac{1}{|A|} (\text{adj } A)$$

i.e. $A^{-1} = \frac{1}{-11} \begin{bmatrix} 2 & -5 \\ -3 & 2 \end{bmatrix}$

Hence using (i) we get

$$X = \frac{1}{-11} \begin{bmatrix} 2 & -5 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \end{bmatrix}$$

$$X = \frac{1}{11} \begin{bmatrix} -2 & 5 \\ 3 & -2 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \end{bmatrix}$$

i.e. $\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 33 \\ -11 \end{bmatrix}$

i.e. $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$

Hence by equality of matrices we get $x = 3$ and $y = -1$.

Ex. 2 : Solve the following equations by the method of inversion

$$x - y + z = 4, 2x + y - 3z = 0, x + y + z = 2.$$

Solution : The required matrix equation is $\begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix}$ i.e. $AX = B$

Hence, by premultiplying the equation by A^{-1} , we get,

i.e. $(A^{-1}A)X = A^{-1}B$

i.e. $IX = A^{-1}B$

i.e. $X = A^{-1}B$ (i)

$$\text{Now as } A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix}, \text{ By definition, } \text{adj } A = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix} \text{ and } |A| = 10$$

$$A^{-1} = \frac{1}{|A|} (\text{adj } A)$$

$$\text{i.e. } A^{-1} = \frac{1}{10} \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix}$$

Hence using (i)

$$X = \frac{1}{10} \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix}$$

$$X = \frac{1}{10} \begin{bmatrix} 20 \\ -10 \\ 10 \end{bmatrix}$$

$$\therefore \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$$

Hence, by equality of matrices we get $x = 2, y = -1$ and $z = 1$

2.3.2 Method of reduction :

From the name of the method, it can be guessed that, the given equations are reduced to a certain form to get the solution.

Here also, we start by converting the given linear equation into matrix equation $AX = B$.

Then we perform the suitable row transformations on the matrix A.

Using the row transformations on A reduce it to an upper triangular matrix or lower triangular matrix or diagonal matrix.

The same row transformations are performed simultaneously on matrix B.

After this step we rewrite the equation in the form of system of linear equations. Now they are in such a form that they can be easily solved by elimination method. Thus, the required solution is obtained.



Solved Examples

Ex. 1 : Solve the equation $2x + 3y = 9$ and $y - x = -2$ using the method of reduction.

Solution : The given equations can be written as

$$\begin{aligned} 2x + 3y &= 9 \\ \text{and } -x + y &= -2 \end{aligned}$$

Hence the matrix equation is
$$\begin{bmatrix} 2 & 3 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \end{bmatrix} \text{ (i.e. } AX = B)$$

Now use $R_2 \rightarrow 2R_2 + R_1$

\therefore We get
$$\begin{bmatrix} 2 & 3 \\ 0 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \end{bmatrix}$$

We rewrite the equations as
$$2x + 3y = 9 \quad \dots\dots \text{ (i)}$$

$$5y = 5 \quad \dots\dots \text{ (ii)}$$

From (ii) $y = 1$ and using (i) we get $x = 3$

$\therefore x = 3, y = 1$ is the required solution.

Ex. 2 : Solve the following equations by the method of reduction.

$x + 3y + 3z = 12, \quad x + 4y + 4z = 15 \quad \text{and} \quad x + 3y + 4z = 13.$

Solution : The above equations can be written in the form $AX = B$

i.e.
$$\begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 4 \\ 1 & 3 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 13 \end{bmatrix}$$

using $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$

we get
$$\begin{bmatrix} 1 & 3 & 3 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 12 \\ 3 \\ 1 \end{bmatrix}$$

Again using $R_1 \rightarrow R_1 - 3R_2$ and $R_2 \rightarrow R_2 - R_3$ We get
$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$$

Hence the required solution is $x = 3, y = 2, z = 1.$ (verify)

Ex. 3 : Solve the following equations by the method of reduction.

$x + y + z = 1, \quad 2x + 3y + 2z = 2 \quad \text{and} \quad x + y + 2z = 4.$

Solution : The above equation can be written in the form $AX = B$ as

$$\begin{bmatrix} 1 & 1 & 1 \\ 2 & 3 & 2 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 4 \end{bmatrix}$$

using $R_2 \rightarrow R_2 - R_3$ and $R_1 \rightarrow R_1 - \frac{1}{2} R_3$

we get
$$\begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ 1 & 2 & 0 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \\ 4 \end{bmatrix}$$

Now using $R_1 \rightarrow R_1 - \frac{1}{4} R_2$ we get
$$\begin{bmatrix} \frac{1}{4} & 0 & 0 \\ 1 & 2 & 0 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ -2 \\ 4 \end{bmatrix}$$

Note that here we have reduced the original matrix A to a lower triangular matrix. Hence we can rewrite the equations in their original form as

$$\begin{aligned} \frac{x}{4} &= -\frac{1}{2} && \dots\dots\dots(i) && \text{i.e. } x = -2 \\ x + 2y &= -2 && \dots\dots\dots(ii) && \\ \therefore 2y &= -2 + 2 = 0 && && \therefore y = 0 \\ \text{and } x + y + 2z &= 4 && && \\ \therefore 2z &= 4 + 2 + 0 && && \\ \therefore 2z &= 6 && && \\ \therefore z &= 3 && && \\ \therefore x = -2, y = 0, z = 3 &\text{ is the required solution.} && && \end{aligned}$$

Ex. 4 : The cost of 2 books and 6 note books is Rs. 34 and the cost of 3 books and 4 notebooks is Rs. 31.

Using matrices, find the cost of one book and one note-book.

Solution : Let Rs. 'x' and ` Rs. 'y' be the costs of one book and one notebook respectively.

Hence, using the above information we get the following equations

$$\begin{aligned} 2x + 6y &= 34 \\ \text{and } 3x + 4y &= 31 \end{aligned}$$

The above equations can be expressed in the form

$$\begin{bmatrix} 2 & 6 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 34 \\ 31 \end{bmatrix} \quad \text{i.e. } AX = B$$


Now using $R_2 \rightarrow R_2 - \frac{3}{2} R_1$ we get

$$\begin{bmatrix} 2 & 6 \\ 0 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 34 \\ -20 \end{bmatrix}$$

As the above matrix 'A' is reduced to an upper triangular matrix, we can write the equations in their original form as $2x + 6y = 34$

$$\begin{aligned} \text{and } -5y &= -20 && \therefore y = 4 \\ \text{and } 2x &= 34 - 6y = 34 - 24 && \therefore 2x = 10 && \therefore x = 5 \end{aligned}$$

\therefore the cost of a book is Rs. 5 and that of a note book is Rs. 4.

 **Exercise 2.3**

1. Solve the following equations by inversion method.

- (i) $x + 2y = 2, \quad 2x + 3y = 3$
- (ii) $x + y = 4, \quad 2x - y = 5$
- (iii) $2x + 6y = 8, \quad x + 3y = 5$

2. Solve the following equations by reduction method.
- (i) $2x + y = 5, \quad 3x + 5y = -3$
(ii) $x + 3y = 2, \quad 3x + 5y = 4$
(iii) $3x - y = 1, \quad 4x + y = 6$
(iv) $5x + 2y = 4, \quad 7x + 3y = 5$
3. The cost of 4 pencils, 3 pens and 2 erasers is Rs. 60. The cost of 2 pencils, 4 pens and 6 erasers is Rs. 90, whereas the cost of 6 pencils, 2 pens and 3 erasers is Rs.70. Find the cost of each item by using matrices.
4. If three numbers are added, their sum is '2'. If 2 times the second number is subtracted from the sum of first and third number we get '8' and if three times the first number is added to the sum of second and third number we get '4'. Find the numbers using matrices.
5. The total cost of 3 T.V. sets and 2 V.C.R.s is Rs. 35000. The shop-keeper wants profit of 1000 per television and Rs. 500 per V.C.R. He can sell 2 T. V. sets and 1 V.C.R. and get the total revenue as Rs. 21,500. Find the cost price and the selling price of a T.V. sets and a V.C.R.



Let's Remember :

- If $A = [a_{ij}]_{m \times n}$ then A' or $A^T = [a_{ji}]_{n \times m}$
- If (i) A is symmetric then $A = A^T$ and (ii) if A is skew-symmetric then $-A = A^T$
- If A is a non singular matrix then $A^{-1} = \frac{1}{|A|}(\text{adj } A)$
- If A, B and C are three matrices of the same order then
 - (i) $A + B = B + A$ (Commutative law of addition)
 - (ii) $(A + B) + C = A + (B + C)$ (Associative law for addition)
- If A, B and C are three matrices of appropriate orders so that the following products are defined then
 - (i) $(AB)C = A(BC)$ (Associative Law of multiplication)
 - (ii) $A(B + C) = AB + AC$ (Left Distributive Law)
 - (iii) $(A + B)C = AC + BC$ (Right Distributive Law)
- The three types of elementary transformations are denoted as
 - (i) $R_i \leftrightarrow R_j$ or $C_i \leftrightarrow C_j$
 - (ii) $R_i \rightarrow kR_j$ or $C_i \rightarrow kC_j$ (k is a scalar), $k \neq 0$
 - (iii) $R_i \rightarrow R_i + kR_j$ or $C_i \rightarrow C_i + kC_j$ (k is a scalar), $k \neq 0$
- If A and B are two square matrices of the same order such that $AB = BA = I$, then A and B are inverses of each other. A is denoted as B^{-1} and B is denoted as A^{-1} .
- For finding the inverse of A , if row transformations are to be used then we consider $AA^{-1} = I$ and if column transformations are to be used then we consider $A^{-1}A = I$.

- For finding the inverse of any nonsingular square matrix, two methods can be used
(i) Elementary transformation method (ii) Adjoint method.
- Using matrices, a system of linear equations can be solved. The two methods are
(i) Method of inversion (ii) Method of reduction.

• If A is a non singular matrix then $A^{-1} = \frac{1}{|A|} (\text{adj } A)$

• A diagonal matrix is invertible iff all the elements in the diagonal are non - zero.

• The inverse of a diagonal matrix is a diagonal matrix.

• If A is a non singular matrix then $\det A^{-1} = \frac{1}{|A|}$

• The adjoint of a square matrix $A = [a_{ij}]_{m \times m}$ is the transpose of the matrix $[A_{ij}]_{m \times m}$, where A_{ij} is the co-factor of the element a_{ij} of A .

• Every square matrix A can be expressed as the sum of symmetric and skew-symmetric matrix. i.e. $A = \frac{1}{2} [A + A^T] + \frac{1}{2} [A - A^T]$

Miscellaneous Exercise 2(B)

1) Choose the correct alternative.

If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $\text{adj } A = \begin{bmatrix} 4 & a \\ -3 & b \end{bmatrix}$ then the values of a and b are,

A) $a = -2$, $b = 1$

B) $a = 2$, $b = 4$

C) $a = 2$, $b = -1$

D) $a = 1$, $b = -2$

2) The inverse of $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ is

A) $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

B) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

C) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

D) None of these

3) If $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $A(\text{adj}A) = kI$ then the value of k is

A) 1

B) -1

C) 0

D) -3

4) If $A = \begin{bmatrix} 2 & -4 \\ 3 & 1 \end{bmatrix}$ then the adjoint of matrix A is

A) $\begin{bmatrix} -1 & 3 \\ -4 & 1 \end{bmatrix}$

B) $\begin{bmatrix} 1 & 4 \\ -3 & 2 \end{bmatrix}$

C) $\begin{bmatrix} 1 & 3 \\ 4 & -2 \end{bmatrix}$

D) $\begin{bmatrix} -1 & -3 \\ -4 & 2 \end{bmatrix}$

5) If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $A(\text{adj } A) = k I$ then the value of k is

A) 2

B) -2

C) 10

D) -10

6) If $A = \begin{bmatrix} \lambda & 1 \\ -1 & -\lambda \end{bmatrix}$ then A^{-1} does not exist if $\lambda =$

A) 0

B) ± 1

C) 2

D) 3

7) If $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$ then $A^{-1} =$

A) $\begin{bmatrix} 1/\cos \alpha & -1/\sin \alpha \\ 1/\sin \alpha & 1/\cos \alpha \end{bmatrix}$

B) $\begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$

C) $\begin{bmatrix} -\cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$

D) $\begin{bmatrix} -\cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{bmatrix}$

8) If $F(\alpha) = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$ where $\alpha \in \mathbb{R}$ then $[F(\alpha)]^{-1}$ is =

A) $F(-\alpha)$

B) $F(\alpha^{-1})$

C) $F(2\alpha)$

D) None of these

9) The inverse of $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ is

A) I

B) A

C) A'

D) $-I$

10) The inverse of a symmetric matrix is -

A) Symmetric

B) Non-symmetric

C) Null matrix

D) Diagonal matrix

11) For a 2×2 matrix A , if $A(\text{adj}A) = \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix}$ then determinant A equals

- A) 20
B) 10
C) 30
D) 40

12) If $A' = -\frac{1}{2} \begin{bmatrix} 1 & -4 \\ -1 & 2 \end{bmatrix}$ then $A =$

- a) $\begin{bmatrix} 2 & 4 \\ -1 & 1 \end{bmatrix}$
b) $\begin{bmatrix} 2 & 4 \\ 1 & -1 \end{bmatrix}$
c) $\begin{bmatrix} 2 & -4 \\ 1 & 1 \end{bmatrix}$
d) $\begin{bmatrix} 2 & 4 \\ 1 & 1 \end{bmatrix}$

II) 1) Solve the following equations by the methods of inversion.

- (i) $2x - y = -2$
(ii) $x + y + z = 1$, $2x + 3y + 2z = 2$ and $ax + ay + 2az = 4$
(iii) $5x - y + 4z = 5$, $2x + 3y + 5z = 2$ and $5z - 2y + 6z = -1$
(iv) $2x + 3y = -5$, $3d + y = 3$
(v) $x + y + z = -1$, $y + z = 2$ and $x + y - z = 3$

2) Express the following equation in matrix form and solve them by the method of reduction.

- (i) $x - y + z = 1$, $2x - y = 1$, $3x + 3y - 4z = 2$
(ii) $x + y = 1$, $y + x =$
(iii) $2x - y + z = 1$, $x + 2y + 3z = 8$ and $3x + y - 4z = 1$
(iv) $x + y + z = 6$, $3x - y + 3z = 10$ and $5z + 5y - 4z = 3$
(v) $x + 2y + z = 8$, $2x + 3y - z = 1$ and $3x - y - 2z = 5$
(vi) $x + 3y + 2z = 6$, $3x - 2y + 5z = 5$ and $2x - 3y + 6z = 7$

3) The sum of three numbers is 6. If we multiply third number by 3 and add it to the second number we get 11.

By adding first and the third numbers we get a number which is double the second number. Use this information and find a system of linear equations. Find the three numbers using matrices.

- 4) The cost of 4 pencils, 3 pens and 2 books is Rs.150. The cost of 1 pencil, 2 pens and 3 books is Rs.125. The cost of 6 pencils, 2 pens and 3 books is Rs.175. Find the cost of each item by using Matrices.
- 5) The sum of three numbers is 6. Thrice the third number when added to the first number gives 7.

On adding three times first number to the sum of second and third number we get 12. Find the three numbers by using Matrices.

- 6) The sum of three numbers is 2. If twice the second number is added to the sum of first and third number, we get 0 adding five times the first number to the sum of second and third we get 6. Find the three numbers by using matrices.
- 7) An amount of Rs.5000 is invested in three types of investments, at interest rates 6.7, 7.7, 8% per annum respectively. The total annual income from these investments is Rs.350/- If the total annual income from first two investments is Rs.70 more than the income from the third, find the amount of each investment using matrix method.
- 8) The sum of the costs of one book each of Mathematics, Physics and Chemistry is Rs.210. Total cost of a mathematics book, 2 physics books, and a chemistry book is Rs. 240/- Also the total cost of a Mathematics book, 3 physics book and chemistry books is Rs. 300/-. Find the cost of each book, using Matrices.



3

Trigonometric Functions



Let's Study

3.1 Trigonometric Equations and their solutions

3.2 Solutions of triangle

3.2.1 Polar co-ordinates

3.2.2 Relation between the polar co-ordinates and the Cartesian co-ordinates

3.2.3 Solving a Triangle

3.2.4 The Sine rule

3.2.5 The Cosine rule

3.2.6 The Projection rule

3.2.7 Applications of the Sine rule, the Cosine rule and the Projection rule.

3.3 Inverse Trigonometric Functions

Properties, Principal values of inverse trigonometric functions

INTRODUCTION :

We are familiar with algebraic equations. In this chapter we will learn how to solve trigonometric equations, their principal and general solutions, their properties. Trigonometric functions play an important role in integral calculus.



Let's learn.

3.1 Trigonometric Equations and their solutions:

Trigonometric equation :

Definition : An equation involving trigonometric function (or functions) is called trigonometric equation.

For example : $\sin\theta = \frac{1}{2}$, $\tan\theta = 2$, $\cos 3\theta = \cos 5\theta$ are all trigonometric equations, $x = a \sin(\omega t + \alpha)$

is also a trigonometric equation.

Solution of Trigonometric equation :

Definition : A value of a variable in a trigonometric equation which satisfies the equation is called a solution of the trigonometric equation.

A trigonometric equation can have more than one solutions.

For example, $\theta = \frac{\pi}{6}$ satisfies the equation, $\sin \theta = \frac{1}{2}$, Therefore $\frac{\pi}{6}$ is a solution of the trigonometric equation $\sin \theta = \frac{1}{2}$,

$\frac{\pi}{4}$ is a solution of the trigonometric equation $\cos \theta = \frac{1}{\sqrt{2}}$.

$\frac{7\pi}{4}$ is a solution of the trigonometric equation $\cos \theta = \frac{1}{\sqrt{2}}$.

Is π a solution of equation $\sin \theta - \cos \theta = 1$? Can you write one more solution of this equation? Equation $\sin \theta = 3$ has no solution. Can you justify it?

Because of periodicity of trigonometric functions, trigonometric equation may have infinite number of solutions. Our interest is in finding solutions in the interval $[0, 2\pi)$.

Principal Solutions :

Definition : A solution α of a trigonometric equation is called a principal solution if $0 \leq \alpha < 2\pi$.

$\frac{\pi}{6}$ and $\frac{5\pi}{6}$ are the principal solutions of trigonometric equation $\sin \theta = \frac{1}{2}$.

Note that $\frac{13\pi}{6}$ is a solution but not principal solution of $\sin \theta = \frac{1}{2}$, $\left\{ \because \frac{13\pi}{6} \notin [0, 2\pi) \right\}$

0 is the principal solution of equation $\sin \theta = 0$ but 2π is not a principal solution.

Trigonometric equation $\cos \theta = -1$ has only one principal solution. $\theta = \pi$ is the only principal solution of this equation.



Solved Examples

Ex. (1) Find the principal solutions of $\sin \theta = \frac{1}{\sqrt{2}}$.

Solution :

As $\sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}$ and $0 \leq \frac{\pi}{4} < 2\pi$, $\frac{\pi}{4}$ is a principal solution.

By allied angle formula, $\sin \theta = \sin (\pi - \theta)$.

$\therefore \sin \frac{\pi}{4} = \sin \left(\pi - \frac{\pi}{4} \right) = \sin \frac{3\pi}{4}$ and $0 \leq \frac{3\pi}{4} < 2\pi$

$\therefore \frac{3\pi}{4}$ is also a principal solution.

$\therefore \frac{\pi}{4}$ and $\frac{3\pi}{4}$ are the principal solutions of $\sin \theta = \frac{1}{\sqrt{2}}$.

Ex.(2) Find the principal solutions of $\cos \theta = \frac{1}{2}$.

Solution : As $\cos \frac{\pi}{3} = \frac{1}{2}$ and $0 \leq \frac{\pi}{3} < 2\pi$, $\frac{\pi}{3}$ is a principal solution.

By allied angle formula, $\cos \theta = \cos(2\pi - \theta)$.

$$\therefore \cos \frac{\pi}{3} = \cos \left(2\pi - \frac{\pi}{3} \right) = \cos \frac{5\pi}{3} \text{ and } 0 \leq \frac{5\pi}{3} < 2\pi$$

$\therefore \frac{5\pi}{3}$ is also a principal solution.

$\therefore \frac{\pi}{3}$ and $\frac{5\pi}{3}$ are the principal solutions of $\cos \theta = \frac{1}{2}$.

Ex. (3) Find the principal solutions of $\cos \theta = -\frac{1}{2}$

Solution : We know that $\cos \frac{\pi}{3} = \frac{1}{2}$

As $\cos(\pi - \theta) = \cos(\pi + \theta) = -\cos \theta$,

$$\cos \left(\pi - \frac{\pi}{3} \right) = -\cos \frac{\pi}{3} = -\frac{1}{2} \text{ and } \cos \left(\pi + \frac{\pi}{3} \right) = -\cos \frac{\pi}{3} = -\frac{1}{2}$$

$$\therefore \cos \frac{2\pi}{3} = -\frac{1}{2} \text{ and } \cos \frac{4\pi}{3} = -\frac{1}{2}$$

Also $0 \leq \frac{2\pi}{3} \leq 2\pi$ and $0 \leq \frac{4\pi}{3} < 2\pi$. Therefore $\frac{2\pi}{3}$ and $\frac{4\pi}{3}$ are principal solutions of $\cos \theta = -\frac{1}{2}$

Ex.(4) Find the principal solutions of $\cot \theta = -\sqrt{3}$

Solution : We know that $\cot \theta = -\sqrt{3}$ if and only if $\tan \theta = -\frac{1}{\sqrt{3}}$

We know that $\tan \frac{\pi}{6} = \frac{1}{\sqrt{3}}$

Using identities, $\tan(\pi - \theta) = -\tan \theta$ and $\tan(2\pi - \theta) = -\tan \theta$, we get

$$\tan \frac{5\pi}{6} = -\frac{1}{\sqrt{3}} \text{ and } \tan \frac{11\pi}{6} = -\frac{1}{\sqrt{3}}$$

$$\text{An } 0 \leq \frac{5\pi}{6} < 2\pi \text{ and } 0 \leq \frac{11\pi}{6} < 2\pi$$

$\therefore \frac{5\pi}{6}$ and $\frac{11\pi}{6}$ are required principal solutions.

The General Solution :

Definition : The solution of a trigonometric equation which is generalized by using its periodicity is called the general solution

For example : All solutions of the equation $\sin \theta = \frac{1}{2}$, are $\left\{ \dots, -\frac{7\pi}{6}, \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}, \dots \right\}$. We can generate all these solutions from the expression $n\pi + (-1)^n \frac{\pi}{6}$, $n \in \mathbb{Z}$. The solution $n\pi + (-1)^n \frac{\pi}{6}$, $n \in \mathbb{Z}$ is called the general solution of $\sin \theta = \frac{1}{2}$.

Theorem 3.1 : The general solution of $\sin \theta = \sin \alpha$ is $\theta = n\pi + (-1)^n \alpha$, where $n \in \mathbb{Z}$.

Proof : As $\sin \theta = \sin \alpha$, α is a solution.

As $\sin(\pi - \alpha) = \sin \alpha$, $\pi - \alpha$ is also a solution. Using periodically, we get

$$\sin \theta = \sin \alpha = \sin(2\pi + \alpha) = \sin(4\pi + \alpha) = \dots \text{ and}$$

$$\sin \theta = \sin(\pi - \alpha) = \sin(3\pi - \alpha) = \sin(5\pi - \alpha) = \dots$$

$$\therefore \sin \theta = \sin \alpha \text{ if and only if } \theta = \alpha, 2\pi + \alpha, 4\pi + \alpha, \dots \text{ or } \theta = \pi - \alpha, 3\pi - \alpha, 5\pi - \alpha, \dots$$

$$\therefore \theta = \dots, \alpha, \pi - \alpha, 2\pi + \alpha, 3\pi - \alpha, 4\pi + \alpha, 5\pi - \alpha, \dots$$

$$\therefore \text{The general solution of } \sin \theta = \sin \alpha \text{ is } \theta = n\pi + (-1)^n \alpha, \text{ where } n \in \mathbb{Z}.$$

Theorem 3.2 : The general solution of $\cos \theta = \cos \alpha$ is $\theta = 2n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Proof : As $\cos \theta = \cos \alpha$, α is a solution.

As $\cos(-\alpha) = \cos \alpha$, $-\alpha$ is also a solution.

Using periodically, we get

$$\cos \theta = \cos \alpha = \cos(2\pi + \alpha) = \cos(4\pi + \alpha) = \dots \text{ and}$$

$$\cos \theta = \cos(-\alpha) = \cos(2\pi - \alpha) = \cos(4\pi - \alpha) = \dots$$

$$\therefore \cos \theta = \cos \alpha \text{ if and only if } \theta = \alpha, 2\pi + \alpha, 4\pi + \alpha, \dots \text{ or } \theta = -\alpha, 2\pi - \alpha, 4\pi - \alpha, 6\pi - \alpha, \dots$$

$$\therefore \text{The general solution of } \cos \theta = \cos \alpha \text{ is } \theta = 2n\pi \pm \alpha, \text{ where } n \in \mathbb{Z}.$$

Theorem 3.3 : The general solution of $\tan \theta = \tan \alpha$ is $\theta = n\pi + \alpha$, where $n \in \mathbb{Z}$.

Proof : We know that $\tan \theta = \tan \alpha$ if and only if $\frac{\sin \theta}{\cos \theta} = \frac{\sin \alpha}{\cos \alpha}$

$$\text{If and only if } \sin \theta \cos \alpha = \cos \theta \sin \alpha$$

$$\text{If and only if } \sin \theta \cos \alpha - \cos \theta \sin \alpha = 0$$

$$\text{If and only if } \sin(\theta - \alpha) = \sin 0$$

$$\text{If and only if } \theta - \alpha = n\pi + (-1)^n \times 0 = n\pi, \text{ where } n \in \mathbb{Z}.$$

$$\text{If and only if } \theta = n\pi + \alpha, \text{ where } n \in \mathbb{Z}.$$

$$\text{The general solution of } \tan \theta = \tan \alpha \text{ is } \theta = n\pi + \alpha, \text{ where } n \in \mathbb{Z}.$$

Remark : For $\theta \in \mathbb{R}$, we have the following :

(i) $\sin \theta = 0$ if and only if $\theta = n\pi$, where $n \in \mathbb{Z}$.

(ii) $\cos \theta = 0$ if and only if $\theta = (2n+1)\frac{\pi}{2}$, where $n \in \mathbb{Z}$.

(iii) $\tan \theta = 0$ if and only if $\theta = n\pi$, where $n \in \mathbb{Z}$.

Theorem 3.4 : The general solution of $\sin^2 \theta = \sin^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Proof : $\sin^2 \theta = \sin^2 \alpha$

- $\therefore \sin \theta = \pm \sin \alpha$
- $\therefore \sin \theta = \sin \alpha$ or $\sin \theta = -\sin \alpha$
- $\therefore \sin \theta = \sin \alpha$ or $\sin \theta = \sin (-\alpha)$
- $\therefore \theta = n\pi + (-1)^n \alpha$ or $\theta = n\pi + (-1)^n (-\alpha)$, where $n \in \mathbb{Z}$.
- $\therefore \theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.
- \therefore The general solution of $\sin^2 \theta = \sin^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Alternative Proof : $\sin^2 \theta = \sin^2 \alpha$

$$\therefore \frac{1 - \cos 2\theta}{2} = \frac{1 - \cos 2\alpha}{2}$$

- $\therefore \cos 2\theta = \cos 2\alpha$
- $\therefore 2\theta = 2n\pi \pm 2\alpha$, where $n \in \mathbb{Z}$.
- $\therefore \theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Theorem 3.5 : The general solution of $\cos^2 \theta = \cos^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Proof : $\cos^2 \theta = \cos^2 \alpha$

- $\therefore \frac{1 + \cos 2\theta}{2} = \frac{1 + \cos 2\alpha}{2}$
- $\therefore \cos 2\theta = \cos 2\alpha$
- $\therefore 2\theta = 2n\pi \pm 2\alpha$, where $n \in \mathbb{Z}$.
- $\therefore \theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Theorem 3.6 : The general solution of $\tan^2 \theta = \tan^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Proof : $\tan^2 \theta = \tan^2 \alpha$

- $\therefore \frac{1 + \tan^2 \theta}{1 - \tan^2 \theta} = \frac{1 + \tan^2 \alpha}{1 - \tan^2 \alpha}$ by componendo and dividendo
- $\therefore \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} = \frac{1 - \tan^2 \alpha}{1 + \tan^2 \alpha}$ by interchanging
- $\therefore \cos 2\theta = \cos 2\alpha$
- $\therefore 2\theta = 2n\pi \pm 2\alpha$, where $n \in \mathbb{Z}$.
- $\therefore \theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

Solved Examples

Ex.(1) Find the general solution of



(i) $\sin \theta = \frac{\sqrt{3}}{2}$ (ii) $\cos \theta = \frac{1}{\sqrt{2}}$ (iii) $\tan \theta = \sqrt{3}$

Solution : (i) We have $\sin \theta = \frac{\sqrt{3}}{2}$

$$\therefore \sin \theta = \sin \frac{\pi}{3}$$

The general solution of $\sin \theta = \sin \alpha$ is $\theta = n\pi + (-1)^n \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \sin \theta = \sin \frac{\pi}{3} \text{ is } \theta = n\pi + (-1)^n \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \text{The general solution of } \sin \theta = \frac{\sqrt{3}}{2} \text{ is } \theta = n\pi + (-1)^n \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}.$$

(ii) We have $\cos \theta = \frac{1}{\sqrt{2}}$

$$\therefore \cos \theta = \cos \frac{\pi}{4}$$

The general solution of $\cos \theta = \cos \alpha$ is $\theta = 2n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \cos \theta = \cos \frac{\pi}{4} \text{ is } \theta = 2n\pi \pm \frac{\pi}{4}, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \text{The general solution of } \cos \theta = \frac{1}{\sqrt{2}} \text{ is } \theta = 2n\pi \pm \frac{\pi}{4}, \text{ where } n \in \mathbb{Z}.$$

(iii) $\tan \theta = \sqrt{3}$

$$\therefore \tan \theta = \tan \frac{\pi}{3}$$

The general solution of $\tan \theta = \tan \alpha$ is $\theta = n\pi + \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \tan \theta = \tan \frac{\pi}{3} \text{ is } \theta = n\pi + \frac{\pi}{3} \text{ where } n \in \mathbb{Z}.$$

$$\therefore \text{The general solution of } \tan \theta = \sqrt{3} \text{ is } \theta = n\pi + \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}.$$

Ex. (2) Find the general solution of

$$(i) \sin \theta = -\frac{\sqrt{3}}{2} \quad (ii) \cos \theta = -\frac{1}{2} \quad (iii) \cot \theta = -\sqrt{3}$$

Solution : (i) $\sin \theta = -\frac{\sqrt{3}}{2}$

$$\therefore \sin \theta = \sin \frac{4\pi}{3} \text{ (As } \sin \frac{4\pi}{3} = \frac{\sqrt{3}}{2} \text{ and } \sin(\pi+A) = -\sin A)$$

The general solution of $\sin \theta = \sin \alpha$ is $\theta = n\pi + (-1)^n \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \sin \theta = \sin \frac{4\pi}{3} \text{ is } \theta = n\pi + (-1)^n \frac{4\pi}{3}, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \text{The general solution of } \sin \theta = -\frac{\sqrt{3}}{2} \text{ is } \theta = n\pi + (-1)^n \frac{4\pi}{3}, \text{ where } n \in \mathbb{Z}.$$

(ii) $\cos \theta = -\frac{1}{2}$

$\therefore \cos \theta = \cos \frac{2\pi}{3}$ (As $\cos \frac{\pi}{3} = \frac{1}{2}$ and $\cos(\pi - A) = -\cos A$)

The general solution of $\cos \theta = \cos \alpha$ is $\theta = 2n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\cos \theta = \cos \frac{2\pi}{3}$ is $\theta = 2n\pi \pm \frac{2\pi}{3}$ where $n \in \mathbb{Z}$.

\therefore The general solution of $\cos \theta = -\frac{1}{2}$ is $\theta = 2n\pi \pm \frac{2\pi}{3}$, where $n \in \mathbb{Z}$.

(iii) $\cot \theta = -\sqrt{3} \quad \therefore \tan \theta = -\frac{1}{\sqrt{3}}$

$\therefore \tan \theta = \tan \frac{5\pi}{6}$ (As $\tan \frac{\pi}{6} = \frac{1}{\sqrt{3}}$ and $\tan(\pi - A) = -\tan A$)

The general solution of $\tan \theta = \tan \alpha$ is $\theta = n\pi + \alpha$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\tan \theta = \tan \frac{5\pi}{6}$ is $\theta = n\pi + \frac{5\pi}{6}$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\cot \theta = -\sqrt{3}$ is $\theta = n\pi + \frac{5\pi}{6}$, where $n \in \mathbb{Z}$.

Ex. (3) Find the general solution of

(i) $\operatorname{cosec} \theta = 2$ (ii) $\sec \theta + \sqrt{2} = 0$

Solution : (i) We have $\operatorname{cosec} \theta = 2 \therefore \sin \theta = \frac{1}{2}$

$\therefore \sin \theta = \sin \frac{\pi}{6}$

The general solution of $\sin \theta = \sin \alpha$ is $\theta = n\pi + (-1)^n \alpha$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\sin \theta = \sin \frac{\pi}{6}$ is $\theta = n\pi + (-1)^n \frac{\pi}{6}$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\operatorname{cosec} \theta = 2$ is $\theta = n\pi + (-1)^n \frac{\pi}{6}$, where $n \in \mathbb{Z}$.

(ii) We have $\sec \theta + \sqrt{2} = 0 \therefore \cos \theta = -\frac{1}{\sqrt{2}}$

$\therefore \cos \theta = \cos \frac{3\pi}{4}$ (As $\cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$ and $\cos(\pi - A) = -\cos A$)

The general solution of $\cos \theta = \cos \alpha$ is $\theta = 2n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\cos \theta = \cos \frac{3\pi}{4}$ is $\theta = 2n\pi \pm \frac{3\pi}{4}$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\sec \theta = -\sqrt{2}$ is $\theta = 2n\pi \pm \frac{3\pi}{4}$, where $n \in \mathbb{Z}$.

Ex. (4) Find the general solution of

$$(i) \cos 2\theta = -\frac{1}{\sqrt{2}} \quad (ii) \tan 3\theta = -1 \quad (iii) \sin 4\theta = \frac{\sqrt{3}}{2}$$

Solution : (i) We have $\cos 2\theta = -\frac{1}{\sqrt{2}}$

$$\therefore \cos 2\theta = \cos \frac{3\pi}{4} \text{ (As } \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}} \text{ and } \cos(\pi - A) = \cos A)$$

The general solution of $\cos \theta = \cos \alpha$ is $\theta = 2n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \cos 2\theta = \cos \frac{3\pi}{4} \text{ is } 2\theta = 2n\pi \pm \frac{3\pi}{4}, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \text{The general solution of } \cos 2\theta = -\frac{1}{\sqrt{2}} \text{ is } \theta = n\pi \pm \frac{3\pi}{8}, \text{ where } n \in \mathbb{Z}.$$

(ii) We have $\tan 3\theta = -1$

$$\therefore \tan 3\theta = \tan \frac{3\pi}{4} \text{ (As } \tan \frac{\pi}{4} = 1 \text{ and } \tan(\pi - A) = -\tan A)$$

The general solution of $\tan \theta = \tan \alpha$ is $\theta = n\pi + \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \tan 3\theta = \tan \frac{3\pi}{4} \text{ is } 3\theta = n\pi + \frac{3\pi}{4} \text{ where } n \in \mathbb{Z}.$$

$$\therefore \text{The general solution of } \tan 3\theta = -1 \text{ is } \theta = \frac{n\pi}{3} + \frac{\pi}{4}, \text{ where } n \in \mathbb{Z}.$$

(iii) $\sin 4\theta = \frac{\sqrt{3}}{2}$

$$\therefore \sin 4\theta = \sin \frac{\pi}{3}$$

The general solution of $\sin \theta = \sin \alpha$ is $\theta = n\pi + (-1)^n \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \sin 4\theta = \sin \frac{\pi}{3} \text{ is } 4\theta = n\pi + (-1)^n \frac{\pi}{3}, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \text{The general solution of } \sin 4\theta = \frac{\sqrt{3}}{2} \text{ is } \theta = \frac{n\pi}{4} + (-1)^n \frac{\pi}{12} \text{ where } n \in \mathbb{Z}.$$

Ex. (5) Find the general solution of

$$(i) 4 \cos^2 \theta = 1 \quad (ii) 4 \sin^2 \theta = 3 \quad (iii) \tan^2 \theta = 1$$

Solution : (i) We have $4 \cos^2 \theta = 1$

$$\therefore \cos^2 \theta = \frac{1}{4} = \left(\frac{1}{2}\right)^2$$

$$\therefore \cos^2 \theta = \cos^2 \frac{\pi}{3}$$

The general solution of $\cos^2 \theta = \cos^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\cos^2 \theta = \cos^2 \frac{\pi}{3}$ is $\theta = n\pi \pm \frac{\pi}{3}$, where $n \in \mathbb{Z}$.

\therefore The general solution of $4 \cos^2 \theta = 1$ is $\theta = n\pi \pm \frac{\pi}{3}$, where $n \in \mathbb{Z}$.

(ii) We have $4\sin^2 \theta = 3$

$$\therefore \sin^2 \theta = \frac{3}{4} = \left(\frac{\sqrt{3}}{2}\right)^2$$

$$\therefore \sin^2 \theta = \sin^2 \frac{\pi}{3}$$

The general solution of $\sin^2 \theta = \sin^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\sin^2 \theta = \sin^2 \frac{\pi}{3}$ is $\theta = n\pi \pm \frac{\pi}{3}$, where $n \in \mathbb{Z}$.

\therefore The general solution of $4\sin^2 \theta = 3$ is $\theta = n\pi \pm \frac{\pi}{3}$, where $n \in \mathbb{Z}$.

(iii) We have $\tan^2 \theta = 1$

$$\therefore \tan^2 \theta = \tan^2 \frac{\pi}{4}$$

The general solution of $\tan^2 \theta = \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\tan^2 \theta = \tan^2 \frac{\pi}{4}$ is $\theta = n\pi \pm \frac{\pi}{4}$, where $n \in \mathbb{Z}$.

\therefore The general solution of $\tan^2 \theta = 1$ is $\theta = n\pi \pm \frac{\pi}{4}$, where $n \in \mathbb{Z}$.

Ex. (6) Find the general solution of $\cos 3\theta = \cos 2\theta$

Solution : We have $\cos 3\theta = \cos 2\theta$

$$\therefore \cos 3\theta - \cos 2\theta = 0$$

$$\therefore -2 \sin \frac{5\theta}{2} \sin \frac{\theta}{2} = 0$$

$$\therefore \sin \frac{5\theta}{2} = 0 \text{ or } \sin \frac{\theta}{2} = 0$$

$$\therefore \frac{5\theta}{2} = n\pi \text{ or } \frac{\theta}{2} = n\pi \text{ where } n \in \mathbb{Z}.$$

$$\therefore \theta = \frac{2n\pi}{5}, n \in \mathbb{Z}.$$

$$\therefore \theta = \frac{2n\pi}{5} \text{ where } n \in \mathbb{Z} \text{ is the required general solution.}$$

Alternative Method : We know that the general solution of $\cos \theta = \cos \alpha$ is $\theta = 2n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

$$\therefore \text{The general solution of } \cos 3\theta = \cos 2\theta \text{ is } 3\theta = 2n\pi \pm 2\theta, \text{ where } n \in \mathbb{Z}.$$

$$\therefore 3\theta = 2n\pi - 2\theta \text{ or } 3\theta = 2n\pi + 2\theta, \text{ where } n \in \mathbb{Z}.$$

$$\therefore 5\theta = 2n\pi \text{ or } \theta = 2n\pi, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \theta = \frac{2n\pi}{5}, n \in \mathbb{Z} \text{ where } n \in \mathbb{Z} \text{ is the required general solution.}$$

Ex. (7) Find the general solution of $\cos 5\theta = \sin 3\theta$

Solution : We have $\cos 5\theta = \sin 3\theta$

$$\therefore \cos 5\theta = \cos \left(\frac{\pi}{2} - 3\theta \right)$$

$$\therefore 5\theta = 2n\pi \pm \left(\frac{\pi}{2} - 3\theta \right)$$

$$\therefore 5\theta = 2n\pi - \left(\frac{\pi}{2} - 3\theta \right) \text{ or } 5\theta = 2n\pi + \left(\frac{\pi}{2} - 3\theta \right)$$

$$\therefore \theta = n\pi - \frac{\pi}{4} \text{ or } \theta = \frac{n\pi}{4} + \frac{\pi}{16}, \text{ where } n \in \mathbb{Z} \text{ are the required general solutions.}$$

Ex. (8) Find the general solution of $\sec^2 2\theta = 1 - \tan 2\theta$

Solution : Given equation is $\sec^2 2\theta = 1 - \tan 2\theta$

$$\therefore 1 + \tan^2 2\theta = 1 - \tan 2\theta$$

$$\therefore \tan^2 2\theta + \tan 2\theta = 0$$

$$\therefore \tan 2\theta (\tan 2\theta + 1) = 0$$

$$\therefore \tan 2\theta = 0 \text{ or } \tan 2\theta + 1 = 0$$

$$\therefore \tan 2\theta = \tan 0 \text{ or } \tan 2\theta = \tan \frac{3\pi}{4}$$

$$\therefore 2\theta = n\pi \text{ or } 2\theta = n\pi + \frac{3\pi}{4}, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \theta = \frac{n\pi}{2} \text{ or } \theta = \frac{n\pi}{2} + \frac{3\pi}{8}, \text{ where } n \in \mathbb{Z} \text{ is the required general solution.}$$

Ex. (9) Find the general solution of $\sin \theta + \sin 3\theta + \sin 5\theta = 0$

Solution : We have $\sin \theta + \sin 3\theta + \sin 5\theta = 0$

$$\therefore (\sin \theta + \sin 5\theta) + \sin 3\theta = 0$$

$$\therefore 2 \sin 3\theta \cos 2\theta + \sin 3\theta = 0$$

$$\therefore (2 \cos 2\theta + 1) \sin 3\theta = 0$$

$$\therefore \sin 3\theta = 0 \text{ or } \cos 2\theta = -\frac{1}{2}$$

$$\therefore \sin 3\theta = 0 \text{ or } \cos 2\theta = \cos \frac{2\pi}{3}$$

$$\therefore 3\theta = n\pi \text{ or } 2\theta = 2n\pi \pm \frac{2\pi}{3}, \text{ where } n \in \mathbb{Z}.$$

$$\therefore \theta = \frac{n\pi}{3} \text{ or } \theta = n\pi \pm \frac{\pi}{3}, \text{ where } n \in \mathbb{Z} \text{ is the required general solution.}$$

Ex. (10) Find the general solution of $\cos\theta - \sin\theta = 1$

Solution : We have $\cos\theta - \sin\theta = 1$

$$\therefore \frac{1}{\sqrt{2}} \cos\theta - \frac{1}{\sqrt{2}} \sin\theta = \frac{1}{\sqrt{2}}$$

$$\therefore \cos\theta \cos \frac{\pi}{4} - \sin\theta \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}$$

$$\therefore \cos\left(\theta + \frac{\pi}{4}\right) = \cos \frac{\pi}{4}$$

$$\therefore \theta + \frac{\pi}{4} = 2n\pi \pm \frac{\pi}{4}$$

$$\therefore \theta + \frac{\pi}{4} = 2n\pi - \frac{\pi}{4} \text{ or } \theta + \frac{\pi}{4} = 2n\pi + \frac{\pi}{4}$$

$$\therefore \theta = 2n\pi - \frac{\pi}{2} \text{ or } \theta = 2n\pi, \text{ where } n \in \mathbb{Z} \text{ is the required general solution.}$$



Exercise 3.1

1) Find the principal solutions of the following equations :

$$(i) \cos\theta = \frac{1}{2} \quad (ii) \sec\theta = \frac{2}{\sqrt{3}} \quad (iii) \cot\theta = \sqrt{3} \quad (iv) \cot\theta = 0$$

2) Find the principal solutions of the following equations:

$$(i) \sin\theta = -\frac{1}{2} \quad (ii) \tan\theta = -1 \quad (iii) \sqrt{3} \operatorname{cosec}\theta + 2 = 0$$

Find the general solutions of the following equations :

$$3) (i) \sin\theta = \frac{1}{2} \quad (ii) \cos\theta = \frac{\sqrt{3}}{2} \quad (iii) \tan\theta = \frac{1}{\sqrt{3}} \quad (iv) \cot\theta = 0$$

$$4) (i) \sec\theta = \sqrt{2} \quad (ii) \operatorname{cosec}\theta = -\sqrt{2} \quad (iii) \tan\theta = -1$$

$$5) (i) \sin 2\theta = \frac{1}{2} \quad (ii) \tan \frac{2\theta}{3} = \sqrt{3} \quad (iii) \cot 4\theta = -1$$

$$6) (i) 4 \cos^2\theta = 3 \quad (ii) 4 \sin^2\theta = 1 \quad (iii) \cos 4\theta = \cos 2\theta$$

$$7) (i) \sin\theta = \tan\theta \quad (ii) \tan^3\theta = 3\tan\theta \quad (iii) \cos\theta + \sin\theta = 1$$

8) Which of the following equations have solutions ?

$$(i) \cos 2\theta = -1 \quad (ii) \cos^2\theta = -1 \quad (iii) 2 \sin\theta = 3 \quad (iv) 3 \tan\theta = 5$$

3.2 Solution of triangle

3.2.1 Polar co-ordinates : Let O be a fixed point in a plane. Let OX be a fixed ray in the plane. O is called the pole and ray OX is called the polar axis. Let P be a point in the plane other than pole O.

Let $OP = r$ and $\angle XOP = \theta$. The ordered pair (r, θ) determines the position of P in the plane. They are called the polar co-ordinates of P. 'r' is called the radius vector and θ is called the vectorial angle of point P.




Fig 3.1

Remarks :

- i) Vectorial angle θ is the smallest non-negative angle made by OP with the ray OX.
- ii) $0 \leq \theta < 2\pi$
- iii) Pole has no polar co-ordinates.

3.2.2 Relation between the Cartesian and the Polar co-ordinates: Let O be the pole and OX be the polar axis of polar co-ordinates system. We take line along OX as the X - axis and line perpendicular to OX through O as the Y - axis.

Let P be any point in the plane other than origin. Let (x, y) and (r, θ) be Cartesian and polar co-ordinates of P. To find the relation between them.

By definition of trigonometric functions, we have $\sin \theta = \frac{y}{r}$ and

$$\cos \theta = \frac{x}{r}$$

$$\therefore x = r \cos \theta \text{ and } y = r \sin \theta$$

This is the relation between Cartesian and polar co-ordinates.




Fig 3.2

Ex. (1) Find the Cartesian co-ordinates of the point whose polar co-ordinates are $\left(2, \frac{\pi}{4}\right)$

Solution : Given $r = 2$ and $\theta = \frac{\pi}{4}$

Using $x = r \cos \theta$ and $y = r \sin \theta$, we get

$$x = 2 \cos \frac{\pi}{4} = 2 \times \frac{1}{\sqrt{2}} = \sqrt{2}$$

$$y = 2 \sin \frac{\pi}{4} = 2 \times \frac{1}{\sqrt{2}} = \sqrt{2}$$

The required Cartesian co-ordinates are $(\sqrt{2}, \sqrt{2})$.

Ex. (2) Find the polar co-ordinates of point whose Cartesian co-ordinates are $\left(\frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}}\right)$

Solution : From the co-ordinates of the given point we observe that point lies in the fourth quadrant.

$$r^2 = x^2 + y^2$$

$$\therefore r^2 = \left(\frac{1}{\sqrt{2}}\right)^2 + \left(\frac{-1}{\sqrt{2}}\right)^2 = \frac{1}{2} + \frac{1}{2} = 1$$

$$\therefore r = 1$$

$$x = r \cos \theta, y = r \sin \theta$$

$$\therefore \frac{1}{\sqrt{2}} = 1 \times \cos \theta \text{ and } -\frac{1}{\sqrt{2}} = 1 \times \sin \theta$$

$$\therefore \cos \theta = \frac{1}{\sqrt{2}} \text{ and } \sin \theta = -\frac{1}{\sqrt{2}}$$

$$\therefore \theta = \frac{7\pi}{4}$$

\therefore The required polar co-ordinates are $\left(1, \frac{7\pi}{4}\right)$.

3.2.3 Solving a Triangle :

Three sides and three angles of a triangle are called the elements of the triangle. If we have a certain set of three elements of a triangle, in which at least one element is a side, then we can determine other three elements of the triangle. To solve a triangle means to find unknown elements of the triangle. Using three angles of a triangle we can't solve it. At least one side should be known. In ΔABC , we use the following notations : $l(BC) = BC = a$, $l(CA) = AC = b$, $l(AB) = AB = c$. This notation is called as the usual notation. Following are some standard relations between elements of triangle.

3.2.4 The Sine Rule : In ΔABC , $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$, where R is the circumradius of ΔABC .

Proof : Let AD be perpendicular to BC .

$$AD = b \sin C$$

$$\therefore A(\Delta ABC) = \frac{1}{2} BC \times AD$$

$$= \frac{1}{2} a \times b \sin C$$

$$\therefore A(\Delta ABC) = \frac{1}{2} ab \sin C$$

$$\therefore 2A(\Delta ABC) = ab \sin C$$

Similarly $2A(\Delta ABC) = ac \sin B$ and $2A(\Delta ABC) = bc \sin A$

$$\therefore bc \sin A = ac \sin B = ab \sin C$$

Divide by abc ,

$$\therefore \frac{bc \sin A}{abc} = \frac{ac \sin B}{abc} = \frac{ab \sin C}{abc}$$

$$\therefore \frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

$$\therefore \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \dots (1)$$





Fig 3.3



To prove that each ratio is equal to $2R$.

As the sum of three angles is 180° , at least one of the angle of the triangle is not right angle.

Suppose A is not right angle.

Draw diameter through A . Let it meet circle in P .

$\therefore AP = 2R$ and $\triangle ACP$ is a right angled triangle. $\angle ABC$ and $\angle APC$ are inscribed in the same arc.

$\therefore m \angle ABC = m \angle APC$

$$\therefore \sin B = \sin P = \frac{b}{AP} = \frac{b}{2R}$$

$$\therefore \sin B = \frac{b}{2R}$$

$$\therefore \frac{b}{\sin B} = 2R \quad \dots (2)$$

From (1) and (2), we get

$$\therefore \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$$

Different forms of Sine rule : Following are the different forms of the Sine rule.

In $\triangle ABC$.

$$(i) \quad \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$$

$$(ii) \quad a = 2R \sin A, b = 2R \sin B, c = 2R \sin C$$

$$(iii) \quad \frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} = k$$

$$(iv) \quad b \sin A = a \sin B, c \sin B = b \sin C, c \sin A = a \sin C$$

$$(v) \quad \frac{a}{b} = \frac{\sin A}{\sin B}, \frac{b}{c} = \frac{\sin B}{\sin C}$$

Ex.(1) In $\triangle ABC$ if $A = 30^\circ$, $B = 60^\circ$ then find the ratio of sides.

Solution : To find $a : b : c$

Given $A = 30^\circ$, $B = 60^\circ$.

As A, B, C are angles of the triangle, $A + B + C = 180^\circ$

$$\therefore C = 90^\circ$$

By Sine rule,

$$\therefore \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$\therefore \frac{a}{\sin 30^\circ} = \frac{b}{\sin 60^\circ} = \frac{c}{\sin 90^\circ}$$

$$\therefore \frac{a}{\frac{1}{2}} = \frac{b}{\frac{\sqrt{3}}{2}} = \frac{c}{1}$$

$$\therefore a : b : c = \frac{1}{2} : \frac{\sqrt{3}}{2} : 1$$

$$\therefore a : b : c = 1 : \sqrt{3} : 2$$

Ex.(2) In $\triangle ABC$ if $a = 2$, $b = 3$ and $\sin A = \frac{2}{3}$ then find B .

Solution : By sine rule, $\frac{a}{\sin A} = \frac{b}{\sin B}$

$$\therefore \frac{2}{\frac{2}{3}} = \frac{3}{\sin B}$$

$$\therefore \sin B = 1$$

$$\therefore B = 90^\circ = \frac{\pi}{2}$$

Ex. (3) In $\triangle ABC$, prove that $a(\sin B - \sin C) + b(\sin C - \sin A) + c(\sin A - \sin B) = 0$

Solution : L.H.S. = $a(\sin B - \sin C) + b(\sin C - \sin A) + c(\sin A - \sin B)$

$$= a \sin B - a \sin C + b \sin C - b \sin A + c \sin A - c \sin B$$

$$= (a \sin B - b \sin A) + (b \sin C - c \sin B) + (c \sin A - a \sin C)$$

$$= 0 + 0 + 0$$

$$= 0 = \text{R.H.S.}$$

Ex.(4) In $\triangle ABC$, prove that $(a - b) \sin C + (b - c) \sin A + (c - a) \sin B = 0$

Solution : L.H.S. = $(a - b) \sin C + (b - c) \sin A + (c - a) \sin B$

$$= (a \sin C - b \sin C) + (b \sin A - c \sin A) + (c \sin B - a \sin B)$$

$$= (a \sin C - c \sin A) + (b \sin A - a \sin B) + (c \sin B - b \sin C)$$

$$= 0 + 0 + 0 = 0 = \text{R.H.S.}$$

3.3.5 The Cosine Rule : In $\triangle ABC$,

$$(i) \quad a^2 = b^2 + c^2 - 2bc \cos A \quad (ii) \quad b^2 = c^2 + a^2 - 2ca \cos B \quad (iii) \quad c^2 = a^2 + b^2 - 2ab \cos C$$

Proof : Take A as the origin, X - axis along AB and the line perpendicular to AB through A as the Y - axis. The co-ordinates of A , B and C are $(0,0)$, $(c, 0)$ and $(b \cos A, b \sin A)$ respectively.

To prove that $a^2 = b^2 + c^2 - 2bc \cos A$

$$\text{L.H.S.} = a^2 = BC^2$$

$$= (c - b \cos A)^2 + (0 - b \sin A)^2 \text{ (by distance formula)}$$

$$= c^2 + b^2 \cos^2 A - 2bc \cos A + b^2 \sin^2 A$$

$$= c^2 + b^2 \cos^2 A + b^2 \sin^2 A - 2bc \cos A$$

$$= c^2 + b^2 - 2bc \cos A$$

$$= \text{R.H.S.}$$

$$\therefore a^2 = b^2 + c^2 - 2bc \cos A$$

Similarly, we can prove that

$$b^2 = c^2 + a^2 - 2ca \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$




Fig 3.5

Remark : The cosine rule can be stated as : In $\triangle ABC$,

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}, \quad \cos B = \frac{c^2 + a^2 - b^2}{2ca}, \quad \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

Ex.(5) In $\triangle ABC$, if $a = 2$, $b = 3$, $c = 4$ then prove that the triangle is obtuse angled.

Solution : We know that the angle opposite to largest side of a triangle is the largest angle of the triangle.

Here side AB is the largest side. C is the largest angle of $\triangle ABC$. To show that C is obtuse angle.

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab} = \frac{2^2 + 3^2 - 4^2}{2(3)(4)} = -\frac{3}{24} = -\frac{1}{8}$$

As $\cos C$ is negative, C is obtuse angle.

$\therefore \triangle ABC$ is obtuse angled triangle.

Ex.(6) In $\triangle ABC$, if $A = 60^\circ$, $b = 3$ and $c = 8$ then find a . Also find the circumradius of the triangle.

Solution : By Cosine rule, $a^2 = b^2 + c^2 - 2bc \cos A$

$$\therefore a^2 = 3^2 + 8^2 - 2(3)(8) \cos(60^\circ)$$

$$= 9 + 64 - 48 \times \frac{1}{2}$$

$$= 73 - 24 = 49$$

$$\therefore a^2 = 49$$

$$\therefore a = 7$$

Now by sine rule $\frac{a}{\sin A} = 2R$

$$\therefore \frac{7}{\sin 60^\circ} = 2R$$

$$\therefore \frac{7}{\frac{\sqrt{3}}{2}} = 2R$$

$$\therefore R = \frac{7}{\sqrt{3}} = \frac{7\sqrt{3}}{3}$$

The circumradius of the $\triangle ABC$ is $\frac{7\sqrt{3}}{3}$

Ex. (7) In $\triangle ABC$ prove that $a(b \cos C - c \cos B) = b^2 - c^2$

Solution :

$$\text{L.H.S.} = a(b \cos C - c \cos B)$$

$$= ab \cos C - ac \cos B$$

$$= \frac{1}{2} (2ab \cos C - 2ac \cos B)$$

$$= \frac{1}{2} \{ (a^2 + b^2 - c^2) - (c^2 + a^2 - b^2) \}$$

$$\begin{aligned}
 &= \frac{1}{2} \{ a^2 + b^2 - c^2 - c^2 - a^2 + b^2 \} \\
 &= \frac{1}{2} \{ 2b^2 - 2c^2 \} \\
 &= b^2 - c^2 = \text{R.H.S.}
 \end{aligned}$$

3.3.6 The projection Rule : In $\triangle ABC$,

- (i) $a = b \cos C + c \cos B$
- (ii) $b = c \cos A + a \cos C$
- (iii) $c = a \cos B + b \cos A$

Proof : Here we give proof of one of these three statements, by considering all possible cases.

To prove that $a = b \cos C + c \cos B$

Let altitude drawn from A meets BC in D.

BD is called the projection of AB on BC.

DC is called the projection of AC on BC.

\therefore Projection of AB on BC = $c \cos B$

And projection of AC on BC = DC = $b \cos C$

Case (i) B and C are acute angles.

\therefore Projection of AB on BC = BD = $c \cos B$

And projection of AC on BC = DC = $b \cos C$

From figure we have,

$$\begin{aligned}
 a &= BC = BD + DC \\
 &= c \cos B + b \cos C \\
 &= b \cos C + c \cos B
 \end{aligned}$$

$\therefore a = b \cos C + c \cos B$

Case (ii) B is obtuse angle.

\therefore Projection AB on BC = BD = $c \cos (\pi - B) = -c \cos B$

And projection of AC on BC = DC = $b \cos C$

From figure we have,

$$\begin{aligned}
 a &= BC = DC - BD \\
 &= b \cos C - (-c \cos B) \\
 &= b \cos C + c \cos B
 \end{aligned}$$

$\therefore a = b \cos C + c \cos B$

Case (iii) B is right angle. In this case D coincides with B.

$$\begin{aligned}
 \text{R.H.S.} &= b \cos C + c \cos B \\
 &= BC + 0 \\
 &= a = \text{L.H.S.}
 \end{aligned}$$

$\therefore a = b \cos C + c \cos B$

Similarly we can prove the cases where C is obtuse angle and C right angle.

Therefore in all possible cases, $a = b \cos C + c \cos B$

Similarly we can prove other statements.




Fig 3.6




Fig 3.7

Ex.(8) In $\triangle ABC$, prove that $(a + b) \cos C + (b + c) \cos A + (c + a) \cos B = a + b + c$

Solution :

$$\begin{aligned} \text{L.H.S.} &= (a + b) \cos C + (b + c) \cos A + (c + a) \cos B \\ &= (a \cos C + b \cos C) + (b \cos A + c \cos A) + (c \cos B + a \cos B) \\ &= (a \cos C + c \cos A) + (b \cos A + a \cos B) + (c \cos B + b \cos C) \\ &= a + b + c = \text{R.H.S.} \end{aligned}$$

Ex.(9) In $\triangle ABC$, prove that $a(\cos C - \cos B) = 2(b - c) \cos^2 \left(\frac{A}{2} \right)$

Solution : By Projection rule, we have $a \cos C + c \cos A = b$ and $a \cos B + b \cos A = c$

$$\therefore a \cos C = b - c \cos A \text{ and } a \cos B = c - b \cos A$$

$$\begin{aligned} \text{L.H.S.} &= a(\cos C - \cos B) \\ &= a \cos C - a \cos B \\ &= (b - c \cos A) - (c - b \cos A) \\ &= b - c \cos A - c + b \cos A \\ &= (b - c) + (b - c) \cos A \\ &= (b - c)(1 + \cos A) \\ &= (b - c) \times 2 \cos^2 \frac{A}{2} \\ &= 2(b - c) \cos^2 \frac{A}{2} \\ &= \text{R.H.S.} \end{aligned}$$

Ex.(10) Prove the Cosine rule using the Projection rule.

Solution : Given: In $\triangle ABC$, $a = b \cos C + c \cos B$

$$b = c \cos A + a \cos C$$

$$c = a \cos B + b \cos A$$

Multiply these equations by a, b, c respectively.

$$a^2 = ab \cos C + ac \cos B$$

$$b^2 = bc \cos A + ab \cos C$$

$$c^2 = ac \cos B + bc \cos A$$

$$a^2 + b^2 - c^2 = (ab \cos C + ac \cos B) + (bc \cos A + ab \cos C) - (ac \cos B + bc \cos A)$$

$$= ab \cos C + ac \cos B + bc \cos A + ab \cos C - ac \cos B - bc \cos A$$

$$= 2ab \cos C$$

$$\therefore a^2 + b^2 - c^2 = 2ab \cos C \quad \therefore c^2 = a^2 + b^2 - 2ab \cos C.$$

Similarly we can prove that

$$a^2 = b^2 + c^2 - 2bc \cos A \text{ and } b^2 = c^2 + a^2 - 2ca \cos B.$$

3.3.7 Applications of Sine rule, Cosine rule and Projection rule:

(1) Half angle formulae : In $\triangle ABC$, if $a + b + c = 2s$ then

$$(i) \quad \sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}} \quad (ii) \quad \cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}} \quad (iii) \quad \tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$$

Proof : (i) We have, $1 \therefore \cos A = 2 \sin^2 \frac{A}{2}$

$$\therefore 1 \therefore \left(\frac{b^2 + c^2 - a^2}{2bc} \right) = 2 \sin^2 \frac{A}{2} \text{ by cosine rule}$$

$$\therefore \frac{2bc - b^2 - c^2 + a^2}{2bc} = 2 \sin^2 \frac{A}{2}$$

$$\therefore \frac{a^2 - (b^2 + c^2 - 2bc)}{2bc} = 2 \sin^2 \frac{A}{2}$$

$$\therefore \frac{a^2 - (b-c)^2}{2bc} = 2 \sin^2 \frac{A}{2}$$

$$\therefore \frac{\{a - (b-c)\} \{a + (b-c)\}}{2bc} = 2 \sin^2 \frac{A}{2}$$

$$\therefore \frac{\{a+b+c-2b\} \{a+b+c-2c\}}{2bc} = 2 \sin^2 \frac{A}{2}$$

$$\therefore \frac{\{2s-2b\} \{2s-2c\}}{2bc} = 2 \sin^2 \frac{A}{2}$$

$$\therefore \frac{(s-b)(s-c)}{bc} = \sin^2 \frac{A}{2}$$

$$\therefore \sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}}$$

(ii) We have, $1 + \cos A = 2 \cos^2 \frac{A}{2}$

$$\therefore 1 + \left(\frac{b^2 + c^2 - a^2}{2bc} \right) = 2 \cos^2 \frac{A}{2} \text{ by cosine rule}$$

$$\therefore \frac{2bc + b^2 + c^2 - a^2}{2bc} = 2 \cos^2 \frac{A}{2}$$

$$\therefore \frac{(b^2 + c^2 + 2bc) - a^2}{2bc} = 2 \cos^2 \frac{A}{2}$$

$$\therefore \frac{(b+c)^2 - a^2}{2bc} = 2 \cos^2 \frac{A}{2}$$

$$\therefore \frac{(b+c+a)(b+c-a)}{2bc} = 2 \cos^2 \frac{A}{2}$$

$$\therefore \frac{(b+c+a)(b+c+a-2a)}{2bc} = 2\cos^2 \frac{A}{2}$$

$$\therefore \frac{(2s)(2s-2a)}{2bc} = 2\cos^2 \frac{A}{2}$$

$$\therefore \cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}$$

$$\begin{aligned} \text{(iii) } \tan \frac{A}{2} &= \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}} = \frac{\sqrt{\frac{(s-b)(s-c)}{bc}}}{\sqrt{\frac{s(s-a)}{bc}}} \\ &= \sqrt{\frac{(s-b)(s-c)}{s(s-a)}} \end{aligned}$$

$$\therefore \tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$$

Similarly we can prove that

$$\sin \frac{B}{2} = \sqrt{\frac{(s-a)(s-c)}{ac}}, \quad \sin \frac{C}{2} = \sqrt{\frac{(s-b)(s-a)}{ab}}$$

$$\cos \frac{B}{2} = \sqrt{\frac{s(s-b)}{ac}}, \quad \cos \frac{C}{2} = \sqrt{\frac{s(s-c)}{ab}}$$

$$\tan \frac{B}{2} = \sqrt{\frac{(s-a)(s-c)}{s(s-b)}}, \quad \tan \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{s(s-c)}}$$

(2) Heron's Formula : If a, b, c are sides of $\triangle ABC$ and $a + b + c = 2s$ then

$$A(\triangle ABC) = \sqrt{s(s-a)(s-b)(s-c)}$$

Proof : We know that $A(\triangle ABC) = \frac{1}{2} ab \sin C$

$$\therefore A(\triangle ABC) = \frac{1}{2} ab 2 \sin \frac{C}{2} \cos \frac{C}{2}$$

$$= ab \sqrt{\frac{(s-b)(s-a)}{ab}} \sqrt{\frac{s(s-c)}{ab}} = \sqrt{s(s-a)(s-b)(s-c)}$$

(3) Napier's Analogy : In $\triangle ABC$, $\tan \left(\frac{B-C}{2} \right) = \frac{(b-c)}{(b+c)} \cot \frac{A}{2}$

Proof : By sine rule $b = 2R \sin B$ and $c = 2R \sin C$

$$\therefore \frac{(b-c)}{(b+c)} = \frac{2R \sin B - 2R \sin C}{2R \sin B + 2R \sin C}$$

$$\therefore \frac{(b-c)}{(b+c)} = \frac{\sin B - \sin C}{\sin B + \sin C}$$

$$\therefore \frac{(b-c)}{(b+c)} = \frac{2 \cos\left(\frac{B+C}{2}\right) \sin\left(\frac{B-C}{2}\right)}{2 \sin\left(\frac{B+C}{2}\right) \cos\left(\frac{B-C}{2}\right)}$$

$$\therefore \frac{(b-c)}{(b+c)} = \cot\left(\frac{B+C}{2}\right) \tan\left(\frac{B-C}{2}\right)$$

$$\therefore \frac{(b-c)}{(b+c)} = \cot\left(\frac{\pi}{2} - \frac{A}{2}\right) \tan\left(\frac{B-C}{2}\right)$$

$$\therefore \frac{(b-c)}{(b+c)} = \tan\left(\frac{A}{2}\right) \tan\left(\frac{B-C}{2}\right)$$

$$\therefore \tan\left(\frac{B-C}{2}\right) = \frac{(b-c)}{(b+c)} \cot \frac{A}{2}$$

Similarly we can prove that

$$\tan\left(\frac{C-A}{2}\right) = \frac{(c-a)}{(c+a)} \cot \frac{B}{2}$$

$$\tan\left(\frac{A-B}{2}\right) = \frac{(a-b)}{(a+b)} \cot \frac{C}{2}$$



Solved Examples

Ex.(1) In ΔABC if $a = 13$, $b = 14$, $c = 15$ then find the values of

- (i) $\cos A$ (ii) $\sin \frac{A}{2}$ (iii) $\cos \frac{A}{2}$ (iv) $\tan \frac{A}{2}$ (v) $A(\Delta ABC)$ (vi) $\sin A$

Solution :

$$s = \frac{a+b+c}{2} = \frac{13+14+15}{2} = 21$$

$$\begin{aligned}(s - a) &= 21 - 13 = 8 \\(s - b) &= 21 - 14 = 7 \\(s - c) &= 21 - 15 = 6\end{aligned}$$

$$\begin{aligned}\text{(i)} \quad \cos A &= \frac{b^2 + c^2 - a^2}{2bc} \\&= \frac{13^2 + 15^2 - 14^2}{2(13)(15)} = \frac{198}{390} = \frac{33}{65}\end{aligned}$$

$$\begin{aligned}\text{(ii)} \quad \sin \frac{A}{2} &= \sqrt{\frac{(s-b)(s-c)}{bc}} \\&= \sqrt{\frac{7 \times 6}{14 \times 15}} = \frac{1}{\sqrt{5}}\end{aligned}$$

$$\begin{aligned}\text{(iii)} \quad \cos \frac{A}{2} &= \sqrt{\frac{s(s-a)}{bc}} \\&= \sqrt{\frac{21 \times 8}{14 \times 15}} = \frac{2}{\sqrt{5}}\end{aligned}$$

$$\text{(iv)} \quad \tan \frac{A}{2} = \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}} = \frac{\frac{1}{\sqrt{5}}}{\frac{2}{\sqrt{5}}} = \frac{1}{2}$$

$$\begin{aligned}\text{(v)} \quad A(\Delta ABC) &= \sqrt{s(s-a)(s-b)(s-c)} \\&= \sqrt{21 \times 8 \times 7 \times 6} = 84 \text{ sq. unit}\end{aligned}$$

$$\begin{aligned}\text{(vi)} \quad \sin A &= 2 \sin \frac{A}{2} \cos \frac{A}{2} \\&= 2 \times \frac{1}{\sqrt{5}} \times \frac{2}{\sqrt{5}} = \frac{4}{5}\end{aligned}$$

Ex.(2) In ΔABC prove that $\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \frac{a+b+c}{b+c-a} \cot \frac{A}{2}$

$$\begin{aligned}\text{Solution : L.H.S.} &= \cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} \\&= \frac{1}{\tan \frac{A}{2}} + \frac{1}{\tan \frac{B}{2}} + \frac{1}{\tan \frac{C}{2}}\end{aligned}$$

$$\begin{aligned}
 &= \sqrt{\frac{s(s-a)}{(s-b)(s-c)}} + \sqrt{\frac{s(s-b)}{(s-a)(s-c)}} + \sqrt{\frac{s(s-c)}{(s-a)(s-b)}} \\
 &= \sqrt{\frac{s(s-a)^2}{(s-b)(s-c)(s-a)}} + \sqrt{\frac{s(s-b)^2}{(s-a)(s-c)(s-b)}} + \sqrt{\frac{s(s-c)^2}{(s-b)(s-a)(s-c)}} \\
 &= \sqrt{\frac{s}{(s-b)(s-a)(s-c)}} \left\{ \sqrt{(s-a)^2} + \sqrt{(s-b)^2} + \sqrt{(s-c)^2} \right\} \\
 &= \sqrt{\frac{s}{(s-b)(s-a)(s-c)}} \{(s-a) + (s-c) + (s-b)\} \\
 &= \sqrt{\frac{s}{(s-b)(s-a)(s-c)}} \{3s - (a+b+c)\} \\
 &= \sqrt{\frac{s}{(s-b)(s-a)(s-c)}} \{3s - 2s\} \\
 &= \sqrt{\frac{s}{(s-b)(s-a)(s-c)}} \times s \\
 &= \sqrt{\frac{s}{(s-b)(s-c)}} \times \frac{s}{\sqrt{(s-a)}} \\
 &= \sqrt{\frac{s(s-a)}{(s-b)(s-c)}} \times \frac{s}{(s-a)} \\
 &= \frac{2s}{(2s-2a)} \times \sqrt{\frac{s(s-a)}{(s-b)(s-c)}} \\
 &= \frac{a+b+c}{(a+b+c-2a)} \times \sqrt{\frac{s(s-a)}{(s-b)(s-c)}} \\
 &= \frac{a+b+c}{b+c-a} \cot \frac{A}{2} = \text{R.H.S.}
 \end{aligned}$$



Exercise 3.2

- Find the Cartesian co-ordinates of the point whose polar co-ordinates are :
 - $(\sqrt{2}, \frac{\pi}{4})$
 - $(4, \frac{\pi}{2})$
 - $(\frac{3}{4}, \frac{3\pi}{4})$
 - $(\frac{1}{2}, \frac{7\pi}{3})$
- Find the of the polar co-ordinates point whose Cartesian co-ordinates are.
 - $(\sqrt{2}, \sqrt{2})$
 - $(0, \frac{1}{2})$
 - $(1, -\sqrt{3})$
 - $(\frac{3}{2}, \frac{3\sqrt{3}}{2})$
- In ΔABC , if $A = 45^\circ$, $B = 60^\circ$ then find the ratio of its sides.
- In ΔABC , prove that $\sin\left(\frac{B-C}{2}\right) = \left(\frac{b-c}{a}\right) \cos \frac{A}{2}$.
- With usual notations prove that $2\left\{a \sin^2 \frac{C}{2} + c \sin^2 \frac{A}{2}\right\} = a - b + c$.
- In ΔABC , prove that $a^3 \sin(B - C) + b^3 \sin(C - A) + c^3 \sin(A - B) = 0$
- In ΔABC , if $\cot A, \cot B, \cot C$ are in A.P. then show that a^2, b^2, c^2 are also in A.P
- In ΔABC , if $a \cos A = b \cos B$ then prove that the triangle is right angled or an isosceles traingle.
- With usual notations prove that $2(bc \cos A + ac \cos B + ab \cos C) = a^2 + b^2 + c^2$
- In ΔABC , if $a = 18, b = 24, c = 30$ then find the values of
 - $\cos A$
 - $\sin \frac{A}{2}$
 - $\cos \frac{A}{2}$
 - $\tan \frac{A}{2}$
 - $A(\Delta ABC)$
 - $\sin A$
- In ΔABC prove that $(b + c - a) \tan \frac{A}{2} = (c + a - b) \tan \frac{B}{2} = (a + b - c) \tan \frac{C}{2}$
- In ΔABC prove that $\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} = \frac{[A(\Delta ABC)]^2}{abcs}$

3.3 Inverse Trigonometric Function :

We know that if a function $f : A \rightarrow B$ is one - one and onto then its inverse function, denoted by $f^{-1} : B \rightarrow A$, exists. For $x \in A$ and $y \in B$ if $y = f(x)$ then $x = f^{-1}(y)$.

Clearly, the domain of f^{-1} = the range of f and the range of f^{-1} = the domain of f . Trigonometric ratios defines functions, called trigonometric functions or circular dunctions. Their inverse functions are called inverse trigonometric functions or inverse cicular functions. Before finding inverse of trigonometric (circular) function , let us revise domain , range and period of the trigonometric function. We summarise them in the following table .

No trigonometric fuction is one-one. An equation of the type $\sin \theta = k$, ($|k| \leq 1$) has infinitely many solutions given by $\theta = n\pi + (-1)^n \alpha$, where $\sin \alpha = k$, $-\frac{\pi}{2} \leq \alpha \leq \frac{\pi}{2}$.

Function	Domain	Range	Period
sin	R	$[-1, 1]$	2π
cos	R	$[-1, 1]$	2π
tan	$R - \left\{ \frac{(2n+1)\pi}{2} : n \in Z \right\}$	R	π
cot	$R - \{n\pi : n \in Z\}$	R	π
sec	$R - \left\{ \frac{(2n+1)\pi}{2} : n \in Z \right\}$	$R - (-1, 1)$	2π
cosec	$R - \{n\pi : n \in Z\}$	$R - (-1, 1)$	2π

There are infinitely many elements in the domain for which the sine function takes the same value. This is true for other trigonometric functions also.

We therefore arrive at the conclusion that inverse of trigonometric functions do not exist. However from the graphs of these functions we see that there are some intervals of their domain, on which they are one-one and onto. Therefore, on these intervals we can define their inverses.

3.3.1 Inverse sine function: Consider the function $\sin : \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \rightarrow [-1, 1]$. It can be verified

from the graph that with this domain and range it is one-one and onto function. Therefore inverse sine function exists. It is denoted by

$$\sin^{-1}: [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$$

$$\text{For } x \in [-1, 1] \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right],$$

we write $\sin^{-1}x = \theta$ if $\sin \theta = x$.

Here θ is known as the principal value of $\sin^{-1}x$.

For example:

$$1) \quad \sin \frac{\pi}{6} = \frac{1}{2}, \text{ where } \frac{1}{2} \in [-1, 1] \text{ and } \frac{\pi}{6} \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$$

$$\therefore \sin^{-1} \frac{1}{2} = \frac{\pi}{6},$$

The principal value of $\sin^{-1} \frac{1}{2}$ is $\frac{\pi}{6}$,

However, though $\sin \frac{5\pi}{6} = \frac{1}{2}$,

we cannot write $\sin^{-1} \frac{1}{2} = \frac{5\pi}{6}$ as $\frac{5\pi}{6} \notin \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$




Fig 3.8(a)




Fig 3.8(b)

2) $\sin\left(-\frac{\pi}{4}\right) = -\frac{1}{\sqrt{2}}$, where $-\frac{1}{\sqrt{2}} \in [-1, 1]$ and $-\frac{\pi}{4} \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

$\therefore \sin^{-1}\left(-\frac{1}{\sqrt{2}}\right) = -\frac{\pi}{4}$

The principal value of $\sin^{-1}\left(-\frac{1}{\sqrt{2}}\right)$ is $-\frac{\pi}{4}$.

Note:

1. $\sin(\sin^{-1}x) = x$, for $x \in [-1, 1]$
2. $\sin^{-1}(\sin y) = y$, for $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

3.3.2 Inverse cosine function: Consider the function $\cos : [0, \pi] \rightarrow [-1, 1]$. It can be verified from the graph that it is a one-one and onto function. Therefore its inverse functions exist. It is denoted by \cos^{-1} .

Thus, $\cos^{-1} : [-1, 1] \rightarrow [0, \pi]$.

For $x \in [-1, 1]$ and $\theta \in [0, \pi]$, we write $\cos^{-1} x = \theta$ is $\cos \theta = x$. Here θ is known as the principal value of $\cos^{-1}x$.




Fig 3.9(a)

For example : $\cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$, where $\frac{1}{\sqrt{2}} \in [-1, 1]$ and $\frac{\pi}{4} \in [0, \pi]$

$\therefore \cos^{-1} \frac{1}{\sqrt{2}} = \frac{\pi}{4}$.

The principal value of $\cos^{-1} \frac{1}{\sqrt{2}}$ is $\frac{\pi}{4}$

Though, $\cos\left(-\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}}$,

We cannot write $\cos^{-1} \frac{1}{\sqrt{2}} = -\frac{\pi}{4}$ as $-\frac{\pi}{4} \notin [0, \pi]$

- Note:**
1. $\cos(\cos^{-1} x) = x$ for $x \in [-1, 1]$
 2. $\cos^{-1}(\cos y) = y$, for $y \in [0, \pi]$




Fig 3.9(b)

3.3.3 Inverse tangent function : Consider the function $\tan \left(-\frac{\pi}{2}, \frac{\pi}{2} \right) \rightarrow \mathbb{R}$. It can be verified from the graph that it is a one-one and onto function .




Fig 3.10(a)




Fig 3.10(b)

Therefore, its inverse function exist. It is denoted by \tan^{-1}

Thus, $\tan^{-1}: \mathbb{R} \rightarrow \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$

For $x \in \mathbb{R}$ and $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$, we write $\tan^{-1} x = \theta$ if $\tan \theta = x$

For example $\tan \frac{\pi}{4} = 1$, where $1 \in \mathbb{R}$ and $\frac{\pi}{4} \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$

$\therefore \tan^{-1} (1) = \frac{\pi}{4}$.

The principal value of $\tan^{-1} 1$ is $\frac{\pi}{4}$.

Note: 1. $\tan(\tan^{-1} x) = x$, for $x \in \mathbb{R}$ 2. $\tan^{-1} (\tan y) = y$, for $y \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$

3.3.4 Inverse cosecant function : Consider the function $\operatorname{cosec} : \left(-\frac{\pi}{2}, \frac{\pi}{2} \right) - \{0\} \rightarrow \mathbb{R} - (-1, 1)$. It can

be verified from the graph that it is a one-one and onto function. Therefore, its inverse function exists. It is denoted by $\operatorname{cosec}^{-1}$

Thus, $\operatorname{cosec}^{-1}: \mathbb{R} - (-1, 1) \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$

For $x \in \mathbb{R} - (-1, 1)$ and

$$\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\},$$

we write $\operatorname{cosec}^{-1} x = \theta$ if $\operatorname{cosec} \theta = x$

For example $\operatorname{cosec} \frac{\pi}{6} = 2,$

where $2 \in \mathbb{R} - (-1, 1)$ and $\frac{\pi}{6} \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$

$$\operatorname{cosec}^{-1}(2) = \frac{\pi}{6}$$

The principal value of $\operatorname{cosec}^{-1} 2$ is $\frac{\pi}{6}$.

- Note:**
- $\operatorname{cosec}(\operatorname{cosec}^{-1}x) = x$, for $x \in \mathbb{R} - (-1, 1)$
 - $\operatorname{cosec}^{-1}(\operatorname{cosec} y) = y$, for $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$

3.3.5 Inverse secant function : consider the function

$$\sec : [0, \pi] - \left\{\frac{\pi}{2}\right\} \rightarrow \mathbb{R} - (-1, 1)$$

It can be verified from the graph that it is a one-one and onto function. Therefore its inverse function exists. It is denoted by \sec^{-1}

$$\text{Thus, } \sec^{-1} : \mathbb{R} - (-1, 1) \rightarrow [0, \pi] - \left\{\frac{\pi}{2}\right\}.$$

For $x \in \mathbb{R} - (-1, 1)$ and $\theta \in [0, \pi] - \left\{\frac{\pi}{2}\right\},$
we write $\sec^{-1} x = \theta$ if $\sec \theta = x$

For example $\sec \pi = -1$, where $-1 \in \mathbb{R} - (-1, 1)$

$$\text{and } \pi \in [0, \pi] - \left\{\frac{\pi}{2}\right\}.$$

$$\therefore \sec^{-1}(-1) = \pi$$

The principal value of $\sec^{-1}(-1)$ is π .

- Note:**
- $\sec(\sec^{-1}x) = x$, for $x \in \mathbb{R} - (-1, 1)$
 - $\sec^{-1}(\sec y) = y$, for $y \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$




Fig 3.11(a)




Fig 3.11(b)




Fig 3.12(a)




Fig 3.12(b)




Fig 3.13(a)

3.4.6 Inverse cotangent function : Consider the function $\cot : (0, \pi) \rightarrow \mathbb{R}$. It can be verified from the graph that it is a one-one and onto function. Therefore, its inverse exists. It is denoted by \cot^{-1} .

Thus, $\cot^{-1} : \mathbb{R} \rightarrow (0, \pi)$

For $x \in \mathbb{R}$ and $\theta \in (0, \pi)$, we write $\cot^{-1} x = \theta$ if $\cot \theta = x$

For example: $\cot \frac{\pi}{3} = \frac{1}{\sqrt{3}}$, where $\frac{1}{\sqrt{3}} \in \mathbb{R}$
and $\frac{\pi}{3} \in (0, \pi)$

$$\therefore \cot^{-1} \left(\frac{1}{\sqrt{3}} \right) = \frac{\pi}{3}$$

The principal value of $\cot^{-1} \left(\frac{1}{\sqrt{3}} \right)$ is $\frac{\pi}{3}$.

- Note:**
1. $\cot(\cot^{-1}x) = x$, for $x \in \mathbb{R}$
 2. $\cot^{-1}(\cot y) = y$, for $y \in (0, \pi)$




Fig 3.13(b)

3.4.7 Principal Values of Inverse Trigonometric Functions :

The following table shows domain and range of all inverse trigonometric functions. The value of function in the range is called the principal value of the function.

$\sin^{-1} : [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$\cos^{-1} : [-1, 1] \rightarrow [0, \pi]$
$\tan^{-1} : \mathbb{R} \rightarrow \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$\operatorname{cosec}^{-1} : \mathbb{R} - (-1, 1) \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
$\sec^{-1} : \mathbb{R} - (-1, 1) \rightarrow [0, \pi] - \left\{\frac{\pi}{2}\right\}$

3.4.8 Properties of inverse trigonometric functions :

i) If $-1 \leq x \leq 1$ and $x \neq 0$ then $\sin^{-1} x = \operatorname{cosec}^{-1} \left(\frac{1}{x}\right)$

Proof : By the conditions on x , $\sin^{-1} x$ and $\operatorname{cosec}^{-1} \left(\frac{1}{x}\right)$ are defined.

As $-1 \leq x \leq 1$ and $x \neq 0$, $\frac{1}{x} \in \mathbb{R} - (-1, 1) \dots(1)$

Let $\sin^{-1} x = \theta$

$\therefore \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ and $\theta \neq 0$ (as $x \neq 0$)

$\therefore \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\} \dots(2)$

$\therefore \sin \theta = x$

$\therefore \operatorname{cosec} \theta = \frac{1}{x} \dots(3)$

From (1), (2) and (3) we get

$\operatorname{cosec}^{-1} \left(\frac{1}{x}\right) = \theta$

$\therefore \theta = \operatorname{cosec}^{-1} \left(\frac{1}{x}\right)$

$\therefore \sin^{-1} x = \operatorname{cosec}^{-1} \left(\frac{1}{x}\right)$

Similarly we can prove the following result.

(i) $\cos^{-1} x = \sec^{-1} \left(\frac{1}{x}\right)$ if $-1 \leq x \leq 1$ and $x \neq 0$

$$(ii) \quad \tan^{-1} x = \cot^{-1} \left(\frac{1}{x} \right) \quad \text{if } x > 0$$

Proof : Let $\tan^{-1} x = \theta$

$$\therefore \theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right). \text{ As } x > 0, \theta \in \left(0, \frac{\pi}{2} \right)$$

$$\therefore \tan \theta = x$$

$$\therefore \cot \theta = \frac{1}{x}, \text{ where } \frac{1}{x} \in \mathbb{R} \dots(1)$$

$$\text{As } \theta \in \left(0, \frac{\pi}{2} \right) \text{ and } \left(0, \frac{\pi}{2} \right) \subset (0, \pi) \dots(2)$$

$$\text{From (1) and (2) we get } \cot^{-1} \left(\frac{1}{x} \right) = \theta$$

$$\therefore \theta = \cot^{-1} \left(\frac{1}{x} \right)$$

$$\therefore \tan^{-1} x = \cot^{-1} \left(\frac{1}{x} \right)$$

(iii) Similarly we can prove that : $\tan^{-1} x = -\pi + \cot^{-1} \left(\frac{1}{x} \right)$ if $x < 0$.

Activity : Verify the above result for $x = -\sqrt{3}$

iv) if $-1 \leq x \leq 1$ then $\sin^{-1}(-x) = -\sin^{-1}(x)$

Proof : As $-1 \leq x \leq 1, x \in [-1, 1]$

$$\therefore -x \in [-1, 1] \dots(1)$$

Let $\sin^{-1}(-x) = \theta$

$$\therefore \theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \text{ and } \sin \theta = x$$

Now $\sin(-\theta) = -\sin \theta = -x$

$$\therefore \sin(-\theta) = -x \dots(2)$$

Also from (1) $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$

$$\therefore \frac{\pi}{2} \geq -\theta \geq -\frac{\pi}{2}$$

$$\therefore -\frac{\pi}{2} \leq -\theta \leq \frac{\pi}{2}$$

$$\therefore -\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right] \dots(3)$$

From (1), (2) and (3) we can write

$$\therefore \sin^{-1}(-x) = -\theta$$

$$\therefore \sin^{-1}(-x) = -\sin^{-1}(x)$$

Similarly we can prove the following results.

v) If $-1 \leq x \leq 1$ then $\cos^{-1}(-x) = \pi - \cos^{-1}(x)$

vi) For all $x \in \mathbb{R}$, $\tan^{-1}(-x) = -\tan^{-1}(x)$

vii) If $|x| \geq 1$ then $\operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1}(x)$

viii) If $|x| \geq 1$ then $\sec^{-1}(-x) = \pi - \sec^{-1}(x)$

ix) For all $x \in \mathbb{R}$, $\cot^{-1}(-x) = \pi - \cot^{-1}(x)$

x) If $-1 \leq x \leq 1$ then $\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}$

Proof : Let $\sin^{-1} x = \theta$, where $x \in [-1, 1]$ and $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

$$\therefore -\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

$$\therefore \frac{\pi}{2} - \theta \in [0, \pi], \text{ the principal domain of the cosine function.}$$

$$\therefore \cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = x$$

$$\therefore \cos^{-1} x = \frac{\pi}{2} - \theta$$

$$\therefore \theta + \cos^{-1} x = \frac{\pi}{2}$$

$$\therefore \sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

Similarly we can prove the following results.

xi) For $x \in \mathbb{R}$, $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$

xii) For $x \geq 1$, $\operatorname{cosec}^{-1} x + \sec^{-1} x = \frac{\pi}{2}$

xiii) If $x > 0$, $y > 0$ and $xy < 1$ then

$$\tan^{-1} x + \tan^{-1} y = \tan^{-1} \left(\frac{x+y}{1-xy} \right)$$

Proof : Let $\tan^{-1} x = \theta$ and $\tan^{-1} y = \phi$

$$\therefore \tan \theta = x \text{ and } \tan \phi = y$$

As $x > 0$ and $y > 0$, we have $0 < \theta < \frac{\pi}{2}$ and $0 < \phi < \frac{\pi}{2}$

$$\therefore 0 < \theta + \phi < \pi \quad \dots(1)$$

$$\text{Also } \tan(\theta + \phi) = \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi} = \frac{x + y}{1 - xy}$$

As $x > 0, y > 0$ and $xy < 1$, x, y and $1 - xy$ are all positive.

$$\therefore \frac{x + y}{1 - xy} \text{ is positive.}$$

$\tan(\theta + \phi)$ is positive.(2)

From (1) and (2) we get $(\theta + \phi) \in \left(0, \frac{\pi}{2}\right)$, the part of the principal domain of the tangent function.

$$\therefore \theta + \phi = \tan^{-1} \left(\frac{x + y}{1 - xy} \right)$$

$$\therefore \tan^{-1} x + \tan^{-1} y = \tan^{-1} \left(\frac{x + y}{1 - xy} \right)$$

Similarly we can prove the following results.

xiv) If $x > 0, y > 0$ and $xy > 1$ then

$$\tan^{-1} x + \tan^{-1} y = \pi + \tan^{-1} \left(\frac{x + y}{1 - xy} \right)$$

xv) If $x > 0, y > 0$ and $xy = 1$ then

$$\tan^{-1} x + \tan^{-1} y = \frac{\pi}{2}$$

xvi) If $x > 0, y > 0$ then

$$\tan^{-1} x - \tan^{-1} y = \tan^{-1} \left(\frac{x - y}{1 + xy} \right)$$

Ex.(1) Find the principal values of the following :

$$(i) \sin^{-1} \left(-\frac{1}{2} \right) \quad (ii) \cos^{-1} \left(\frac{\sqrt{3}}{2} \right) \quad (iii) \cot^{-1} \left(-\frac{1}{\sqrt{3}} \right)$$

Solution : (i) $\sin^{-1} \left(-\frac{1}{2} \right)$

$$\text{We have, } \sin \left(-\frac{1}{2} \right) = -\frac{1}{2}, \text{ where } -\frac{1}{2} \in [-1, 1] \text{ and } -\frac{\pi}{6} \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$$

$$\therefore \sin^{-1} \left(-\frac{1}{2} \right) = -\frac{\pi}{6}$$

The principal value of $\sin^{-1} \left(-\frac{1}{2} \right)$ is $-\frac{\pi}{6}$.

$$(ii) \cos^{-1} \left(\frac{\sqrt{3}}{2} \right)$$

We have, $\cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$, where $\frac{\sqrt{3}}{2} \in [-1, 1]$ and $\frac{\pi}{6} \in [0, \pi]$

$$\cos^{-1} \left(\frac{\sqrt{3}}{2} \right) = \frac{\pi}{6}$$

The principal value of $\cos^{-1} \left(\frac{\sqrt{3}}{2} \right)$ is $\frac{\pi}{6}$.

$$(iii) \cot^{-1} \left(-\frac{1}{\sqrt{3}} \right)$$

We have, $\cot \frac{2\pi}{3} = -\frac{1}{\sqrt{3}}$, where $-\frac{1}{\sqrt{3}} \in \mathbb{R}$ and $\frac{2\pi}{3} \in (0, \pi)$

\therefore The principal value of $\cot^{-1} \left(-\frac{1}{\sqrt{3}} \right)$ is $\frac{2\pi}{3}$

Ex.(2) Find the values of the following

$$(i) \sin^{-1} \left(\sin \frac{5\pi}{3} \right) \quad (ii) \tan^{-1} \left(\tan \frac{\pi}{4} \right)$$

$$(iii) \sin \left(\cos^{-1} \left(-\frac{1}{\sqrt{2}} \right) \right)$$

$$(iv) \sin \left(\cos^{-1} \frac{4}{5} + \tan^{-1} \frac{5}{12} \right)$$

Solution : (i) $\sin^{-1} \left(\sin \left(\frac{5\pi}{3} \right) \right) = \sin^{-1} \left(\sin \left(2\pi - \frac{\pi}{3} \right) \right)$

$$= \sin^{-1} \left(\sin \left(-\frac{\pi}{3} \right) \right) = -\frac{\pi}{3}, \text{ as } -\frac{\pi}{3} \in \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$$

$$\therefore \sin^{-1} \left(\sin \frac{5\pi}{3} \right) = -\frac{\pi}{3}$$

$$(ii) \tan^{-1} \left(\tan \frac{\pi}{4} \right) = \frac{\pi}{4}, \text{ as } \frac{\pi}{4} \in \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$$

(iii) We have $\cos^{-1}(-x) = \pi - \cos^{-1} x$

$$\therefore \sin \left(\cos^{-1} \left(-\frac{1}{\sqrt{2}} \right) \right) = \sin \left(\pi - \cos^{-1} \frac{1}{\sqrt{2}} \right)$$

$$= \sin \left(\pi - \frac{\pi}{4} \right) = \sin \left(\frac{3\pi}{4} \right) = \frac{1}{\sqrt{2}}$$

$$\therefore \sin \left(\cos^{-1} \left(-\frac{1}{\sqrt{2}} \right) \right) = \frac{1}{\sqrt{2}}$$

(iv) Let $\cos^{-1} \frac{4}{5} = \theta$ and $\tan^{-1} \frac{5}{12} = \phi$

$$\therefore \cos \theta = \frac{4}{5} \text{ and } \tan \phi = \frac{5}{12}$$

$$\therefore \sin \theta = \frac{3}{5} \text{ and } \sin \phi = \frac{5}{13}, \cos \phi = \frac{12}{13}$$

$$\sin \left(\cos^{-1} \frac{4}{5} + \tan^{-1} \frac{5}{12} \right) = \sin (\theta + \phi)$$

$$= \sin \theta \cos \phi + \cos \theta \sin \phi$$

$$= \frac{3}{5} \frac{12}{13} + \frac{4}{5} \frac{5}{13} = \frac{56}{65}$$

$$\therefore \sin \left(\cos^{-1} \frac{4}{5} + \tan^{-1} \frac{5}{12} \right) = \frac{56}{65}$$

Ex.(3) Find the values of the following :

(i) $\sin \left[\sin^{-1} \left(\frac{3}{5} \right) + \cos^{-1} \left(\frac{3}{5} \right) \right]$

(ii) $\cos \left[\cos^{-1} \left(-\frac{1}{2} \right) + \tan^{-1} \sqrt{3} \right]$

Solution : (i) We have if $-1 \leq x \leq 1$ then

$$\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

Here $-1 < \frac{3}{5} < 1$

$$\therefore \sin^{-1} \left(\frac{3}{5} \right) + \cos^{-1} \left(\frac{3}{5} \right) = \frac{\pi}{2}$$

$$\sin \left[\sin^{-1} \left(\frac{3}{5} \right) + \cos^{-1} \left(\frac{3}{5} \right) \right] = \sin \left(\frac{\pi}{2} \right) = 1.$$

$$(ii) \quad \cos^{-1} \left(-\frac{1}{2} \right) = \frac{2\pi}{3} \text{ and } \tan^{-1} \sqrt{3} = \frac{\pi}{3}.$$

$$\therefore \cos \left[\cos^{-1} \left(-\frac{1}{2} \right) + \tan^{-1} \sqrt{3} \right]$$

$$= \cos \left(\frac{2\pi}{3} + \frac{\pi}{3} \right)$$

$$= \cos \pi$$

$$= -1.$$

Ex.(4) If $|x| \leq 1$, show that
 $\sin(\cos^{-1}x) = \cos(\sin^{-1}x).$

Solution :

We have for $|x| \leq 1$.

$$\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

$$\therefore \cos^{-1} x = \frac{\pi}{2} - \sin^{-1} x,$$

Now using $\sin \left(\frac{\pi}{2} - \theta \right) = \cos \theta$

$$\text{We have } \sin(\cos^{-1}x) = \sin \left(\frac{\pi}{2} - \sin^{-1} x \right)$$

$$= \cos(\sin^{-1}x)$$

$$\therefore \sin(\cos^{-1}x) = \cos(\sin^{-1}x)$$

Ex.(5) Prove the following

$$(i) \quad 2\tan^{-1} \left(-\frac{1}{3} \right) + \cos^{-1} \left(\frac{3}{5} \right) = \frac{\pi}{2}$$

$$(ii) \quad 2\tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{7} \right) = \frac{\pi}{4}$$

Solution :

$$(i) \quad 2\tan^{-1} \left(\frac{1}{3} \right) = \tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{3} \right), \text{ as } xy = \frac{1}{3} \times \frac{1}{3} = \frac{1}{9} < 1$$

$$= \tan^{-1} \left(\frac{\frac{1}{3} + \frac{1}{3}}{1 - \frac{1}{3} \cdot \frac{1}{3}} \right) = \tan^{-1} \frac{3}{4} = \theta, \text{ (say)}$$

$$\therefore \tan \theta = \frac{3}{4} \quad \therefore 0 < \theta < \frac{\pi}{2}$$

$$\therefore \sin \theta = \frac{3}{5} \quad \therefore \theta = \sin^{-1} \frac{3}{5}$$

$$\therefore 2 \tan^{-1} \frac{1}{3} = \tan^{-1} \frac{3}{4} = \sin^{-1} \frac{3}{5}$$

$$\therefore 2 \tan^{-1} \frac{1}{3} + \cos^{-1} \frac{3}{5} = \sin^{-1} \frac{3}{5} + \cos^{-1} \frac{3}{5} = \frac{\pi}{2}$$

(ii) $2 \tan^{-1} \frac{1}{3} = \tan^{-1} \frac{3}{4}$ as seen in (i)

$$\therefore 2 \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{7} = \tan^{-1} \frac{3}{4} + \tan^{-1} \frac{1}{7} \text{ and } xy = \frac{3}{4} \times \frac{1}{7} = \frac{3}{28} < 1$$

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{1}{7}}{1 - \frac{3}{4} \cdot \frac{1}{7}} \right) = \tan^{-1} 1 = \frac{\pi}{4}$$

Ex.(6) Prove that $\tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3 = \pi$

Solution: We use the result:

$$\tan^{-1} x + \tan^{-1} y = \pi + \tan^{-1} \left(\frac{x+y}{1-xy} \right) \text{ if } xy > 1$$

$$\text{Here } xy = 1 \times 2 = 2 > 1$$

$$\therefore \tan^{-1} 1 + \tan^{-1} 2 = \pi + \tan^{-1} \left(\frac{1+2}{1-(1)(2)} \right)$$

$$= \pi + \tan^{-1} \left(\frac{3}{1-2} \right)$$

$$= \pi + \tan^{-1} (-3)$$

$$= \pi - \tan^{-1} 3 \text{ (As, } \tan^{-1}(-x) = -\tan^{-1} x)$$

$$\therefore \tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3 = \pi$$

Ex.(7) Prove that $\cos^{-1} \frac{4}{5} + \cos^{-1} \frac{12}{13} = \cos^{-1} \frac{33}{65}$

Solution: Let $\cos^{-1} \frac{4}{5} = \theta$

$$\text{Then } 0 < \theta < \frac{\pi}{2} \text{ and } \cos \theta = \frac{4}{5}$$

$$\therefore \sin \theta = \frac{3}{5}$$

Let $\cos^{-1} \frac{12}{13} = \phi$

Then $0 < \phi < \frac{\pi}{2}$ and $\cos \phi = \frac{12}{13}$

$$\sin \phi = \frac{5}{13}$$

$$\cos(\theta + \phi) = \cos \theta \cos \phi - \sin \theta \sin \phi$$

$$= \left(\frac{4}{5}\right)\left(\frac{12}{13}\right) - \left(\frac{3}{5}\right)\left(\frac{5}{13}\right)$$

$$= \frac{48}{65} - \frac{15}{65} = \frac{33}{65}$$


$$\therefore \cos(\theta + \phi) = \frac{33}{65} \quad \dots(1)$$

Also $0 < \theta < \frac{\pi}{2}$ and $0 < \phi < \frac{\pi}{2}$

$$\therefore 0 < \theta + \phi < \pi.$$

$$\therefore \text{from (1), } \theta + \phi = \cos^{-1} \frac{33}{65}$$

$$\therefore \cos^{-1} \frac{4}{5} + \cos^{-1} \frac{12}{13} = \cos^{-1} \frac{33}{65}$$

 **Exercise 3.3**

1) Find the principal values of the following :

(i) $\sin^{-1} \left(\frac{1}{2}\right)$ (ii) $\operatorname{cosec}^{-1}(2)$ (iii) $\tan^{-1}(-1)$

(iv) $\tan^{-1}(-\sqrt{3})$ (v) $\sin^{-1} \left(\frac{1}{\sqrt{2}}\right)$ (vi) $\cos^{-1} \left(-\frac{1}{2}\right)$

2) Evaluate the following :

(i) $\tan^{-1}(1) + \cos^{-1} \left(\frac{1}{2}\right) + \sin^{-1} \left(\frac{1}{2}\right)$

(ii) $\cos^{-1} \left(\frac{1}{2}\right) + 2 \sin^{-1} \left(\frac{1}{2}\right)$

(iii) $\tan^{-1} \sqrt{3} - \sec^{-1}(-2)$

$$(iv) \operatorname{cosec}^{-1}(-\sqrt{2}) + \cot^{-1}(\sqrt{3})$$

3) Prove the following :

$$(i) \sin^{-1}\left(-\frac{1}{\sqrt{2}}\right) - 3 \sin^{-1}\left(\frac{\sqrt{3}}{2}\right) = -\frac{3\pi}{4}$$

$$(ii) \sin^{-1}\left(-\frac{1}{2}\right) + \cos^{-1}\left(-\frac{\sqrt{3}}{2}\right) = \cos^{-1}\left(-\frac{1}{2}\right)$$

$$(iii) \sin^{-1}\left(\frac{3}{5}\right) + \cos^{-1}\left(\frac{12}{13}\right) = \sin^{-1}\left(\frac{56}{65}\right)$$

$$(iv) \cos^{-1}\left(\frac{3}{5}\right) + \cos^{-1}\left(\frac{4}{5}\right) = \frac{\pi}{2}$$

$$(v) \tan^{-1}\left(\frac{1}{2}\right) + \tan^{-1}\left(\frac{1}{3}\right) = \frac{\pi}{4}$$

$$(vi) 2 \tan^{-1}\left(\frac{1}{3}\right) = \tan^{-1}\left(\frac{3}{4}\right)$$

$$(vii) \tan^{-1}\left[\frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta}\right] = \frac{\pi}{4} + \theta \text{ if } \theta \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$$

$$(viii) \tan^{-1}\sqrt{\frac{1-\cos \theta}{1+\cos \theta}} = \frac{\theta}{2}, \text{ if } \theta \in (-\pi, \pi)$$



Let's remember!

- * An equation involving trigonometric function (or functions) is called a trigonometric equation.
- * A value of α variable in a trigonometric equation which satisfies the equation is called a solution of the trigonometric equation.
- * A solution α of a trigonometric equation is called a principal solution if $0 \leq \alpha < 2\pi$.
- * The solution of a trigonometric equation which is generalized by using its periodicity is called the general solution.
- * The general solution of $\sin \theta = \sin \alpha$ is $\theta = n\pi + (-1)^n \alpha$, where $n \in \mathbb{Z}$.
- * The general solution of $\cos \theta = \cos \alpha$ is $\theta = 2n\pi + \alpha$, where $n \in \mathbb{Z}$.
- * The general solution of $\tan \theta = \tan \alpha$ is $\theta = n\pi + \alpha$, where $n \in \mathbb{Z}$.
- * The general solution of $\sin^2 \theta = \sin^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.
- * The general solution of $\cos^2 \theta = \cos^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.
- * The general solution of $\tan^2 \theta = \tan^2 \alpha$ is $\theta = n\pi \pm \alpha$, where $n \in \mathbb{Z}$.

* The Sine Rule : In ΔABC , $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$, where R is the circumradius of ΔABC .

Following are the different forms of the Sine rule.

(i) $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R$

(ii) $a = 2R \sin A$, $b = 2R \sin B$, $c = 2R \sin C$

(iii) $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} = k$

(iv) $b \sin A = a \sin B$, $c \sin B = b \sin C$, $c \sin A = a \sin C$

(v) $\frac{a}{b} = \frac{\sin A}{\sin B}$, $\frac{b}{c} = \frac{\sin B}{\sin C}$

* The Cosine Rule : In ΔABC ,

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = c^2 + a^2 - 2ca \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

* The Projection Rule : In ΔABC .

$$a = b \cos C + c \cos B$$

$$b = c \cos A + a \cos C$$

$$c = a \cos B + b \cos A$$

* Half angle formulae : In ΔABC , if $a + b + c = 2s$ then

(i) $\sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}}$, $\sin \frac{B}{2} = \sqrt{\frac{(s-a)(s-c)}{ac}}$, $\sin \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{ab}}$

(ii) $\cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}$, $\cos \frac{B}{2} = \sqrt{\frac{s(s-b)}{ac}}$, $\cos \frac{C}{2} = \sqrt{\frac{s(s-c)}{ab}}$

(iii) $\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$, $\tan \frac{B}{2} = \sqrt{\frac{(s-a)(s-c)}{s(s-b)}}$, $\tan \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{s(s-c)}}$

* Heron's Formula : If a,b,c are sides of ΔABC and $a + b + c = 2s$ then

$$A(\Delta ABC) = \sqrt{s(s-a)(s-b)(s-c)}$$

* Napier's Analogy : In ΔABC , $\tan \left(\frac{B-C}{2} \right) = \frac{(b-c)}{(b+c)} \cot \frac{A}{2}$

* Inverse Trigonometric functions :

(i) $\sin(\sin^{-1}x) = x$, for $x \in [-1,1]$

- (ii) $\sin^{-1}(\sin y) = y$, for $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
- (iii) $\cos(\cos^{-1}x) = x$, for $x \in [-1, 1]$
- (iv) $\cos^{-1}(\cos y) = y$, for $y \in [0, \pi]$
- (v) $\tan(\tan^{-1}x) = x$, for $x \in \mathbb{R}$
- (vi) $\tan^{-1}(\tan y) = y$, for $y \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
- (vii) $\operatorname{cosec}(\operatorname{cosec}^{-1}x) = x$, for $x \in \mathbb{R} - (-1, 1)$
- (viii) $\operatorname{cosec}^{-1}(\operatorname{cosec} y) = y$, for $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
- (ix) $\sec(\sec^{-1}x) = x$, for $x \in \mathbb{R} - (-1, 1)$
- (x) $\sec^{-1}(\sec y) = y$, for $y \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$
- (xi) $\cot(\cot^{-1}x) = x$, for $x \in \mathbb{R}$
- (xii) $\cot^{-1}(\cot y) = y$, for $y \in (0, \pi)$

*** Properties of inverse trigonometric functions :**

- (i) $\sin^{-1}x = \operatorname{cosec}^{-1}\left(\frac{1}{x}\right)$ if $-1 \leq x \leq 1$ and $x \neq 0$
- (ii) $\cos^{-1}x = \sec^{-1}\left(\frac{1}{x}\right)$ if $-1 \leq x \leq 1$ and $x \neq 0$
- (iii) $\tan^{-1}x = \cot^{-1}\left(\frac{1}{x}\right)$ if $x > 0$
- (iv) If $-1 \leq x \leq 1$ then $\sin^{-1}(-x) = -\sin^{-1}(x)$
- (v) If $-1 \leq x \leq 1$ then $\cos^{-1}(-x) = \pi - \cos^{-1}(x)$
- (vi) For all $x \in \mathbb{R}$, $\tan^{-1}(x) = -\tan^{-1}(-x)$
- (vii) If $|x| \geq 1$ then $\operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1}(x)$
- (viii) If $|x| \geq 1$ then $\sec^{-1}(x) = \pi - \sec^{-1}(-x)$
- (ix) For all $x \in \mathbb{R}$, $\cot^{-1}(-x) = \pi - \cot^{-1}(x)$
- (x) If $-1 \leq x \leq 1$ then $\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}$
- (xi) For $x \in \mathbb{R}$, $\tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}$
- (xii) For $x \geq 1$, $\operatorname{cosec}^{-1}x + \sec^{-1}x = \frac{\pi}{2}$
- (xiii) If $x > 0$, $y > 0$ and $xy < 1$ then $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\frac{x+y}{1-xy}$

(xiv) If $x > 0, y > 0$ and $xy > 1$ then $\tan^{-1}x + \tan^{-1}y = \pi + \tan^{-1} \frac{x+y}{1-xy}$

(xv) If $x > 0, y > 0$ and $xy = 1$ then $\tan^{-1}x + \tan^{-1}y = \frac{\pi}{2}$

(xvi) If $x > 0, y > 0$ then $\tan^{-1}x - \tan^{-1}y = \tan^{-1} \left(\frac{x-y}{1+xy} \right)$

Miscellaneous Exercise 3

I) Select the correct option from the given alternatives.

1) The principal of solutions equation $\sin\theta = \frac{-1}{2}$ are _____.

- a) $\frac{5\pi}{6}, \frac{\pi}{6}$ b) $\frac{7\pi}{6}, \frac{11\pi}{6}$ c) $\frac{\pi}{6}, \frac{7\pi}{6}$ d) $\frac{7\pi}{6}, \frac{\pi}{6}$

2) The principal solution of equation $\cot\theta = \sqrt{3}$ _____.

- a) $\frac{\pi}{6}, \frac{7\pi}{6}$ b) $\frac{\pi}{6}, \frac{5\pi}{6}$
 c) $\frac{\pi}{6}, \frac{8\pi}{6}$ d) $\frac{7\pi}{6}, \frac{\pi}{6}$

3) The general solution of $\sec x = \sqrt{2}$ is _____.

- a) $2n\pi \pm \frac{\pi}{4}, n \in Z$ b) $2n\pi \pm \frac{\pi}{2}, n \in Z$
 c) $n\pi \pm \frac{\pi}{2}, n \in Z$ d) $2n\pi \pm \frac{\pi}{3}, n \in Z$

4) If $\cos p\theta = \cos q\theta, p \neq q$ then _____.

- a) $\theta = \frac{2n\pi}{p \pm q}$ b) $\theta = 2n\pi$
 c) $\theta = 2n\pi \pm p$ d) $n\pi \pm q$

5) If polar co-ordinates of a point are $\left(2, \frac{\pi}{4}\right)$ then its cartesian co-ordinates are _____.

- a) $(2, \sqrt{2})$ b) $(\sqrt{2}, 2)$
 c) $(2, 2)$ d) $(\sqrt{2}, \sqrt{2})$

6) If $\sqrt{3} \cos x - \sin x = 1$, then general value of x is _____.

- a) $2n\pi \pm \frac{\pi}{3}$ b) $2n\pi \pm \frac{\pi}{6}$
 c) $2n\pi \pm \frac{\pi}{3} - \frac{\pi}{6}$ d) $n\pi + (-1)^n \frac{\pi}{3}$

- 7) In ΔABC if $\angle A = 45^\circ$, $\angle B = 60^\circ$ then the ratio of its sides are _____.
- a) $2 : \sqrt{2} : \sqrt{3} + 1$ b) $\sqrt{2} : 2 : \sqrt{3} + 1$
 c) $2\sqrt{2} : \sqrt{2} : \sqrt{3}$ d) $2 : 2\sqrt{2} : \sqrt{3} + 1$
- 8) In ΔABC , if $c^2 + a^2 - b^2 = ac$, then $\angle B =$ _____.
- a) $\frac{\pi}{4}$ b) $\frac{\pi}{3}$ c) $\frac{\pi}{2}$ d) $\frac{\pi}{6}$
- 9) In ΔABC , $ac \cos B - bc \cos A =$ _____.
- a) $a^2 - b^2$ b) $b^2 - c^2$
 c) $c^2 - a^2$ d) $a^2 - b^2 - c^2$
- 10) If in a triangle, the angles are in A.P. and $b : c = \sqrt{3} : \sqrt{2}$ then $\angle A$ is equal to _____.
- a) 30° b) 60°
 c) 75° d) 45°
- 11) $\cos^{-1} \left(\cos \frac{7\pi}{6} \right) =$ _____.
- a) $\frac{7\pi}{6}$ b) $\frac{5\pi}{6}$
 c) $\frac{\pi}{6}$ d) $\frac{3\pi}{2}$
- 11) The value of $\cot(\tan^{-1} 2x + \cot^{-1} 2x)$ is _____.
- a) 0 b) $2x$
 c) $\pi + 2x$ d) $\pi - 2x$
- 12) The principal value of $\sin^{-1} \left(-\frac{\sqrt{3}}{2} \right)$ is _____.
- a) $\left(-\frac{2\pi}{3} \right)$ b) $\frac{4\pi}{3}$
 c) $\frac{5\pi}{3}$ d) $-\frac{\pi}{3}$
- 13) If $\sin^{-1} \frac{4}{5} + \cos^{-1} \frac{12}{13} = \sin^{-1} \alpha$, then $\alpha =$ _____.
- a) $\frac{63}{65}$ b) $\frac{62}{65}$
 c) $\frac{61}{65}$ d) $\frac{60}{65}$
- 14) If $\tan^{-1}(2x) + \tan^{-1}(3x) = \frac{\pi}{4}$, then $x =$
- a) -1 b) $\frac{1}{6}$ c) $\frac{2}{6}$ d) $\frac{3}{2}$

- 15) $2 \tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{7} \right) =$ _____.
- a) $\tan^{-1} \left(\frac{4}{5} \right)$ b) $\frac{\pi}{2}$
 c) 1 d) $\frac{\pi}{4}$
- 16) $\tan \left(2 \tan^{-1} \left(\frac{1}{5} \right) - \frac{\pi}{4} \right) =$ _____.
- a) $\frac{17}{7}$ b) $-\frac{17}{7}$
 c) $\frac{7}{17}$ d) $-\frac{7}{17}$
- 17) The principal value branch of $\sec^{-1} x$ is _____.
- a) $\left[-\frac{\pi}{2}, \frac{\pi}{2} \right] - \{0\}$ b) $[0, \pi] - \left[\frac{\pi}{2} \right]$
 c) $(0, \pi)$ d) $\left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$
- 18) $\cos \left[\tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{2} \right] =$ _____.
- a) $\frac{1}{\sqrt{2}}$ b) $\frac{\sqrt{3}}{2}$ c) $\frac{1}{2}$ d) $\frac{\pi}{4}$
- 19) If $\tan \theta + \tan 2\theta + \tan 3\theta = \tan \theta \tan 2\theta \tan 3\theta$, then the general value of the θ is _____.
- a) $n\pi$ b) $\frac{n\pi}{6}$
 c) $n\pi \pm \frac{\pi}{4}$ d) $\frac{n\pi}{2}$
- 20) If any ΔABC , if $a \cos B = b \cos A$, then the triangle is _____.
- a) Equilateral triangle b) Isosceles triangle
 c) Scalene d) Right angled

II: Solve the following

- 1) Find the principal solutions of the following equations :
- (i) $\sin 2\theta = -\frac{1}{2}$ (ii) $\tan 3\theta = -1$ (iii) $\cot \theta = 0$
- 2) Find the principal solutions of the following equations :
- (i) $\sin 2\theta = -\frac{1}{\sqrt{2}}$ (ii) $\tan 5\theta = -1$ (iii) $\cot 2\theta = 0$

3) Which of the following equations have no solutions ?

(i) $\cos 2\theta = \frac{1}{3}$ (ii) $\cos^2 \theta = -1$ (iii) $2 \sin \theta = 3$ (iv) $3 \sin \theta = 5$

4) Find the general solutions of the following equations :

i) $\tan \theta = -\sqrt{3}$ ii) $\tan^2 \theta = 3$ iii) $\sin \theta - \cos \theta = 1$ iv) $\sin^2 \theta - \cos^2 \theta = 1$

5) In ΔABC prove that $\cos \left(\frac{A-B}{2} \right) = \left(\frac{a+b}{c} \right) \sin \frac{C}{2}$

6) With usual notations prove that $\frac{\sin (A-B)}{\sin (A+B)} = \frac{a^2 - b^2}{c^2}$.

7) In ΔABC prove that $(a-b)^2 \cos^2 \frac{C}{2} + (a+b)^2 \sin^2 \frac{C}{2} = c^2$

8) In ΔABC if $\cos A = \sin B - \cos C$ then show that it is a right angled triangle.

9) If $\frac{\sin A}{\sin C} = \frac{\sin(A-B)}{\sin(B-C)}$ then show that a^2, b^2, c^2 , are in A.P.

10) Solve the triangle in which $a = \sqrt{3} + 1$, $b = \sqrt{3} - 1$ and $C = 60^\circ$

11) In ΔABC prove the following :

(i) $a \sin A - b \sin B = c \sin (A - B)$

(ii) $\frac{c - b \cos A}{b - c \cos A} = \frac{\cos B}{\cos C}$

(iii) $a^2 \sin (B - C) = (b^2 - c^2) \sin A$

(iv) $ac \cos B - bc \cos A = (a^2 - b^2)$

(v) $\frac{\cos A}{a} + \frac{\cos B}{b} + \frac{\cos C}{c} = \frac{a^2 + b^2 + c^2}{2abc}$

(vi) $\frac{\cos 2A}{a^2} - \frac{\cos 2B}{b^2} = \frac{1}{a^2} - \frac{1}{b^2}$

(vii) $\frac{b-c}{a} = \frac{\tan \frac{B}{2} - \tan \frac{C}{2}}{\tan \frac{B}{2} + \tan \frac{C}{2}}$

12) In ΔABC if a^2, b^2, c^2 , are in A.P. then $\cot \frac{A}{2}, \cot \frac{B}{2}, \cot \frac{C}{2}$ are also in A.P.

13) In ΔABC if $C = 90^\circ$ then prove that $\sin(A - B) = \frac{a^2 - b^2}{a^2 + b^2}$

- 14) In ΔABC if $\frac{\cos A}{a} = \frac{\cos B}{b}$ then show that it is an isosceles triangle.
- 15) In ΔABC if $\sin^2 A + \sin^2 B = \sin^2 C$ then prove that the triangle is a right angled triangle.
- 16) In ΔABC prove that $a^2 (\cos^2 B - \cos^2 C) + b^2 (\cos^2 C - \cos^2 A) + c^2 (\cos^2 A - \cos^2 B) = 0$
- 17) With usual notations show that $(c^2 - a^2 + b^2) \tan A = (a^2 - b^2 + c^2) \tan B = (b^2 - c^2 + a^2) \tan C$
- 18) In ΔABC if $a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2} = \frac{3b}{2}$ then prove that a, b, c are in A.P.
- 19) Show that $2 \sin^{-1} \frac{3}{5} = \tan^{-1} \frac{24}{7}$
- 20) Show that $\tan^{-1} \frac{1}{5} + \tan^{-1} \frac{1}{7} + \tan^{-1} \frac{1}{3} + \tan^{-1} \frac{1}{8} = \frac{\pi}{4}$
- 21) Prove that $\tan^{-1} \sqrt{x} = \frac{1}{2} \cos^{-1} \left(\frac{1-x}{1+x} \right)$ if $x \in [0, 1]$
- 22) Show that $\frac{9\pi}{5} - \frac{9}{4} \sin^{-1} \frac{1}{3} = \frac{9}{4} \sin^{-1} \frac{2\sqrt{2}}{3}$
- 23) Show that $\tan^{-1} \left(\frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt{1+x} + \sqrt{1-x}} \right) = \frac{\pi}{4} - \frac{1}{2} \cos^{-1} x$, for $-\frac{1}{\sqrt{2}} \leq x \leq 1$
- 24) If $\sin(\sin^{-1} \frac{1}{5} + \cos^{-1} x) = 1$ then find the value of x .
- 25) If $\tan^{-1} \left(\frac{x-1}{x-2} \right) + \tan^{-1} \left(\frac{x+1}{x+2} \right) = \frac{\pi}{4}$ then find the value of x .
- 26) If $2 \tan^{-1} (\cos x) = \tan^{-1} (\operatorname{cosec} x)$ then find the value of x .
- 27) Solve: $\tan^{-1} \left(\frac{1-x}{1+x} \right) = \frac{1}{2} \tan^{-1} x$, for $x > 0$
- 28) If $\sin^{-1}(1-x) - 2 \sin^{-1} x = \frac{\pi}{2}$ then find the value of x .
- 29) If $\tan^{-1} 2x + \tan^{-1} 3x = \frac{\pi}{2}$ then find the value of x .
- 30) Show that $\tan^{-1} \frac{1}{2} - \tan^{-1} \frac{1}{4} = \tan^{-1} \frac{2}{9}$
- 31) Show that $\cot^{-1} \frac{1}{3} - \tan^{-1} \frac{1}{3} = \cot^{-1} \frac{3}{4}$
- 32) Show that $\tan^{-1} \frac{1}{2} = \frac{1}{3} \tan^{-1} \frac{11}{2}$

33) Show that $\cos^{-1} \frac{\sqrt{3}}{2} + 2 \sin^{-1} \frac{\sqrt{3}}{2} = \frac{5\pi}{6}$

34) Show that $2 \cot^{-1} \frac{3}{2} + \sec^{-1} \frac{13}{12} = \frac{\pi}{2}$

35) Prove the following :

(i) $\cos^{-1} x = \tan^{-1} \frac{\sqrt{1-x^2}}{x}$ if $x < 0$. (ii) $\cos^{-1} x = \pi + \tan^{-1} \frac{\sqrt{1-x^2}}{x}$ if $x < 0$.

36) If $|x| < 1$, then prove that $2 \tan^{-1} x = \tan^{-1} \frac{2x}{1+x^2} = \sin^{-1} \frac{2x}{1+x^2} = \cos^{-1} \frac{1-x^2}{1+x^2}$

37) If x, y, z , are positive then prove that $\tan^{-1} \frac{x-y}{1+xy} + \tan^{-1} \frac{y-z}{1+yz} + \tan^{-1} \frac{z-x}{1+zx} = 0$

38) If $\tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \frac{\pi}{2}$ then, show that $xy + yz + zx = 1$

39) If $\cos^{-1} x + \cos^{-1} y + \cos^{-1} z = \pi$ then show that $x^2 + y^2 + z^2 + 2xyz = 1$.



4

Pair of Straight Lines



Let's Study

- 4.1 Combined equation of a pair lines.
- 4.2 Homogeneous equation of degree two.
- 4.3 Angle between lines.
- 4.4 General second degree equation in x and y .

4.1 INTRODUCTION

We know that equation $ax + by + c = 0$, where $a, b, c, \in R$, (a and b not zero simultaneously), represents a line in XY plane. We are familiar with different forms of equations of line. Now let's study two lines simultaneously. For this we need the concept of the combined equation of two lines.



Let's learn.

4.1 Combined equation of a pair of lines :

An equation which represents two lines is called the combined equation of those two lines. Let $u \equiv a_1x + b_1y + c_1$ and $v \equiv a_2x + b_2y + c_2$. Equation $u = 0$ and $v = 0$ represent lines. We know that equation $u + kv = 0$, $k \in R$ represents a family of lines. Let us interpret the equation $uv = 0$.

Theorem 4.1:

The equation $uv = 0$ represents, the combined equation of lines $u = 0$ and $v = 0$

Proof : Consider the lines represented by $u = 0$ and $v = 0$

$$\therefore a_1x + b_1y + c_1 = 0 \text{ and } a_2x + b_2y + c_2 = 0.$$

Let $P(x_p, y_p)$ be a point on the line $u = 0$.

$$\therefore (x_p, y_p) \text{ satisfy the equation } a_1x + b_1y + c_1 = 0$$

$$\therefore a_1x_1 + b_1y_1 + c_1 = 0$$

To show that (x_p, y_p) satisfy the equation $uv = 0$.

$$\begin{aligned} (a_1x_1 + b_1y_1 + c_1)(a_2x_1 + b_2y_1 + c_2) \\ = 0(a_2x_1 + b_2y_1 + c_2) \\ = 0 \end{aligned}$$

Therefore (x_1, y_1) satisfy the equation $uv = 0$.

This proves that every point on the line $u = 0$ satisfy the equation $uv = 0$.

Similarly we can prove that every point on the line $v = 0$ satisfies the equation $uv = 0$.

Now let $R(x', y')$ be any point which satisfy the equation $uv = 0$.

$$\therefore (a_1x' + b_1y' + c_1)(a_2x' + b_2y' + c_2) = 0$$




Figure 4.1

$$\therefore (a_1x' + b_1y' + c_1) = 0 \text{ or } (a_2x' + b_2y' + c_2) = 0$$

Therefore $R(x', y')$ lies on the line $u = 0$ or $v = 0$.

Every points which satisfy the equation $uv = 0$ lies on the line $u = 0$ or the line $v = 0$.

Therefore equation $uv = 0$ represents the combined equation of lines $u = 0$ and $v = 0$.

Remark :

- 1) The combined equation of a pair of lines is also called as the joint equation of a pair of lines.
- 2) Equations $u = 0$ and $v = 0$ are called separate equations of lines represented by $uv = 0$.



Solved Examples :

Ex. 1) Find the combined equation of lines $x + y - 2 = 0$ and $2x - y + 2 = 0$

Solution : The combined equation of lines $u = 0$ and $v = 0$ is $uv = 0$

\therefore The combined equation of lines $x + y - 2 = 0$ and $2x - y + 2 = 0$ is

$$(x + y - 2)(2x - y + 2) = 0$$

$$\therefore x(2x - y + 2) + y(2x - y + 2) - 2(2x - y + 2) = 0$$

$$\therefore 2x^2 - xy + 2x + 2xy - y^2 + 2y - 4x + 2y - 4 = 0$$

$$\therefore 2x^2 + xy - y^2 - 2x + 4y - 4 = 0$$

Ex. 2) Find the combined equation of lines $x - 2 = 0$ and $y + 2 = 0$.

Solution : The combined equation of lines $u = 0$ and $v = 0$ is $uv = 0$.

\therefore The combined equation of lines $x - 2 = 0$ and $y + 2 = 0$ is

$$(x - 2)(y + 2) = 0$$

$$\therefore xy + 2x - 2y - 4 = 0$$

Ex. 3) Find the combined equation of lines $x - 2y = 0$ and $x + y = 0$.

Solution : The combined equation of lines $u = 0$ and $v = 0$ is $uv = 0$.

\therefore The combined equation of lines $x - 2y = 0$ and $x + y = 0$ is

$$(x - 2y)(x + y) = 0$$

$$\therefore x^2 - xy - 2y^2 = 0$$

Ex. 4) Find separate equation of lines represented by $x^2 - y^2 + x + y = 0$.

Solution : We factorize equation $x^2 - y^2 + x + y = 0$ as

$$(x + y)(x - y) + (x + y) = 0$$

$$\therefore (x + y)(x - y + 1) = 0$$

Required separate equations are $x + y = 0$ and $x - y + 1 = 0$.

4.2 Homogeneous equation of degree two:

4.2.1 Degree of a term:

Definition: The sum of the indices of all variables in a term is called the degree of the term.

For example, in the expression $x^2 + 3xy - 2y^2 + 5x + 2$ the degree of the term x^2 is two, the degree of the term $3xy$ is two, the degree of the term $-2y^2$ is two, the degree of $5x$ is one. The degree of constant term 2 is zero. Degree of '0' is not defined.

4.2.2 Homogeneous Equation :

Definition: An equation in which the degree of every term is same, is called a homogeneous equation.

For example: $x^2 + 3xy = 0$, $7xy - 2y^2 = 0$, $5x^2 + 3xy - 2y^2 = 0$ are homogeneous equations.
 But $3x^2 + 2xy + 2y^2 + 5x = 0$ is not a homogeneous equation.

Homogeneous equation of degree two in x and y has form $ax^2 + 2hxy + by^2 = 0$.

Theorem 4.2 :

The combined equation of a pair of lines passing through the origin is a homogeneous equation of degree two in x and y .

Proof : Let $a_1x + b_1y = 0$ and $a_2x + b_2y = 0$ be any two lines passing through the origin.

Their combined equation is $(a_1x + b_1y)(a_2x + b_2y) = 0$

$$a_1a_2x^2 + a_1b_2xy + a_2b_1xy + b_1b_2y^2 = 0$$

$$(a_1a_2)x^2 + (a_1b_2 + a_2b_1)xy + (b_1b_2)y^2 = 0$$

In this if we put $a_1a_2 = a$, $a_1b_2 + a_2b_1 = 2h$, $b_1b_2 = b$, we get, $ax^2 + 2hxy + by^2 = 0$, which is a homogeneous equation of degree two in x and y .




Figure 4.2

Ex.1) Verify that the combined equation of lines $2x + 3y = 0$ and $x - 2y = 0$ is a homogeneous equation of degree two.

Solution :

The combined equation of lines $u = 0$ and $v = 0$ is $uv = 0$.

∴ The combined equation of lines $2x + 3y = 0$ and $x - 2y = 0$ is

$$(2x + 3y)(x - 2y) = 0$$

$$2x^2 - xy - 6y^2 = 0, \text{ which is a homogeneous equation of degree two.}$$

Remark :

The combined equation of a pair of lines passing through the origin is a homogeneous equation of degree two. But every homogeneous equation of degree two *need not* represent a pair of lines.

Equation $x^2 + y^2 = 0$ is a homogeneous equation of degree two but it does not represent a pair of lines.

How to test whether given homogeneous equation of degree two represents a pair of lines or not?

Let's have a theorem.

Theorem 3 : Homogeneous equation of degree two in x and y , $ax^2 + 2hxy + by^2 = 0$ represents a pair of lines passing through the origin if $h^2 - ab \geq 0$.

Proof : Consider the homogeneous equation of degree two in x and y , $ax^2 + 2hxy + by^2 = 0$ (1)

Consider two cases $b = 0$ and $b \neq 0$. These two cases are exhaustive.

Case 1: If $b = 0$ then equation (1) becomes $ax^2 + 2hxy = 0$

∴ $x(ax + 2hy) = 0$, which is the combined equation of lines




Figure 4.3

$$x = 0 \text{ and } ax + 2hy = 0.$$

We observe that these lines pass through the origin.

Case 2: If $b \neq 0$ then we multiply equation (1) by b .

$$abx^2 + 2hbxy + b^2y^2 = 0$$

$$b^2y^2 + 2hbxy = -abx^2$$

To make L.H.S. complete square we add h^2x^2 to both sides.

$$b^2y^2 + 2hbxy + h^2x^2 = h^2x^2 - abx^2$$

$$(by + hx)^2 = (h^2 - ab)x^2$$

$$(by + hx)^2 = \left(\sqrt{h^2 - ab}\right)^2 x^2, \text{ as } h^2 - ab \geq 0$$

$$(by + hx)^2 - \left(\sqrt{h^2 - ab}\right)^2 x^2 = 0$$

$$(by + hx + \sqrt{h^2 - ab} x)(by + hx - \sqrt{h^2 - ab} x) = 0$$

$$[(h + \sqrt{h^2 - ab})x + by] \times [(h - \sqrt{h^2 - ab})x + by] = 0$$

Which is the combined equation of lines $(h + \sqrt{h^2 - ab})x + by = 0$ and $(h - \sqrt{h^2 - ab})x + by = 0$.

As $b \neq 0$, we can write these equations in the form $y = m_1x$ and $y = m_2x$, where $m_1 = \frac{-h - \sqrt{h^2 - ab}}{b}$
and $m_2 = \frac{-h + \sqrt{h^2 - ab}}{b}$.

We observe that these lines pass through the origin.

Therefore equation $abx^2 + 2hbxy + by^2 = 0$ represents a pair of lines passing through the origin if $h^2 - ab \geq 0$.

Remarks:

- 1) If $h^2 - ab > 0$ then line represented by (1) are distinct.
- 2) If $h^2 - ab = 0$ then lines represented by (1) are coincident.
- 3) If $h^2 - ab < 0$ then equation (1) does not represent a pair of lines.
- 4) If $b = 0$ then one of the lines is the Y - axis, whose slope is not defined and the slope of the other line is $-\frac{a}{2h}$ (provided that $h \neq 0$).

- 5) If $h^2 - ab \geq 0$ and $b \neq 0$ then slopes of the lines are $m_1 = \frac{-h - \sqrt{h^2 - ab}}{b}$ and

$$m_2 = \frac{-h + \sqrt{h^2 - ab}}{b}$$

$$\text{Their sum is } m_1 + m_2 = -\frac{2h}{b} \text{ and product is } m_1 m_2 = \frac{a}{b}$$

The quadratic equation in m whose roots are m_1 and m_2 is given by

$$m^2 - (m_1 + m_2)m + m_1 m_2 = 0$$

$$\therefore m^2 - \left(-\frac{2h}{b}\right)m + \frac{a}{b} = 0$$

$$bm^2 + 2hm + a = 0 \quad \dots\dots (2)$$

Equation (2) is called the **auxiliary** equation of equation (1). Roots of equation (2) are slopes of lines represented by equation (1).



Solved Examples

Ex. 1) Show that lines represented by equation $x^2 - 2xy - 3y^2 = 0$ are distinct.

Solution : Comparing equation $x^2 - 2xy - 3y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get

$$a = 1, h = -1 \text{ and } b = -3.$$

$$\begin{aligned} h^2 - ab &= (-1)^2 - (1)(-3) \\ &= 1 + 3 \\ &= 4 > 0 \end{aligned}$$

As $h^2 - ab > 0$, lines represented by equation $x^2 - 2xy - 3y^2 = 0$ are distinct.

Ex. 2) Show that lines represented by equation $x^2 - 6xy + 9y^2 = 0$ are coincident.

Solution : Comparing equation $x^2 - 6xy + 9y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get

$$a = 1, h = -3 \text{ and } b = 9.$$

$$\begin{aligned} h^2 - ab &= (-3)^2 - (1)(9) \\ &= 9 - 9 = 0 \end{aligned}$$

As $h^2 - ab = 0$, lines represented by equation $x^2 - 6xy + 9y^2 = 0$ are coincident.

Ex. 3) Find the sum and the product of slopes of lines represented by $x^2 + 4xy - 7y^2 = 0$.

Solution : Comparing equation $x^2 + 4xy - 7y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get $a = 1, h = 2$ and $b = -7$.

If m_1 and m_2 are slopes of lines represented by this equation then

$$m_1 + m_2 = -\frac{2h}{b} \text{ and } m_1 m_2 = \frac{a}{b}.$$

$$\therefore m_1 + m_2 = \frac{-4}{-7} = \frac{4}{7} \text{ and } m_1 m_2 = \frac{1}{-7} = -\frac{1}{7}$$

$$\text{Their sum is } \frac{4}{7} \text{ and products is } -\frac{1}{7}$$

Ex. 4) Find the separate equations of lines represented by

i) $x^2 - 4y^2 = 0$

ii) $3x^2 - 7xy + 4y^2 = 0$

iii) $x^2 + 2xy - y^2 = 0$

iv) $5x^2 - 3y^2 = 0$

Solution : i) $x^2 - 4y^2 = 0$

$$\therefore (x - 2y)(x + 2y) = 0$$

Required separate equations are

$$x - 2y = 0 \text{ and } x + 2y = 0$$

ii) $3x^2 - 7xy + 4y^2 = 0$

$$\therefore 3x^2 - 3xy - 4xy + 4y^2 = 0$$

$$\therefore 3x(x - y) - 4y(x - y) = 0$$

$$\therefore (x - y)(3x - 4y) = 0$$

Required separate equations are

$$x - y = 0 \text{ and } 3x - 4y = 0$$

$$\text{iii) } x^2 + 2xy - y^2 = 0$$

The corresponding auxiliary equation is $bm^2 + 2hm + a = 0$

$$\therefore -m^2 + 2m + 1 = 0$$

$$m^2 - 2m - 1 = 0$$

$$\therefore m = \frac{2 \pm \sqrt{8}}{2} = \frac{2 \pm 2\sqrt{2}}{2}$$

$$= 1 \pm \sqrt{2}$$

Slopes of these lines are

$$m_1 = 1 + \sqrt{2} \text{ and } m_2 = 1 - \sqrt{2}$$

\therefore Required separate equations are

$$y = m_1 x \text{ and } y = m_2 x$$

$$\therefore y = (1 + \sqrt{2})x \text{ and } y = (1 - \sqrt{2})x$$

$$\therefore (1 + \sqrt{2})x - y = 0 \text{ and } (1 - \sqrt{2})x - y = 0$$

$$\text{iv) } 5x^2 - 3y^2 = 0$$

$$\therefore (\sqrt{5}x)^2 - (\sqrt{3}y)^2 = 0$$

$$\therefore (\sqrt{5}x - \sqrt{3}y)(\sqrt{5}x + \sqrt{3}y) = 0$$

\therefore Required separate equations are

$$\sqrt{5}x - \sqrt{3}y = 0 \text{ and } \sqrt{5}x + \sqrt{3}y = 0$$

Ex. 5 Find the value of k if $2x + y = 0$ is one of the lines represented by $3x^2 + kxy + 2y^2 = 0$.

Solution : Slope of the line $2x + y = 0$ is -2

As $2x + y = 0$ is one of the lines represented by $3x^2 + kxy + 2y^2 = 0$, -2 is a root of the auxiliary equation $2m^2 + km + 3 = 0$

$$\therefore 2(-2)^2 + k(-2) + 3 = 0$$

$$\therefore 8 - 2k + 3 = 0$$

$$\therefore -2k + 11 = 0$$

$$\therefore 2k = 11 \qquad \therefore k = \frac{11}{2}$$

Alternative Method : As $2x + y = 0$ is one of the lines represented by $3x^2 + kxy + 2y^2 = 0$, co-ordinates of every point on the line $2x + y = 0$ satisfy the equation $3x^2 + kxy + 2y^2 = 0$.

As $(1, -2)$ is a point on the line $2x + y = 0$, it must satisfy the combined equation.

$$\therefore 3(1)^2 + k(1)(-2) + 2(-2)^2 = 0$$

$$\therefore -2k + 11 = 0$$

$$\therefore 2k = 11 \qquad \therefore k = \frac{11}{2}$$

Ex. 6) Find the condition that the line $3x - 2y = 0$ coincides with one of the lines represented by $ax^2 + 2hxy + by^2 = 0$.

Solution : The corresponding auxiliary equation is $bm^2 + 2hm + a = 0$.

As line $3x - 2y = 0$ coincides with one of the lines given by $ax^2 + 2hxy + by^2 = 0$, its slope $\frac{3}{2}$ is a root of the auxiliary equation.

$$\therefore \frac{3}{2} \text{ is a root of } bm^2 + 2hm + a = 0$$

$$\therefore b \left(\frac{3}{2}\right)^2 + 2h \left(\frac{3}{2}\right) + a = 0$$

$$\therefore \frac{9}{4}b + 3h + a = 0$$

$$\therefore 4a + 12h + 9b = 0 \text{ is the required condition.}$$

Ex.7) Find the combined equation of the pair of lines passing through the origin and perpendicular to the lines represented by $3x^2 + 2xy - y^2 = 0$.

Solution : Let m_1 and m_2 are slopes of lines represented by $3x^2 + 2xy - y^2 = 0$.

$$\therefore m_1 + m_2 = -\frac{2h}{b} = \frac{-2}{-1} = 2$$

$$\text{and } m_1 m_2 = \frac{a}{b} = \frac{3}{-1} = -3$$

Now required lines are perpendicular to given lines.

$$\therefore \text{Their slopes are } -\frac{1}{m_1} \text{ and } -\frac{1}{m_2}$$

And required lines pass through the origin.

$$\therefore \text{Their equations are } y = -\frac{1}{m_1}x \text{ and } y = -\frac{1}{m_2}x$$

$$\therefore m_1 y = -x \text{ and } m_2 y = -x$$

$$\therefore x + m_1 y = 0 \text{ and } x + m_2 y = 0$$

Their combined equation is $(x + m_1 y)(x + m_2 y) = 0$

$$\therefore x^2 + (m_1 + m_2)xy + m_1 m_2 y^2 = 0$$

$$\therefore x^2 + (2)xy + (-3)y^2 = 0$$

$$\therefore x^2 + 2xy - 3y^2 = 0$$

Ex.8) Find the value of k , if slope of one of the lines represented by $4x^2 + kxy + y^2 = 0$ is four times the slope of the other line.

Solution : Let slopes of the lines represented by $4x^2 + kxy + y^2 = 0$ be m and $4m$,

$$\text{their sum is } m + 4m = 5m$$

$$\text{But their sum is } \frac{-2h}{b} = \frac{-k}{1} = -k$$

$$\therefore 5m = -k$$

$$\therefore m = \frac{-k}{5} \dots (1)$$

Now their product is $(m)(4m) = 4m^2$

But their product is $\frac{a}{b} = \frac{4}{1} = 4$

$$\therefore 4m^2 = 4$$

$$\therefore m^2 = 1 \dots (2)$$

From (1) and (2), we get

$$\left(\frac{-k}{5}\right)^2 = 1$$

$$\therefore k^2 = 25$$

$$\therefore k = \pm 5$$



Exercise 4.1

1) Find the combined equation of the following pairs of lines:

i) $2x + y = 0$ and $3x - y = 0$

ii) $x + 2y - 1 = 0$ and $x - 3y + 2 = 0$

iii) Passing through (2,3) and parallel to the co-ordinate axes.

iv) Passing through (2,3) and perpendicular to lines $3x + 2y - 1 = 0$ and $x - 3y + 2 = 0$

v) Passing through (-1,2), one is parallel to $x + 3y - 1 = 0$ and the other is perpendicular to $2x - 3y - 1 = 0$.

2) Find the separate equations of the lines represented by following equations:

i) $3y^2 + 7xy = 0$

ii) $5x^2 - 9y^2 = 0$

iii) $x^2 - 4xy = 0$

iv) $3x^2 - 10xy - 8y^2 = 0$

v) $3x^2 - 2\sqrt{3}xy - 3y^2 = 0$

vi) $x^2 + 2(\operatorname{cosec} \alpha)xy + y^2 = 0$

vii) $x^2 + 2xy \tan \alpha - y^2 = 0$

3) Find the combined equation of a pair of lines passing through the origin and perpendicular to the lines represented by following equations :

i) $5x^2 - 8xy + 3y^2 = 0$

ii) $5x^2 + 2xy - 3y^2 = 0$

iii) $xy + y^2 = 0$

iv) $3x^2 - 4xy = 0$

4) Find k if,

i) the sum of the slopes of the lines represented by $x^2 + kxy - 3y^2 = 0$ is twice their product.

ii) slopes of lines represent by $3x^2 + kxy - y^2 = 0$ differ by 4.

iii) slope of one of the lines given by $kx^2 + 4xy - y^2 = 0$ exceeds the slope of the other by 8.

- 5) Find the condition that :
- the line $4x + 5y = 0$ coincides with one of the lines given by $ax^2 + 2hxy + by^2 = 0$.
 - the line $3x + y = 0$ may be perpendicular to one of the lines given by $ax^2 + 2hxy + by^2 = 0$.
- 6) If one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is perpendicular to $px + qy = 0$ then show that $ap^2 + 2hpq + by^2 = 0$.
- 7) Find the combined equation of the pair of lines passing through the origin and making an equilateral triangle with the line $y = 3$.
- 8) If slope of one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is four times the other then show that $16h^2 = 25ab$.
- 9) If one of the lines given by $ax^2 + 2hxy + by^2 = 0$ bisects an angle between co-ordinate axes then show that $(a + b)^2 = 4h^2$.

4.3 Angle between lines represented by $ax^2 + 2hxy + by^2 = 0$:

If we know slope of a line then we can find the angles made by the line with the co-ordinate axes. In equation $ax^2 + 2hxy + by^2 = 0$ if $b = 0$ then one of the lines is the Y - axis. Using the slope of the other line we can find the angle between them. In the following discussion we assume that $b \neq 0$, so that slopes of both lines will be defined.

If m_1 and m_2 are slopes of these lines then $m_1 m_2 = \frac{a}{b}$

We know that lines having slopes m_1 and m_2 are perpendicular to each other if and only if $m_1 m_2 = -1$.

$$\therefore \frac{a}{b} = -1$$

$$\therefore a = -b$$

$$\therefore a + b = 0$$

Thus lines represented by $ax^2 + 2hxy + by^2 = 0$ are perpendicular to each other if and only if $a + b = 0$.

If lines are **not perpendicular** to each other then the acute angle between them can be obtained by using the following theorem.

Theorem 4.4 : The acute angle θ between the lines represented by $ax^2 + 2hxy + by^2 = 0$ is given by

$$\tan\theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$

Proof : Let m_1 and m_2 be slopes of lines represented by the equation $ax^2 + 2hxy + by^2 = 0$.

$$\therefore m_1 + m_2 = -\frac{2h}{b} \text{ and } m_1 m_2 = \frac{a}{b}$$

$$\therefore (m_1 - m_2)^2 = (m_1 + m_2)^2 - 4m_1 m_2$$

$$= \left(\frac{2h}{b} \right)^2 - 4 \left(\frac{a}{b} \right)$$

$$= \frac{4h^2}{b^2} - \frac{4ab}{b^2}$$

$$\begin{aligned}
 &= \frac{4h^2 - 4ab}{b^2} \\
 &= \frac{4(h^2 - ab)}{b^2} \\
 \therefore m_1 - m_2 &= \pm \frac{2\sqrt{h^2 - ab}}{b}
 \end{aligned}$$

As θ is the acute angle between the lines,

$$\begin{aligned}
 \tan \theta &= \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| \\
 &= \left| \frac{\pm \frac{2\sqrt{h^2 - ab}}{b}}{1 + \frac{a}{b}} \right| \\
 &= \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|
 \end{aligned}$$




Figure 4.4

Remark : Lines represented by $ax^2 + 2hxy + by^2 = 0$ are coincident if and only if $m_1 = m_2$

$$\begin{aligned}
 \therefore m_1 - m_2 &= 0 \\
 \therefore \frac{2\sqrt{h^2 - ab}}{b} &= 0 \\
 \therefore h^2 - ab &= 0 \\
 \therefore h^2 &= ab
 \end{aligned}$$

Lines represented by $ax^2 + 2hxy + by^2 = 0$ are coincident if and only if $h^2 = ab$.



Solved Examples

Ex.1) Show that lines represented by $3x^2 - 4xy - 3y^2 = 0$ are perpendicular to each other.

Solution : Comparing given equation with $ax^2 + 2hxy + by^2 = 0$ we get $a = 3$, $h = -2$ and $b = -3$.
As $a + b = 3 + (-3) = 0$, lines represented by $3x^2 - 4xy - 3y^2 = 0$ are perpendicular to each other.

Ex. 2) Show that lines represented by $x^2 + 4xy + 4y^2 = 0$ are coincident.

Solution : Comparing given equation with $ax^2 + 2hxy + by^2 = 0$, we get $a = 1$, $h = 2$ and $b = 4$.

$$\begin{aligned}
 \text{As, } h^2 - ab &= (2)^2 - (1)(4) \\
 &= 4 - 4 = 0
 \end{aligned}$$

\therefore Lines represented by $x^2 + 4xy + 4y^2 = 0$ are coincident.

Ex.3) Find the acute angle between lines represented by:

i) $x^2 + xy = 0$

ii) $x^2 - 4xy + y^2 = 0$

iii) $3x^2 + 2xy - y^2 = 0$

iv) $2x^2 - 6xy + y^2 = 0$

v) $xy + y^2 = 0$

Solution :

- i) Comparing equation $x^2 + xy = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get $a = 1$, $h = \frac{1}{2}$ and $b = 0$.
Let θ be the acute angle between them.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right| = \left| \frac{2\sqrt{\frac{1}{4} - 0}}{1} \right| = 1$$

$$\therefore \theta = 45^\circ = \frac{\pi}{4}$$

- ii) Comparing equation $x^2 - 4xy + y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get $a = 1$, $h = -2$ and $b = 1$.
Let θ be the acute angle between them.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right|$$

$$= \left| \frac{2\sqrt{4-1}}{2} \right| = \sqrt{3}$$

$$\therefore \theta = 60^\circ = \frac{\pi}{3}$$

- iii) Comparing equation $3x^2 + 2xy - y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get $a = 3$, $h = 1$ and $b = -1$.
Let θ be the acute angle between them.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right|$$

$$= \left| \frac{2\sqrt{1+3}}{2} \right| = 2.$$

$$\therefore \theta = \tan^{-1}(2)$$

- iv) Comparing equation $2x^2 - 6xy + y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get $a = 2$, $h = -3$ and $b = 1$.
Let θ be the acute angle between them.

$$\therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right|$$

$$= \left| \frac{2\sqrt{9-2}}{3} \right| = \frac{2\sqrt{7}}{3} \quad \therefore \theta = \tan^{-1} \left(\frac{2\sqrt{7}}{3} \right).$$

v) Comparing equation $xy + y^2 = 0$ with $ax^2 + 2hxy + by^2 = 0$, we get $a = 0$, $h = \frac{1}{2}$ and $b = 1$.

Let θ be the acute angle between them.

$$\begin{aligned}\therefore \tan \theta &= \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right| \\ &= \left| \frac{2\sqrt{\frac{1}{4} - 0}}{1} \right| = 1\end{aligned}$$

$$\therefore \theta = 45^\circ = \frac{\pi}{4}.$$

Ex.4) Find the combined equation of lines passing through the origin and making angle $\frac{\pi}{6}$ with the line $3x + y - 6 = 0$.

Solution : Let m be the slope of one of the lines which make angle $\frac{\pi}{6}$ with the line $3x + y - 6 = 0$. Slope of the given line is -3 .

$$\therefore \tan \frac{\pi}{6} = \left| \frac{m - (-3)}{1 + m(-3)} \right|$$

$$\therefore \frac{1}{\sqrt{3}} = \left| \frac{m+3}{1-3m} \right|$$

$$\therefore (1 - 3m)^2 = 3(m + 3)^2$$

$$\therefore 9m^2 - 6m + 1 = 3(m^2 + 6m + 9)$$

$$\therefore 9m^2 - 6m + 1 = 3m^2 + 18m + 27$$

$$\therefore 6m^2 - 24m - 26 = 0$$

$$\therefore 3m^2 - 12m - 13 = 0$$

This is the auxiliary equation of the required combined equation.

The required combined equation is $-13x^2 - 12xy + 3y^2 = 0$

$$\therefore 13x^2 + 12xy - 3y^2 = 0$$

Ex. 5) Find the combined equation of lines passing through the origin and each of which making angle 60° with the X - axis.

Solution :

Let m be the slope of one of the required lines.

The slope of the X - axis is 0. As required lines make angle 60° with the X - axis,

$$\tan 60^\circ = \left| \frac{m - 0}{1 + (m)(0)} \right|$$

$$\therefore \sqrt{3} = |m|$$

$$\therefore m^2 = 3$$

$\therefore m^2 + 0m - 3 = 0$ is the auxiliary equation.

\therefore The required combined equation is

$$-3x^2 + 0xy + y^2 = 0$$

$$\therefore 3x^2 - y^2 = 0$$

Alternative Method: As required lines make angle 60° with the X - axis, their inclination are 60° and 120° . Hence their slopes are $\sqrt{3}$ and $-\sqrt{3}$.

Lines pass through the origin. Their equations are $y = \sqrt{3}x$ and $y = -\sqrt{3}x$

$$\therefore \sqrt{3}x - y = 0 \text{ and } \sqrt{3}x + y = 0$$

Their combined equation is $(\sqrt{3}x - y)(\sqrt{3}x + y) = 0$

$$\therefore 3x^2 - y^2 = 0$$



Exercise 4.2

- Show that lines represented by $3x^2 - 4xy - 3y^2 = 0$ are perpendicular to each other.
- Show that lines represented by $x^2 + 6xy + 9y^2 = 0$ are coincident.
- Find the value of k if lines represented by $kx^2 + 4xy - 4y^2 = 0$ are perpendicular to each other.
- Find the measure of the acute angle between the lines represented by:
 - $3x^2 - 4\sqrt{3}xy + 3y^2 = 0$
 - $4x^2 + 5xy + 4y^2 = 0$
 - $2x^2 + 7xy + 3y^2 = 0$
 - $(a^2 - 3b^2)x^2 + 8abxy + (b^2 - 3a^2)y^2 = 0$
- Find the combined equation of lines passing through the origin each of which making an angle of 30° with the line $3x + 2y - 11 = 0$
- If the angle between lines represented by $ax^2 + 2hxy + by^2 = 0$ is equal to the angle between lines represented by $2x^2 - 5xy + 3y^2 = 0$ then show that $100(h^2 - ab) = (a + b)^2$.
- Find the combined equation of lines passing through the origin and each of which making angle 60° with the Y- axis.

4.4 General Second Degree Equation in x and y:

Equation of the form $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$, where at least one of a, b, h is not zero, is called a general second degree equation in x and y .

Theorem 4.5 : The combined equation of two lines is a general second degree equation in x and y .

Proof: Let $u \equiv a_1x + b_1y + c_1$ and $v \equiv a_2x + b_2y + c_2$. Equations $u = 0$ and $v = 0$ represent lines. Their combined equation is $uv = 0$.

$$\therefore (a_1x + b_1y + c_1)(a_2x + b_2y + c_2) = 0$$

$$a_1a_2x^2 + a_1b_2xy + a_1c_2x + b_1a_2xy + b_1b_2y^2 + b_1c_2y + c_1a_2x + c_1b_2y + c_1c_2 = 0$$

$$\text{Writing } a_1a_2 = a, b_1b_2 = b, a_1b_2 + a_2b_1 = 2h, a_1c_2 + a_2c_1 = 2g, b_1c_2 + b_2c_1 = 2f, c_1c_2 = c,$$

we get, $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$, which is the general equation of degree two in x and y .

Remark : The converse of the above theorem is **not true**. Every general second degree equation in x and y **need not** represent a pair of lines. For example $x^2 + y^2 = 25$ is a general second degree equation in x and y but it does not represent a pair of lines. It represents a circle.

Equation $x^2 + y^2 - 4x + 6y + 13 = 0$ is also a general second degree equation which does not represent a pair of lines. How to identify that whether the given equation represents a pair of lines or not?

4.4.1 The necessary conditions for a general second degree equation.

$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ to represent a pair of lines are:

i) $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$ ii) $h^2 - ab \geq 0$

Remarks :

If equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of lines then

1) These lines are parallel to the line represented by $ax^2 + 2hxy + by^2 = 0$

2) The acute angle between them is given by $\tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right|$

3) Condition for lines to be perpendicular to each other is $a + b = 0$.

4) Condition for lines to be parallel to each other is $h^2 - ab = 0$.

5) Condition for lines to intersect each other is $h^2 - ab > 0$ and the co-ordinates of their point

of intersection are $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$

6) The expression $abc + 2fgh - af^2 - bg^2 - ch^2$ is the expansion of the determinant $\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}$

7) The joint equation of the bisector of the angle between the lines represented by $ax^2 + 2hxy + by^2 = 0$ is $hx^2 - (a-b)xy - hy^2 = 0$. Here coefficient of $x^2 +$ coefficient of $y^2 = 0$.

Hence bisectors are perpendicular to each other



Solved Examples

Ex.1) Show that equation $x^2 - 6xy + 5y^2 + 10x - 14y + 9 = 0$ represents a pair of lines. Find the acute angle between them. Also find the point of their intersection.

Solution: We have $x^2 - 6xy + 5y^2 = (x - 5y)(x - y)$

Suppose $x^2 - 6xy + 5y^2 + 10x - 14y + 9 = (x - 5y + c)(x - y + k)$

$$\therefore x^2 - 6xy + 5y^2 + 10x - 14y + 9 = x^2 - 6xy + 5y^2 + (c + k)x - (c + 5k)y + ck$$

$$\therefore c + k = 10, c + 5k = 14 \text{ and } ck = 9$$

We observe that $c = 9$ and $k = 1$ satisfy all three equations.

$$\therefore \text{Given general equation can be factorized as } (x - 5y + 9)(x - y + 1) = 0$$

\therefore Given equation represents a pair of intersecting lines.

The acute angle between them is given by

$$\tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right| = \left| \frac{2\sqrt{(-3)^2 - (1)(5)}}{1+5} \right| = \frac{2}{3}$$

$$\therefore \theta = \tan^{-1} \left(\frac{2}{3} \right)$$

Their point of intersection is given by

$$\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right) = \left(\frac{21 - 25}{5 - 9}, \frac{-15 + 7}{5 - 9} \right) = (1, 2)$$

Remark :

Note that condition $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$ is not sufficient for equation to represent a pair of lines. We can't use this condition to show that given equation represents a pair of lines.

Ex.2) Find the value of k if the equation $2x^2 + 4xy - 2y^2 + 4x + 8y + k = 0$ represents a pair of lines.

Solution: Comparing given equation with $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ we get

$$a = 2, b = -2, c = k, f = 4, g = 2, h = 2.$$

As given equation represents a pair of lines, it must satisfy the necessary condition.

$$\therefore abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$\therefore (2)(-2)(k) + 2(4)(2)(2) - 2(4)^2 - (-2)(2)^2 - (k)(2)^2 = 0$$

$$\therefore -4k + 32 - 32 + 8 - 4k = 0$$

$$\therefore 8k = 8$$

$$\therefore k = 1.$$

Ex.3) Find p and q if the equation $2x^2 + 4xy - py^2 + 4x + qy + 1 = 0$ represents a pair of perpendicular lines.

Solution: Comparing given equation with $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ we get

$$a = 2, b = -p, c = 1, f = \frac{q}{2}, g = 2, h = 2$$

As lines are perpendicular to each other, $a + b = 0$

$$\therefore 2 + (-p) = 0$$

$$\therefore p = 2$$

As given equation represents a pair of lines, it must satisfy the necessary condition.

$$\therefore abc + 2fgh - af^2 - bg^2 - ch^2 = 0$$

$$\therefore (2)(-p)(1) + 2\left(\frac{q}{2}\right)(2)(2) - 2\left(\frac{q}{2}\right)^2 - (-p)(2)^2 - 1(2)^2 = 0$$

$$\therefore -2p + 4q - \frac{q^2}{2} + 4p - 4 = 0$$

$$\therefore 2p + 4q - \frac{q^2}{2} - 4 = 0 \quad \dots(1)$$

substituting $p = 2$ in (1), we get

$$\therefore 2(2) + 4q - \frac{q^2}{2} - 4 = 0$$

$$\therefore 4q - \frac{q^2}{2} = 0 \therefore 8q - q^2 = 0$$

$$\therefore q(8 - q) = 0$$

$$\therefore q = 0 \text{ or } q = 8$$

Ex.4) Δ OAB is formed by lines $x^2 - 4xy + y^2 = 0$ and the line $x + y - 2 = 0$. Find the equation of the median of the triangle drawn from O.

Solution : Let the co-ordinates of A and B be (x_1, y_1) and (x_2, y_2) respectively.

The midpoint of segment AB is

$$P \left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2} \right)$$

The co-ordinates of A and B can be obtained by solving equations $x + y - 2 = 0$ and $x^2 - 4xy + y^2 = 0$ simultaneously.

put $y = 2 - x$ in $x^2 - 4xy + y^2 = 0$.

$$x^2 - 4x(2 - x) + (2 - x)^2 = 0$$

$$\therefore 6x^2 - 12x + 4 = 0$$

$$\therefore 3x^2 - 6x + 2 = 0$$

x_1 and x_2 are roots of this equation.

$$x_1 + x_2 = -\frac{-6}{3} = 2$$

$$\therefore \frac{x_1+x_2}{2} = 1$$

The x co-ordinate of P is 1.

As P lies on the line $x + y - 2 = 0$

$$\therefore 1 + y - 2 = 0 \quad \therefore y = 1$$

\therefore Co-ordinates of P are (1,1).

The equation of the median OP is $\frac{y-0}{1-0} = \frac{x-0}{1-0}$

$$\therefore y = x \quad \therefore x - y = 0.$$




Figure 4.5



Exercise 4.3

- Find the joint equation of the pair of lines:
 - Through the point $(2, -1)$ and parallel to lines represented by $2x^2 + 3xy - 9y^2 = 0$
 - Through the point $(2, -3)$ and parallel to lines represented by $x^2 + xy - y^2 = 0$
- Show that equation $x^2 + 2xy + 2y^2 + 2x + 2y + 1 = 0$ does not represent a pair of lines.
- Show that equation $2x^2 - xy - 3y^2 - 6x + 19y - 20 = 0$ represents a pair of lines.
- Show the equation $2x^2 + xy - y^2 + x + 4y - 3 = 0$ represents a pair of lines. Also find the acute angle between them.
- Find the separate equation of the lines represented by the following equations :
 - $(x - 2)^2 - 3(x - 2)(y + 1) + 2(y + 1)^2 = 0$
 - $10(x + 1)^2 + (x + 1)(y - 2) - 3(y - 2)^2 = 0$
- Find the value of k if the following equations represent a pair of lines :
 - $3x^2 + 10xy + 3y^2 + 16y + k = 0$
 - $kxy + 10x + 6y + 4 = 0$
 - $x^2 + 3xy + 2y^2 + x - y + k = 0$

- 7) Find p and q if the equation $px^2 - 8xy + 3y^2 + 14x + 2y + q = 0$ represents a pair of perpendicular lines.
- 8) Find p and q if the equation $2x^2 + 8xy + py^2 + qx + 2y - 15 = 0$ represents a pair of parallel lines.
- 9) Equations of pairs of opposite sides of a parallelogram are $x^2 - 7x + 6 = 0$ and $y^2 - 14y + 40 = 0$. Find the joint equation of its diagonals.
- 10) ΔOAB is formed by lines $x^2 - 4xy + y^2 = 0$ and the line $2x + 3y - 1 = 0$. Find the equation of the median of the triangle drawn from O .
- 11) Find the co-ordinates of the points of intersection of the lines represented by $x^2 - y^2 - 2x + 1 = 0$.



Let's remember!

- An equation which represents two lines is called the combined equation of those two lines.
- The equation $uv = 0$ represents the combined equation of lines $u = 0$ and $v = 0$.
- The sum of the indices of all variables in a term is called the degree of the term.
- An equation in which the degree of every term is same, is called a homogeneous equation.
- The combined equation of a pair of lines passing through the origin is a homogeneous equation of degree two in x and y .
- Every homogeneous equation of degree two need not represent a pair of lines.
- A homogeneous equation of degree two in x and y , $ax^2 + 2hxy + by^2 = 0$ represents a pair of lines passing through the origin if $h^2 - ab \geq 0$.
- If $h^2 - ab > 0$ then lines are distinct.
- If $h^2 - ab = 0$ then lines are coincident.
- Slopes of these lines are $m_1 = \frac{-h - \sqrt{h^2 - ab}}{b}$ and $m_2 = \frac{-h + \sqrt{h^2 - ab}}{b}$
- Their sum is, $m_1 + m_2 = -\frac{2h}{b}$ and product is, $m_1 m_2 = \frac{a}{b}$
- The quadratic equation in m whose roots are m_1 and m_2 is given by $bm^2 + 2hm + a = 0$, called the **auxiliary** equation.
- Lines represented by $ax^2 + 2hxy + by^2 = 0$ are perpendicular to each other if and only if $a + b = 0$.
- Lines represented by $ax^2 + 2hxy + by^2 = 0$ are coincident if and only if $h^2 - ab = 0$.
- The acute angle θ between the lines represented by $ax^2 + 2hxy + by^2 = 0$ is given by

$$\tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$
- Equation of the form $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ is called a general second degree equation in x and y .
- The combined equation of two lines is a general second degree equation in x and y .
- The necessary conditions for a general second degree equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ to represent a pair of lines are:
 - i) $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$
 - ii) $h^2 - ab \geq 0$

- The expression $abc + 2fgh - af^2 - bg^2 - ch^2$ is the expansion of the determinant $\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}$
- If equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of lines then
 - These lines are parallel to the lines represented by $ax^2 + 2hxy + by^2 = 0$.
 - The acute angle between them is given by $\tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a+b} \right|$
 - Condition for lines to be perpendicular to each other is $a + b = 0$.
 - Condition for lines to be parallel to each other is $h^2 - ab = 0$.
 - Condition for lines to intersect each other is $h^2 - ab \geq 0$ and the co-ordinates of their point of intersection are $\left(\frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right)$

MISCELLANEOUS EXERCISE 4

I : Choose correct alternatives.

- If the equation $4x^2 + hxy + y^2 = 0$ represents two coincident lines, then $h =$ _____.

A) ± 2	B) ± 3
C) ± 4	D) ± 5
- If the lines represented by $kx^2 - 3xy + 6y^2 = 0$ are perpendicular to each other then _____.

A) $k = 6$	B) $k = -6$
C) $k = 3$	D) $k = -3$
- Auxiliary equation of $2x^2 + 3xy - 9y^2 = 0$ is _____.

A) $2m^2 + 3m - 9 = 0$	B) $9m^2 - 3m - 2 = 0$
C) $2m^2 - 3m + 9 = 0$	D) $-9m^2 - 3m + 2 = 0$
- The difference between the slopes of the lines represented by $3x^2 - 4xy + y^2 = 0$ is _____.

A) 2	B) 1
C) 3	D) 4
- If the two lines $ax^2 + 2hxy + by^2 = 0$ make angles α and β with X-axis, then $\tan(\alpha + \beta) =$ _____.

A) $\frac{h}{a+b}$	B) $\frac{h}{a-b}$
C) $\frac{2h}{a+b}$	D) $\frac{2h}{a-b}$
- If the slope of one of the two lines $\frac{x^2}{a} + \frac{2xy}{h} + \frac{y^2}{b} = 0$ is twice that of the other, then $ab:h^2 =$ _____.

A) 1 : 2	B) 2 : 1
----------	----------

- C) 8 : 9
D) 9 : 8
- 7) The joint equation of the lines through the origin and perpendicular to the pair of lines $3x^2 + 4xy - 5y^2 = 0$ is _____.
- A) $5x^2 + 4xy - 3y^2 = 0$
B) $3x^2 + 4xy - 5y^2 = 0$
C) $3x^2 - 4xy + 5y^2 = 0$
D) $5x^2 + 4xy + 3y^2 = 0$
- 8) If acute angle between lines $ax^2 + 2hxy + by^2 = 0$ is, $\frac{\pi}{4}$ then $4h^2 =$ _____.
- A) $a^2 + 4ab + b^2$
B) $a^2 + 6ab + b^2$
C) $(a + 2b)(a + 3b)$
D) $(a - 2b)(2a + b)$
- 9) If the equation $3x^2 - 8xy + 4y^2 + 2x + 14y + p = 1$ represents a pair of perpendicular lines then the values of p and q are respectively _____.
- A) -3 and -7
B) -7 and -3
C) 3 and 7
D) -7 and 3
- 10) The area of triangle formed by the lines $x^2 + 4xy + y^2 = 0$ and $x - y - 4 = 0$ is _____.
- A) $\frac{4}{\sqrt{3}}$ Sq. units
B) $\frac{8}{\sqrt{3}}$ Sq. units
C) $\frac{16}{\sqrt{3}}$ Sq. units
D) $\frac{15}{\sqrt{3}}$ Sq. units
- 11) The combined equation of the co-ordinate axes is _____.
- A) $x + y = 0$
B) $xy = k$
C) $xy = 0$
D) $x - y = k$
- 12) If $h^2 = ab$, then slope of lines $ax^2 + 2hxy + by^2 = 0$ are in the ratio _____.
- A) 1 : 2
B) 2 : 1
C) 2 : 3
D) 1 : 1
- 13) If slope of one of the lines $ax^2 + 2hxy + by^2 = 0$ is 5 times the slope of the other, then $5h^2 =$ _____.
- A) ab
B) $2 ab$
C) $7 ab$
D) $9 ab$
- 14) If distance between lines $(x - 2y)^2 + k(x - 2y) = 0$ is 3 units, then $k =$
- A) ± 3
B) $\pm 5\sqrt{5}$
C) 0
D) $\pm 3\sqrt{5}$

II. Solve the following.

1) Find the joint equation of lines:

i) $x - y = 0$ and $x + y = 0$

ii) $x + y - 3 = 0$ and $2x + y - 1 = 0$

iii) Passing through the origin and having slopes 2 and 3.

iv) Passing through the origin and having inclinations 60° and 120° .

v) Passing through $(1,2)$ and parallel to the co-ordinate axes.

vi) Passing through $(3,2)$ and parallel to the line $x = 2$ and $y = 3$.

- vii) Passing through $(-1,2)$ and perpendicular to the lines $x + 2y + 3 = 0$ and $3x - 4y - 5 = 0$.
- viii) Passing through the origin and having slopes $1 + \sqrt{3}$ and $1 - \sqrt{3}$
- ix) Which are at a distance of 9 units from the Y - axis.
- x) Passing through the point $(3,2)$, one of which is parallel to the line $x - 2y = 2$ and other is perpendicular to the line $y = 3$.
- xi) Passing through the origin and perpendicular to the lines $x + 2y = 19$ and $3x + y = 18$.
- 2) Show that each of the following equation represents a pair of lines.
- $x^2 + 2xy - y^2 = 0$
 - $4x^2 + 4xy + y^2 = 0$
 - $x^2 - y^2 = 0$
 - $x^2 + 7xy - 2y^2 = 0$
 - $x^2 - 2\sqrt{3}xy - y^2 = 0$
- 3) Find the separate equations of lines represented by the following equations:
- $6x^2 - 5xy - 6y^2 = 0$
 - $x^2 - 4y^2 = 0$
 - $3x^2 - y^2 = 0$
 - $2x^2 + 2xy - y^2 = 0$
- 4) Find the joint equation of the pair of lines through the origin and perpendicular to the lines given by :
- $x^2 + 4xy - 5y^2 = 0$
 - $2x^2 - 3xy - 9y^2 = 0$
 - $x^2 + xy - y^2 = 0$
- 5) Find k if
- The sum of the slopes of the lines given by $3x^2 + kxy - y^2 = 0$ is zero.
 - The sum of slopes of the lines given by $x^2 + kxy - 3y^2 = 0$ is equal to their product.
 - The slope of one of the lines given by $3x^2 - 4xy + 5y^2 = 0$ is 1.
 - One of the lines given by $3x^2 - kxy + 5y^2 = 0$ is perpendicular to the $5x + 3y = 0$.
 - The slope of one of the lines given by $3x^2 + 4xy + ky^2 = 0$ is three times the other.
 - The slopes of lines given by $kx^2 + 5xy + y^2 = 0$ differ by 1.
 - One of the lines given by $6x^2 + kxy + y^2 = 0$ is $2x + y = 0$.
- 6) Find the joint equation of the pair of lines which bisect angle between the lines given by $x^2 + 3xy + 2y^2 = 0$
- 7) Find the joint equation of the pair of lines through the origin and making equilateral triangle with the line $x = 3$.
- 8) Show that the lines $x^2 - 4xy + y^2 = 0$ and $x + y = 10$ contain the sides of an equilateral triangle. Find the area of the triangle.
- 9) If the slope of one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ is three times the other then prove that $3h^2 = 4ab$.

- 10) Find the combined equation of the bisectors of the angles between the lines represented by $5x^2 + 6xy - y^2 = 0$.
- 11) Find a if the sum of slope of lines represented by $ax^2 + 8xy + 5y^2 = 0$ is twice their product.
- 12) If line $4x - 5y = 0$ coincides with one of the lines given by $ax^2 + 2hxy + by^2 = 0$ then show that $25a + 40h + 16b = 0$.
- 13) Show that the following equations represent a pair of lines, find the acute angle between them.
- $9x^2 - 6xy + y^2 + 18x - 6y + 8 = 0$
 - $2x^2 + xy - y^2 + x - 4y + 3 = 0$
 - $(x - 3)^2 + (x - 3)(y - 4) - 2(y - 4)^2 = 0$
- 14) Find the combined equation of pair of lines through the origin each of which makes angle of 60° with the Y-axis.
- 15) If lines represented by $ax^2 + 2hxy + by^2 = 0$ make angles of equal measures with the co-ordinate axes then show that $a = \pm b$.
- 16) Show that the combined equation of a pair of lines through the origin and each making an angle of α with the line $x + y = 0$ is $x^2 + 2(\sec 2\alpha)xy + y^2 = 0$.
- 17) Show that the line $3x + 4y + 5 = 0$ and the lines $(3x + 4y)^2 - 3(4x - 3y)^2 = 0$ form an equilateral triangle.
- 18) Show that lines $x^2 - 4xy + y^2 = 0$ and $x + y = \sqrt{6}$ form an equilateral triangle. Find its area and perimeter.
- 19) If the slope of one of the lines given by $ax^2 + 2hxy + by^2 = 0$ is square of the other then show that $a^2b + ab^2 + 8h^3 = 6abh$.
- 20) Prove that the product of lengths of perpendiculars drawn from P (x_1, y_1) to the lines represented by $ax^2 + 2hxy + by^2 = 0$ is $\left| \frac{ax_1^2 + 2hx_1y_1 + by_1^2}{\sqrt{(a-b)^2 + 4h^2}} \right|$
- 21) Show that the difference between the slopes of lines given by $(\tan^2\theta + \cos^2\theta)x^2 - 2xy \tan \theta + (\sin^2 \theta)y^2 = 0$ is two.
- 22) Find the condition that the equation $ay^2 + bxy + ex + dy = 0$ may represent a pair of lines.
- 23) If the lines given by $ax^2 + 2hxy + by^2 = 0$ form an equilateral triangle with the line $lx + my = 1$ then show that $(3a + b)(a + 3b) = 4h^2$.
- 24) If line $x + 2 = 0$ coincides with one of the lines represented by the equation $x^2 + 2xy + 4y + k = 0$ then show that $k = -4$.
- 25) Prove that the combined equation of the pair of lines passing through the origin and perpendicular to the lines represented by $ax^2 + 2hxy + by^2 = 0$ is $bx^2 - 2hxy + ay^2 = 0$
- 26) If equation $ax^2 - y^2 + 2y + c = 1$ represents a pair of perpendicular lines then find a and c .



5

Vectors



Let's Study

1. Vectors and their types.
2. Section formula.
3. Dot Product of Vectors.
4. Cross Product of Vectors.
5. Triple Product of Vectors.



Let's recall.

Scalar quantity : A quantity which can be completely described by magnitude only is called a scalar quantity. e.g. mass, length, temperature, area, volume, time distance, speed, work, money, voltage, density, resistance etc. In this book, scalars are given by real numbers.

Vector quantity : A quantity which needs to be described using both magnitude and direction is called a vector quantity. e.g. displacement, velocity, force, electric field, acceleration, momentum etc.



Let's learn.

5.1 Representation of Vector :

Vector is represented by a directed line segment.

If AB is a segment and its direction is shown with an arrowhead as in figure, then the directed segment AB has magnitude as well as direction. This is an example of vector.

The segment AB with direction from A to B denotes the vector \overline{AB} read as \overline{AB} read as 'AB bar' while direction from B to A denotes the vector \overline{BA} .

In vector \overline{AB} , the point A is called the initial point and the point B is called the terminal point

The directed line segment is a part of a line of unlimited length which is called the line of support or the line of action of the given vector.

If the initial and terminal point are not specified then the vectors are denoted by \vec{a} , \vec{b} , \vec{c} or **a**, **b**, **c** (bold face) etc.




Fig. 5.1

5.1.1 Magnitude of a Vector :

The magnitude (or size or length) of \overline{AB} is denoted by $|\overline{AB}|$ and is defined as the length of segment AB. *i.e.* $|\overline{AB}| = l(AB)$

Magnitudes of vectors $\vec{a}, \vec{b}, \vec{c}$ are $|\vec{a}|, |\vec{b}|, |\vec{c}|$ respectively.

The magnitude of a vector does not depend on its direction. Since the length is never negative, $|\vec{a}| \geq 0$

5.1.2 Types of Vectors :

i) **Zero Vector :** A vector whose initial and terminal points coincide, is called a zero vector (or null vector) and denoted as $\vec{0}$. Zero vector cannot be assigned a definite direction and it has zero magnitude or it may be regarded as having any suitable direction. The vectors $\overline{AA}, \overline{BB}$ represent the zero vector and $|\overline{AA}| = 0$.

ii) **Unit Vector :** A vector whose magnitude is unity (*i.e.* 1 unity) is called a unit vector. The unit vector in the direction of a given vector \vec{a} is denoted by \hat{a} , read as 'a-cap' or 'a-hat'.




Fig. 5.2

iii) **Co-initial and Co-terminus Vectors :** Vectors having same initial point are called co-initial vectors, whereas vectors having same terminal point are called co-terminus vectors. Here \vec{a} and \vec{b} are co-initial vectors. \vec{c} and \vec{d} are co-terminus vectors.

iv) **Equal Vectors :** Two or more vectors are said to be equal vectors if they have same magnitude and direction.

- As $|\vec{a}| = |\vec{b}|$, and their directions are same regardless of initial point, we write as $\vec{a} = \vec{b}$.
- Here $|\vec{a}| = |\vec{c}|$, but directions are not same, so $\vec{a} \neq \vec{c}$.
- Here directions of \vec{a} and \vec{d} same but $|\vec{a}| \neq |\vec{d}|$, so $\vec{a} \neq \vec{d}$.




Fig 5.3

v) **Negative of a Vector :** If \vec{a} is a given vector, then the negative of \vec{a} is vector whose magnitude is same as that of \vec{a} but whose direction is opposite to that of \vec{a} . It is denoted by $-\vec{a}$.

Thus, if $\overline{PQ} = \vec{a}$, then $\overline{QP} = -\vec{a} = -\overline{PQ}$.

Here $|\overline{PQ}| = |-\overline{QP}|$.




Fig. 5.4

vi) **Collinear Vectors :** Vectors are said to be collinear vectors if they are parallel to same line or they are along the same line.

vii) **Free Vectors :** If a vector can be translated anywhere in the space without changing its magnitude and direction then such a vector is called free vector. In other words, the initial point of free vector can be taken anywhere in the space keeping magnitude and direction same.




Fig. 5.5

viii) **Localised Vectors :** For a vector of given magnitude and direction, if it's initial point is fixed in space, then such a vector is called localised vector.

For example, if there are two stationary cars A and B on the road and a force is applied to car A, it is a localised vector and only car A moves, while car B is not affected.



Fig 5.6

Note that in this chapter vectors are treated as free vectors unless otherwise stated.

Activity 1 :

Write the following vectors in terms of vectors \vec{p} , \vec{q} and \vec{r} .

- i) $\vec{AB} = \square$ ii) $\vec{BA} = \square$ iii) $\vec{BC} = \square$
- iv) $\vec{CB} = \square$ v) $\vec{CA} = \square$ vi) $\vec{AC} = \square$




Fig 5.7

Algebra of Vectors :

5.1.3 Scalar Multiplication :

$2\vec{a}$ has the same direction as \vec{a} but is twice as long as \vec{a} .

Let \vec{a} be any vector and k be a scalar, then vector $k\vec{a}$, the scalar multiple is defined a vector whose magnitude is $|k\vec{a}| = |k| |\vec{a}|$ and vectors \vec{a} and $k\vec{a}$ have the same direction if $k > 0$ and opposite direction if $k < 0$.




Fig 5.8

Note :

- i) If $k = 0$, then $k\vec{a} = \vec{0}$.
- ii) \vec{a} and $k\vec{a}$ are collinear or parallel vectors.
- iii) Two non zero vectors \vec{a} and \vec{b} are collinear or parallel if $\vec{a} = m\vec{b}$, where $m \neq 0$.
- iv) Let \hat{a} be the unit vector along non-zero




Fig 5.9

vector \vec{a} then $\vec{a} = |\vec{a}| \hat{a}$ or $\frac{\vec{a}}{|\vec{a}|} = \hat{a}$.

- v) A vector of length k in the same direction as \vec{a} is $k\hat{a} = k \left(\frac{\vec{a}}{|\vec{a}|} \right)$.




Fig 5.10

Now, consider a boat in a river going from one bank of the river to the other in a direction perpendicular to the flow of the river. Then, it is acted upon by two velocity vectors, one is the velocity imparted to the boat by its engine and other one is the velocity of the flow of river water. Under the simultaneous influence of these two velocities, the boat starts travelling with a different velocity. To have a precise velocity (i.e. resultant velocity) of the boat we use the law of addition of vectors.

5.1.4 Addition of Two Vectors : If \vec{a} and \vec{b} are any two vectors then their addition (or resultant) is denoted by $\vec{a} + \vec{b}$.

There are two laws of addition of two vectors.

Parallelogram Law : Let \vec{a} and \vec{b} be two vectors. Consider \overline{AB} and \overline{AD} along two adjacent sides of a parallelogram, such that $\overline{AB} = \vec{a}$ and $\overline{AD} = \vec{b}$ then $\vec{a} + \vec{b}$ lies along the diagonal of a parallelogram with \vec{a} and \vec{b} as sides.




Fig 5.11

$$\overline{AC} = \overline{AB} + \overline{AD} \text{ i.e. } \vec{c} = \vec{a} + \vec{b}.$$

Triangle Law of addition of two vectors : Let \vec{a} , \vec{b} be any two vectors then consider triangle ABC as shown in figure such that $\overline{AB} = \vec{a}$ and $\overline{BC} = \vec{b}$ then $\vec{a} + \vec{b}$ is given by vector \overline{AC} along the third side of triangle ABC.

$$\text{Thus, } \vec{a} + \vec{b} = \overline{AB} + \overline{BC} = \overline{AC}.$$

This is known as the triangle law of addition of two vectors \vec{a} and \vec{b} . The triangle law can also be applied to the ΔADC .

$$\text{Here, } \overline{AD} = \overline{BC} = \vec{b}, \overline{DC} = \overline{AB} = \vec{a}$$

$$\text{Hence, } \overline{AD} + \overline{DC} = \vec{b} + \vec{a} = \overline{AC}$$

$$\text{Thus, } \overline{AC} = \vec{b} + \vec{a} = \vec{a} + \vec{b}$$




Fig 5.12

5.1.5 Subtraction of two vectors : If \vec{a} and \vec{b} are two vectors, then $\vec{a} - \vec{b} = \vec{a} + (-\vec{b})$, where $-\vec{b}$ is the negative vector of vector \vec{b} . Let $\overline{AB} = \vec{a}$, $\overline{BC} = \vec{b}$, now construct a vector \overline{BD} such that its magnitude is same as the vector \overline{BC} , but the direction is opposite to that of it.

$$\text{i.e. } \overline{BD} = -\overline{BC}.$$

$$\therefore \overline{BD} = -\vec{b}.$$

Thus applying triangle law of addition.

We have

$$\begin{aligned} \therefore \overline{AD} &= \overline{AB} + \overline{BD} \\ &= \overline{AB} - \overline{BC} \\ &= \vec{a} - \vec{b} \end{aligned}$$




Fig 5.13

Note :

- i) If (velocity) vectors \vec{a} and \vec{b} are acting simultaneously then we use parallelogram law of addition.
- ii) If (velocity) vectors \vec{a} and \vec{b} are acting one after another then we use triangle law of addition.
- iii) Adding vector to its opposite vector gives $\vec{0}$

$$\text{As } \overline{PQ} + \overline{QP} = \overline{PP} = \vec{0} \text{ or}$$

$$\text{As } \overline{PQ} = -\overline{QP}, \text{ then } \overline{PQ} + \overline{QP} = -\overline{QP} + \overline{QP} = \vec{0}.$$



Fig 5.14

iv) In ΔABC , $\overline{AC} = -\overline{CA}$, so $\overline{AB} + \overline{BC} + \overline{CA} = \overline{AA} = \vec{0}$. This means that when the vectors along the sides of a triangle are in order, their resultant is zero as initial and terminal points become same.

v) The addition law of vectors can be extended to a polygon :

Let \vec{a} , \vec{b} , \vec{c} and \vec{d} be four vectors. Let $\overline{PQ} = \vec{a}$, $\overline{QR} = \vec{b}$, $\overline{RS} = \vec{c}$ and $\overline{ST} = \vec{d}$.

$$\therefore \vec{a} + \vec{b} + \vec{c} + \vec{d}$$

$$= \overline{PQ} + \overline{QR} + \overline{RS} + \overline{ST}$$

$$= (\overline{PQ} + \overline{QR}) + \overline{RS} + \overline{ST}$$

$$= (\overline{PR} + \overline{RS}) + \overline{ST}$$

$$= \overline{PS} + \overline{ST}$$

$$= \overline{PT}$$




Fig 5.15

Thus, the vector \overline{PT} represents sum of all vectors \vec{a} , \vec{b} , \vec{c} and \vec{d} .

This is also called as extended law of addition of vectors or polygonal law of addition of vectors.

- vi) $\vec{a} + \vec{b} = \vec{b} + \vec{a}$ (commutative)
- vii) $(\vec{a} + \vec{b}) + \vec{c} = \vec{a} + (\vec{b} + \vec{c})$ (associative)
- viii) $\vec{a} + \vec{0} = \vec{a}$ ($\vec{0}$ is additive identify)
- ix) $\vec{a} + (-\vec{a}) = \vec{0}$ ($-\vec{a}$ is additive inverse)
- x) If \vec{a} and \vec{b} are vectors and m and n are scalars, then
 - i) $(m + n) \vec{a} = m \vec{a} + n \vec{a}$ (distributive)
 - ii) $m(\vec{a} + \vec{b}) = m \vec{a} + m \vec{b}$ (distributive)
 - iii) $m(n \vec{a}) = (mn) \vec{a} = n(m \vec{a})$
- xi) $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$, this is known as "Triangle Inequality".

This is obtained from the triangle law, as the length of any side of triangle is less than the sum of the other two sides. *i.e.* in triangle ABC, $AC < AB + BC$,

where, $AC = |\vec{a} + \vec{b}|$, $AB = |\vec{a}|$, $BC = |\vec{b}|$

- xii) Any two vectors \vec{a} and \vec{b} determine a plane and vectors $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ lie in the same plane.

Activity 2 :

In quadrilateral PQRS, find a resultant vector

- i) $\overline{QR} + \overline{RS} = \square$
- ii) $\overline{PQ} + \overline{QR} = \square$
- iii) $\overline{PS} + \overline{SR} + \overline{RQ} = \square$
- iv) $\overline{PR} + \overline{RQ} + \overline{QS} = \square$
- v) $\overline{QR} - \overline{SR} - \overline{PS} = \square$
- vi) $\overline{QP} - \overline{RP} + \overline{RS} = \square$




Fig 5.16

Theorem 1:

Two non-zero vectors \vec{a} and \vec{b} are collinear if and only if there exist scalars m and n , at least one of them is non-zero such that $m \vec{a} + n \vec{b} = \vec{0}$.

Proof : Only If - part :

Suppose \vec{a} and \vec{b} are collinear.

\therefore There exists a scalar $t \neq 0$ such that $\vec{a} = t \vec{b}$

$$\therefore \bar{a} - t\bar{b} = \bar{0}$$

i.e. $m\bar{a} + n\bar{b} = \bar{0}$, where $m = 1$ and $n = -t$.

If - part :

Conversely, suppose $m\bar{a} + n\bar{b} = \bar{0}$ and $m \neq 0$.

$$\therefore m\bar{a} = -n\bar{b}$$

$$\therefore \bar{a} = \left(-\frac{n}{m}\right)\bar{b}, \text{ where } t = \left(-\frac{n}{m}\right) \text{ is a scalar.}$$

$$\text{i.e. } \bar{a} = t\bar{b},$$

$\therefore \bar{a}$ is scalar multiple of \bar{b} .

$\therefore \bar{a}$ and \bar{b} are collinear.

Corollary 1 : If two vectors \bar{a} and \bar{b} are not collinear and $m\bar{a} + n\bar{b} = \bar{0}$, then $m = 0, n = 0$. (This can be proved by contradiction assuming $m \neq 0$ or $n \neq 0$).

Corollary 2 : If two vectors \bar{a} and \bar{b} are not collinear and $m\bar{a} + n\bar{b} = p\bar{a} + q\bar{b}$, then $m = p, n = q$.

For example, If two vectors \bar{a} and \bar{b} are not collinear and $3\bar{a} + y\bar{b} = x\bar{a} + 5\bar{b}$, then $3 = x, y = 5$.

5.1.6 Coplanar Vectors :

Two or more vectors are coplanar, if they lie in the same plane or in parallel plane.

Vector \bar{a} and \bar{b} are coplanar.

Vector \bar{a} and \bar{c} are coplanar.

Are vectors \bar{a} and \bar{e} coplanar ?




Fig 5.17

Remark : Any two intersecting straight lines OA and OB in space determine a plane. We may choose for convenience the coordinate axes of the plane so that O is origin and axis OX is along one of OA or OB.

Theorem 2 : Let \bar{a} and \bar{b} be non-collinear vectors. A vector \bar{r} is coplanar with \bar{a} and \bar{b} if and only if there exist unique scalars t_1, t_2 , such that $\bar{r} = t_1\bar{a} + t_2\bar{b}$.

Proof : Only If-part :

Suppose \bar{r} is coplanar with \bar{a} and \bar{b} . To show that there exist unique scalars t_1 and t_2 such that $\bar{r} = t_1\bar{a} + t_2\bar{b}$.

Let \bar{a} be along OA and \bar{b} be along OB. Given a vector \bar{r} , with initial point O, Let $\overline{OP} = \bar{r}$, draw lines parallel to OB, meeting OA in M and parallel to OA, meeting OB in N.




Fig 5.18

Then $ON = t_2\bar{b}$ and $OM = t_1\bar{a}$ for some $t_1, t_2 \in \mathbb{R}$. By triangle law or parallelogram law, we have $\bar{r} = t_1\bar{a} + t_2\bar{b}$

If Part : Suppose $\bar{r} = t_1\bar{a} + t_2\bar{b}$, and we have to show that \bar{r}, \bar{a} and \bar{b} are co-planar.

As \bar{a}, \bar{b} are coplanar, $t_1\bar{a}, t_2\bar{b}$ are also coplanar. Therefore $t_1\bar{a} + t_2\bar{b}, \bar{a}, \bar{b}$ are coplanar.

Therefore $\bar{a}, \bar{b}, \bar{r}$ are coplanar.

Uniqueness :

Suppose vector $\vec{r} = t_1 \vec{a} + t_2 \vec{b}$... (1)

can also be written as $\vec{r} = s_1 \vec{a} + s_2 \vec{b}$... (2)

Subtracting (2) from (1) we get,

$$\vec{0} = (t_1 - s_1) \vec{a} + (t_2 - s_2) \vec{b}$$

But, \vec{a} and \vec{b} are non-collinear, vectors

By Corollary 1 of Theorem 1,

$$\therefore t_1 - s_1 = 0 = t_2 - s_2$$

$$\therefore t_1 = s_1 \text{ and } t_2 = s_2.$$

Therefore, the uniqueness follows.




Fig 5.19

Remark :

Linear combination of vectors : If $\vec{a}_1, \vec{a}_2, \vec{a}_3, \dots, \vec{a}_n$ are n vectors and $m_1, m_2, m_3, \dots, m_n$ are n scalars, then the vector $m_1 \vec{a}_1 + m_2 \vec{a}_2 + m_3 \vec{a}_3 + \dots + m_n \vec{a}_n$ is called a linear combination of vectors $\vec{a}_1, \vec{a}_2, \vec{a}_3, \dots, \vec{a}_n$. If atleast one m_i is not zero then the linear combination is non zero linear combination

For example : Let \vec{a}, \vec{b} be vectors and m, n are scalars then the vector $\vec{c} = m\vec{a} + n\vec{b}$ is called a linear combination of vector \vec{a} and \vec{b} . Vectors \vec{a}, \vec{b} and \vec{c} are coplanar vectors.

Theorem 3 : Three vectors \vec{a}, \vec{b} and \vec{c} are coplanar, if and only if there exists a non-zero linear combination $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$ with $(x, y, z) \neq (0, 0, 0)$.

Proof : Only If - part :

Assume that \vec{a}, \vec{b} and \vec{c} are coplanar.

Case - 1 : Suppose that any two of \vec{a}, \vec{b} and \vec{c} are collinear vectors, say \vec{a} and \vec{b} .

\therefore There exist scalars x, y at least one of which is non-zero such that $x\vec{a} + y\vec{b} = \vec{0}$
i.e. $x\vec{a} + y\vec{b} + 0\vec{c} = \vec{0}$ and $(x, y, 0)$ is the required solution for $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$.

Case - 2 : No two vectors \vec{a}, \vec{b} and \vec{c} are collinear.

As \vec{c} is coplanar with \vec{a} and \vec{b} ,

\therefore we have scalars x, y such that $\vec{c} = x\vec{a} + y\vec{b}$ (using Theorem 2).

$\therefore x\vec{a} + y\vec{b} - \vec{c} = \vec{0}$ and $(x, y, -1)$ is the required solution for $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$.

If - part : Conversely, suppose $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$ where one of x, y, z is non-zero, say $z \neq 0$.

$$\therefore \vec{c} = \frac{-x}{z} \vec{a} - \frac{y}{z} \vec{b}$$

$\therefore \vec{c}$ is coplanar with \vec{a} and \vec{b} .

$\therefore \vec{a}, \vec{b}$ and \vec{c} are coplanar vectors.

Corollary 1 : If three vectors \vec{a}, \vec{b} and \vec{c} are not coplanar and $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$, then $x = 0, y = 0$ and $z = 0$ because if $(x, y, z) \neq (0, 0, 0)$ then $\vec{a}, \vec{b}, \vec{c}$ are coplanar.

Corollary 2 : The vectors \vec{a}, \vec{b} and $x\vec{a} + y\vec{b}$ are coplanar for all values of x and y .

5.1.7 Vector in Two Dimensions (2-D) :

The plane spanned (covered) by non collinear vectors \vec{a} and \vec{b} is $\{x\vec{a} + y\vec{b} \mid x, y \in \mathbb{R}\}$, where \vec{a} and \vec{b} have same initial point.

This is 2-D space where generators are \vec{a} and \vec{b} or its basis is $\{\vec{a}, \vec{b}\}$

For example, in XY plane, let $M = (1, 0)$ and $N = (0, 1)$ be two points along X and Y axis respectively.

Then, we define unit vectors \hat{i} and \hat{j} as $\overline{OM} = \hat{i}$, $\overline{ON} = \hat{j}$.

Given any other vector say \overline{OP} , where $P = (3, 4)$ then

$$\overline{OP} = 3\hat{i} + 4\hat{j}$$




Fig 5.20

5.1.8 Three Dimensional (3-D) Coordinate System :

Any point in the plane is represented as an ordered pair (a, b) where a and b are distances (with suitable sign) of point (a, b) from Y-axis and X-axis respectively.

To locate a point in space, three numbers are required. Here, we need three coordinate axes OX, OY and OZ and to determine a point we need distances of it from three planes formed by these axes.

We represent any point in space by an ordered triple (a, b, c) of real numbers.




Fig 5.21

O is the origin and three directed lines through O that are perpendicular to each other are the coordinate axes.

Label them as X-axis (XOX'), Y-axis (YOY') and Z-axis (ZOZ'). The direction of Z-axis is determined by right hand rule *i.e.* When you hold your right hand so that the fingers curl from the positive X-axis toward the positive Y-axis, your thumb points along the positive Z-axis, as shown in figure.

Co-ordinates of a point in space :

Let P be a point in the space. Draw perpendiculars PL, PM, PN through P to XY-plane, YZ-plane and XZ-plane respectively, where points L, M and N are feet of perpendiculars in XY, YZ and XZ planes respectively.




Fig 5.22

For point $P(x, y, z)$, x , y and z are x -coordinate, y -coordinate and z -coordinate respectively.

Point of intersection of all 3 planes is origin $O(0, 0, 0)$.




Fig 5.23

Co-ordinates of points on co-ordinate axes :

Points on X-axis, Y-axis and Z-axis have coordinates given by $A(x, 0, 0)$, $B(0, y, 0)$ and $C(0, 0, z)$.

Co-ordinates of points on co-ordinate planes :

Points in XY-plane, YZ-plane and ZX-plane are given by $L(x, y, 0)$, $M(0, y, z)$, $N(x, 0, z)$ respectively.

Distance of P(x, y, z) from co-ordinate planes :

- i) Distance of P from XY plane = |PL| = |z|.
- ii) Distance of P from YZ plane = |PM| = |x|.
- iii) Distance of P from XZ plane = |PN| = |y|.

Distance of any point from origin :

Distance of P (x, y, z) from the origin O(0, 0, 0) from figure 5.23 we have,

$$\begin{aligned}
 l(OP) &= \sqrt{OL^2 + LP^2} \quad (\Delta OLP \text{ right angled triangle}) \\
 &= \sqrt{OA^2 + AL^2 + LP^2} \\
 &= \sqrt{OA^2 + OB^2 + OC^2} \\
 &= \sqrt{x^2 + y^2 + z^2}
 \end{aligned}$$

Distance between any two points in space

Distance between two points A(x₁, y₁, z₁) and B(x₂, y₂, z₂) in space is given by distance formula

$$l(AB) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$$

Distance of a point P(x, y, z) from coordinate axes.

In fig. 5.23, PA is perpendicular to X-axis. Hence distance of P from X-axis is PA.

$$\begin{aligned}
 \therefore PA &= \sqrt{(x - x)^2 + (y - 0)^2 + (z - 0)^2} \\
 &= \sqrt{y^2 + z^2}
 \end{aligned}$$

- iv) In a right-handed system. Octants II, III and IV are found by rotating anti-clockwise around the positive Z-axis. Octant V is vertically below Octant I. Octants VI, VII and VIII are then found by rotating anti-clockwise around the negative Z-axis.

Signs of coordinates of a point P(x, y, z) in different octants :

Octant (x, y, z)	(I) O-XYZ (+, +, +)	(II) O-X'YZ (-, +, +)	(III) O-XY'Z (+, -, +)	(IV) O-X'Y'Z (-, -, +)
Octant (x, y, z)	(V) O-XYZ' (+, +, -)	(VI) O-X'YZ' (-, +, -)	(VII) O-XY'Z' (+, -, -)	(VIII) O-X'Y'Z' (-, -, -)

Point in Octants




Fig 5.24

Various shapes in b space




Fig 5.25

5.1.9 Components of Vector :

In order to be more precise about the direction of a vector we can represent a vector as a linear combination of basis vectors.

Take the points A(1, 0, 0), B(0, 1, 0) and C(0, 0, 1) on the X-axis, Y-axis and Z-axis, respectively.

Then $|\overline{OA}| = |\overline{OB}| = |\overline{OC}| = 1$

The vectors \overline{OA} , \overline{OB} , \overline{OC} each having magnitude 1 are called unit vectors along the axes X, Y, Z respectively. These vectors are denoted by \hat{i} , \hat{j} , \hat{k} respectively and also called as standard basis vectors or standard unit vectors.

Any vector, along X-axis is a scalar multiple of unit vector \hat{i} , along Y-axis is a scalar multiple of \hat{j} and along Z-axis is a scalar multiple of \hat{k} . (Collinearity property).

e.g.i) $3\hat{i}$ is a vector along OX with magnitude 3.

ii) $5\hat{j}$ is a vector along OY with magnitude 5.

iii) $4\hat{k}$ is a vector along OZ with magnitude 4.




Fig 5.26

Theorem 4 :

If \vec{a} , \vec{b} , \vec{c} are three non-coplanar vectors, then any vector \vec{r} in the space can be uniquely expressed as a linear combination of \vec{a} , \vec{b} , \vec{c} .

Proof : Let A be any point in the space, take the vectors \vec{a} , \vec{b} , \vec{c} and \vec{r} , so that A becomes their initial point (Fig.5.27).

Let $\overline{AP} = \vec{r}$. As \vec{a} , \vec{b} , \vec{c} are non-coplanar vectors, they determine three distinct planes intersecting at the point A. Through the point P, draw the plane parallel to the plane formed by vectors \vec{b} , \vec{c} .

This plane intersects line containing \vec{a} at point B. Similarly, draw the other planes and complete the parallelepiped.




Fig 5.27

Now, \overline{AB} and \overline{a} are collinear.

$\therefore \overline{AB} = x\overline{a}$, where x is a scalar.

Similarly, we have $\overline{AQ} = y\overline{b}$ and $\overline{AS} = z\overline{c}$, where y and z are scalars.

Also, by triangle law of addition of vectors,

$$\begin{aligned} \overline{AP} &= \overline{AC} + \overline{CP} \quad (\text{In } \triangle ACP) \\ &= (\overline{AB} + \overline{BC}) + \overline{CP} \quad (\text{In } \triangle ABC) \\ &= \overline{AB} + \overline{BC} + \overline{CP} \\ \overline{AP} &= \overline{AB} + \overline{BC} + \overline{CP} \end{aligned}$$

$$\therefore \overline{r} = x\overline{a} + y\overline{b} + z\overline{c}. \quad (\because \overline{AQ} = \overline{BC} \text{ and } \overline{AS} = \overline{CP})$$

Therefore, any vector \overline{r} in the space can be expressed as a linear combination of \overline{a} , \overline{b} and \overline{c} .

Uniqueness :

Suppose $\overline{r} = x_1\overline{a} + x_2\overline{b} + x_3\overline{c}$ and also $\overline{r} = y_1\overline{a} + y_2\overline{b} + y_3\overline{c}$ for some scalars x_1, x_2, x_3 and y_1, y_2, y_3 .

We need to prove $x_1 = y_1, x_2 = y_2$ and $x_3 = y_3$.

Subtracting one expression from the other we have $(x_1 - y_1)\overline{a} + (x_2 - y_2)\overline{b} + (x_3 - y_3)\overline{c} = \overline{0}$.

By Corollary 1 of Theorem 3

As $\overline{a}, \overline{b}, \overline{c}$ are non-coplanar we must have $x_1 - y_1 = x_2 - y_2 = x_3 - y_3 = 0$ that is $x_1 = y_1, x_2 = y_2, x_3 = y_3$, as desired.

5.1.10 Position vector of a point P(x, y, z) in space :

Consider a point P in space, having coordinates (x, y, z) with respect to the origin O(0, 0, 0). Then the vector \overline{OP} having O and P as its initial and terminal points, respectively is called the position vector of the point P with respect to O.

Let P (x, y, z) be a point in space.

$$\therefore OA = PM = x, OB = PN = y, OC = PL = z.$$

i.e. A \equiv (x, 0, 0), B \equiv (0, y, 0) and C \equiv (0, 0, z).

Let $\hat{i}, \hat{j}, \hat{k}$ be unit vectors along positive directions of X-axis, Y-axis and Z-axis is respectively.

$$\therefore \overline{OA} = x\hat{i}, \overline{OB} = y\hat{j} \text{ and } \overline{OC} = z\hat{k}$$

\overline{OP} is the position vector of point P in space with respect to origin O.

Representation of \overline{OP} in terms of unit vector $\hat{i}, \hat{j}, \hat{k}$.

In $\triangle OLP$ we have (see fig. 5.28)

$$\begin{aligned} \overline{OP} &= \overline{OL} + \overline{LP} \\ &= \overline{OA} + \overline{AL} + \overline{LP} && (\text{from } \triangle OAL) \\ &= \overline{OA} + \overline{OB} + \overline{OC} && (\because \overline{AL} = \overline{OB}) \\ &= x\hat{i} + y\hat{j} + z\hat{k} && \dots \dots \dots (1) \end{aligned}$$




Fig 5.28

Magnitude of \overline{OP}

Now, $OP^2 = OL^2 + LP^2$ (In right angled ΔOLP)
 $= OA^2 + AL^2 + LP^2$ (In right angled ΔOAL)
 $= OA^2 + OB^2 + OC^2$
 $= x^2 + y^2 + z^2.$ ($\because AL = OB$)

$\therefore l(OP) = \sqrt{x^2 + y^2 + z^2}$

$\therefore (\overline{OP}) = \sqrt{x^2 + y^2 + z^2}$

5.1.11 Component form of \vec{r} :

If \vec{r} is a position vector (p.v.) of point P w.r.t. O then $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

In this representation x, y, z are called the components of \vec{r} along OX, OY and OZ.

Note that any vector in space is unique linear combination of \hat{i}, \hat{j} and \hat{k} .

Note : Some authors represent vector $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ using angle brackets as $\vec{r} = \langle x, y, z \rangle$ for the point (x, y, z) .

5.1.12 Vector joining two points :

Let O be the origin then \overline{OA} and \overline{OB} are the position vectors of points A and B w.r.t. origin 'O'.

In ΔAOB , we have by triangle law.

$$\begin{aligned} \overline{AB} &= \overline{AO} + \overline{OB} \\ &= -\overline{OA} + \overline{OB} \quad (\because \overline{AO} = -\overline{OA}) \\ &= \overline{OB} - \overline{OA} \\ &= \text{position vector of B} - \text{position vector of A} \end{aligned}$$

That is, $\overline{AB} = \vec{b} - \vec{a}$

If $A \equiv (x_1, y_1, z_1)$ and $B \equiv (x_2, y_2, z_2)$

Then $\overline{OA} = x_1\hat{i} + y_1\hat{j} + z_1\hat{k}$ and $\overline{OB} = x_2\hat{i} + y_2\hat{j} + z_2\hat{k}$

$$\begin{aligned} \therefore \overline{AB} &= \overline{OB} - \overline{OA} \\ &= (x_2\hat{i} + y_2\hat{j} + z_2\hat{k}) - (x_1\hat{i} + y_1\hat{j} + z_1\hat{k}) \\ &= (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k} \end{aligned}$$

$$|\overline{AB}| = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$




Fig 5.29

In general, any non zero vector \vec{a} in space can be expressed uniquely as the linear combination of $\hat{i}, \hat{j}, \hat{k}$ as $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ where a_1, a_2, a_3 are scalars.

$\therefore |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$

Note :

(i) If $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ and $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ then $\vec{a} = \vec{b}$ if $a_1 = b_1, a_2 = b_2, a_3 = b_3$.

(ii) If $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ and $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ then

$$\vec{a} + \vec{b} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) + (b_1\hat{i} + b_2\hat{j} + b_3\hat{k}) = (a_1 + b_1)\hat{i} + (a_2 + b_2)\hat{j} + (a_3 + b_3)\hat{k}.$$

(iii) If k is any scalar then $k\vec{a} = k(a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) = ka_1\hat{i} + ka_2\hat{j} + ka_3\hat{k}$

Also if \vec{b} and \vec{a} are collinear i.e. $\vec{b} = k\vec{a}$ then $\frac{b_1}{a_1} = \frac{b_2}{a_2} = \frac{b_3}{a_3} = k$.

(iv) Let \hat{a} be a unit vector along a non zero vector \vec{a} in space, Then $\hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{a_1\hat{i} + a_2\hat{j} + a_3\hat{k}}{\sqrt{a_1^2 + a_2^2 + a_3^2}}$.

(v) **(Collinearity of 3-points)** Three distinct points A, B and C with position vectors \vec{a} , \vec{b} and \vec{c} respectively are collinear if and only if there exist three non-zero scalars x, y and z such that $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$ and $x + y + z = 0$. (Use Theorem 1 and the fact that \overline{AB} and \overline{BC} are collinear).

(vi) **(Coplanarity of 4-points)** Four distinct points A, B, C and D (no three of which are collinear) with position vectors \vec{a} , \vec{b} , \vec{c} and \vec{d} respectively are coplanar if and only if there exist four scalars x, y, z and w , not all zero, such that $x\vec{a} + y\vec{b} + z\vec{c} + w\vec{d} = \vec{0}$,

where $x + y + z + w = 0$. (Use Theorem 2 and the fact that \overline{AB} , \overline{AC} and \overline{AD} are coplanar).

(vii) **Linearly dependent vectors :** A set of non-zero vectors \vec{a} , \vec{b} and \vec{c} is said to be linearly dependent if there exist scalars x, y, z not all zero such that $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$.

Such vectors \vec{a} , \vec{b} and \vec{c} are coplanar.

(viii) **Linearly independent vectors :** A set of non-zero vectors \vec{a} , \vec{b} and \vec{c} is said to be linearly independent if $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$, then $x = y = z = 0$.

Such vectors \vec{a} , \vec{b} and \vec{c} are non-coplanar.



Solved Examples

Ex.1. State the vectors which are :

- (i) equal in magnitude
- (ii) parallel
- (iii) in the same direction
- (iv) equal
- (v) negatives of one another




Fig 5.30

Solution :

- (i) \vec{a} , \vec{c} and \vec{e} ; \vec{b} and \vec{d}
- (ii) \vec{a} , \vec{b} , \vec{c} and \vec{d}
- (iii) \vec{a} and \vec{b} ; \vec{c} and \vec{d}
- (iv) none are equal
- (v) \vec{a} and \vec{c} , \vec{b} and \vec{d}

Ex. 2. In the diagram $\vec{KL} = \vec{a}$, $\vec{LN} = \vec{b}$, $\vec{NM} = \vec{c}$ and $\vec{KT} = \vec{d}$. Find in terms of \vec{a} , \vec{b} , \vec{c} and \vec{d} . (i) \vec{LT} (ii) \vec{KM} (iii) \vec{TN} (iv) \vec{MT}




Fig 5.31

Solution :

- (i) In ΔPTQ , using triangle law
 $\vec{KL} + \vec{LT} = \vec{KT}$ i.e. $\vec{a} + \vec{LT} = \vec{d}$
 $\vec{LT} = \vec{d} - \vec{a}$
- (ii) Using polygonal law of addition of vectors for polygon KLMN
 $\vec{KM} = \vec{KL} + \vec{LN} + \vec{NM}$.
 $= \vec{a} + \vec{b} + \vec{c}$
- (iii) Using polygonal law of addition of vectors for polygon TKLN
 $\vec{TN} = \vec{TK} + \vec{KL} + \vec{LN}$
 $= -\vec{d} + \vec{a} + \vec{b} = \vec{a} + \vec{b} - \vec{d}$
- (iv) Using polygonal law of addition of vectors for polygon TKLNM
 $\vec{MT} = \vec{MN} + \vec{NL} + \vec{LK} + \vec{KT}$
 $= -\vec{c} - \vec{b} - \vec{a} + \vec{d}$
 $= \vec{d} - \vec{a} - \vec{b} - \vec{c}$.

Ex.3. Find the magnitude of following vectors :

- (i) $\vec{a} = \hat{i} - 2\hat{j} + 4\hat{k}$ (ii) $\vec{b} = 4\hat{i} - 3\hat{j} - 7\hat{k}$
- (iii) a vector with initial point : (1, -3, 4); terminal point : (1, 0, -1).

Solution :

- (i) $|\vec{a}| = \sqrt{1^2 + (-2)^2 + 4^2} = \sqrt{21}$
- (ii) $|\vec{b}| = \sqrt{4^2 + (-3)^2 + (-7)^2}$
 $|\vec{b}| = \sqrt{16 + 9 + 49} = \sqrt{74}$

$$(iii) |\vec{c}| = (\hat{i} - \hat{k}) - (\hat{i} - 3\hat{j} + 4\hat{k}) = 3\hat{j} - 5\hat{k}$$

$$|\vec{c}| = \sqrt{0 + 3^2 + (-5)^2} = \sqrt{34}$$

Ex. 4. A(2, 3), B(-1, 5), C(-1, 1) and D(-7, 5) are four points in the Cartesian plane.

- (i) Find \overline{AB} and \overline{CD} .
 (ii) Check if, \overline{CD} is parallel to \overline{AB} .
 (iii) E is the point (k, 1) and \overline{AC} is parallel to \overline{BE} . Find k.

Solution : (i) $\vec{a} = 2\hat{i} + 3\hat{j}, \vec{b} = -\hat{i} + 5\hat{j}, \vec{c} = -\hat{i} + \hat{j}, \vec{d} = -7\hat{i} +$

$$\overline{AB} = \vec{b} - \vec{a} = (-\hat{i} + 5\hat{j}) - (2\hat{i} + 3\hat{j}) = -3\hat{i} + 2\hat{j}$$

$$\overline{CD} = \vec{d} - \vec{c} = (-7\hat{i} + 5\hat{j}) - (-\hat{i} + \hat{j}) = -6\hat{i} + 4\hat{j}$$

(ii) $\overline{CD} = -6\hat{i} + 4\hat{j} = 2(-3\hat{i} + 2\hat{j}) = 2 \overline{AB}$ therefore \overline{CD} and \overline{AB} are parallel.

(iii) $\overline{BE} = (k\hat{i} + \hat{j}) - (-\hat{i} + 5\hat{j}) = (k+1)\hat{i} - 4\hat{j}$

$$\overline{AC} = (-\hat{i} + \hat{j}) - (2\hat{i} + 3\hat{j}) = -3\hat{i} - 2\hat{j}$$

$$\overline{BE} = m\overline{AC}$$

$$(k+1)\hat{i} - 4\hat{j} = m(-3\hat{i} - 2\hat{j})$$

So -4 therefore $2 = m$

and $k+1 = -3m$

$$k+1 = -3(2)$$

$$k = -6 - 1$$

$$k = -7$$

Ex. 5. Determine the values of c that satisfy $|c\vec{u}| = 3, \vec{u} = \hat{i} + 2\hat{j} + 3\hat{k}$

Solution : $|c\vec{u}| = \sqrt{c^2 + 4c^2 + 9c^2} = |c|\sqrt{14} = 3$

$$\therefore c = \pm \frac{3}{\sqrt{14}}$$

Ex. 6. Find a unit vector (i) in the direction of \vec{u} and (ii) in the direction opposite of \vec{u} . where $\vec{u} = 8\hat{i} + 3\hat{j} - \hat{k}$.

Solution : (i) $\hat{u} = \frac{\vec{u}}{|\vec{u}|} = \frac{8\hat{i} + 3\hat{j} - \hat{k}}{\sqrt{74}}$

$$= \frac{1}{\sqrt{74}}(8\hat{i} + 3\hat{j} - \hat{k}) \text{ is the unit vector in direction of } \vec{u}.$$

(ii) $-\hat{u} = -\frac{1}{\sqrt{74}}(8\hat{i} + 3\hat{j} - \hat{k})$ is the unit vector in opposite direction of \vec{u} .

Ex. 7. Show that the vectors $2\hat{i} - 3\hat{j} + 4\hat{k}$ and $-4\hat{i} + 6\hat{j} - 8\hat{k}$ are parallel.

Solution : $\vec{a} = 2\hat{i} - 3\hat{j} + 4\hat{k}$

$$\vec{b} = -4\hat{i} + 6\hat{j} - 8\hat{k} = -2(2\hat{i} - 3\hat{j} + 4\hat{k}) = -2\vec{a}$$

As \vec{b} is scalar multiple of \vec{a}

$\therefore \vec{b}$ and \vec{a} are parallel.

Ex. 8. The non-zero vectors \vec{a} and \vec{b} are not collinear find the value of λ and μ :

(i) $\vec{a} + 3\vec{b} = 2\lambda\vec{a} - \mu\vec{b}$

(ii) $(1 + \lambda)\vec{a} + 2\lambda\vec{b} = \mu\vec{a} + 4\mu\vec{b}$

(iii) $(3\lambda + 5)\vec{a} + \vec{b} = 2\mu\vec{a} + (\lambda - 3)\vec{b}$

Solution : (i) $2\lambda = 1, 3 = -\mu \therefore \lambda = \frac{1}{2}, \mu = -3$

(ii) $1 + \lambda = \mu, 2\lambda = 4\mu, \lambda = 2\mu$

$$1 + 2\mu = \mu, 1 = -\mu$$

$$\therefore \mu = -1, \lambda = -2$$

(iii) $3\lambda + 5 = 2\mu, 1 = \lambda - 3, \therefore \lambda = 1 + 3 = 4$

$$\text{and } 3(4) + 5 = 2\mu$$

$$\therefore 2\mu = 17. \text{ So } \mu = \frac{17}{2}$$

Ex. 9. Are the following set of vectors linearly independent?

(i) $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}, \vec{b} = 3\hat{i} - 6\hat{j} + 9\hat{k}$

(ii) $\vec{a} = -2\hat{i} - 4\hat{k}, \vec{b} = \hat{i} - 2\hat{j} - \hat{k}, \vec{c} = \hat{i} - 4\hat{j} + 3\hat{k}$. Interpret the results.

Solution : (i) $\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}, \vec{b} = 3\hat{i} - 6\hat{j} + 9\hat{k}$

$$\vec{b} = 3(\hat{i} - 2\hat{j} + 3\hat{k})$$

$\vec{b} = 3\vec{a}$ Here \vec{a} and \vec{b} linearly dependent vectors. Hence \vec{a} and \vec{b} are collinear.

(ii) $\vec{a} = -2\hat{i} - 4\hat{k}, \vec{b} = \hat{i} - 2\hat{j} - \hat{k}, \vec{c} = \hat{i} - 4\hat{j} + 3\hat{k}$

$$\text{Let } x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$$

$$\therefore x(-2\hat{i} - 4\hat{k}) + y(\hat{i} - 2\hat{j} - \hat{k}) + z(\hat{i} - 4\hat{j} + 3\hat{k}) = \vec{0}$$

$$-2x + y + z = 0$$

$$-2y - 4z = 0$$

$$-4x - y + 3z = 0$$

$$\therefore x = 0, y = 0, z = 0. \text{ Here } \vec{a}, \vec{b} \text{ and } \vec{c} \text{ are linearly independent vectors.}$$

Hence \vec{a}, \vec{b} and \vec{c} are non-coplanar.

Ex. 10. If $\vec{a} = 4\hat{i} + 3\hat{k}$ and $\vec{b} = -2\hat{i} + \hat{j} + 5\hat{k}$ find (i) $|\vec{a}|$, (ii) $\vec{a} + \vec{b}$, (iii) $\vec{a} - \vec{b}$, (iv) $3\vec{b}$, (v) $2\vec{a} + 5\vec{b}$

Solution : (i) $|\vec{a}| = \sqrt{4^2 + 0^2 + 3^2} = \sqrt{25} = 5$

(ii) $\vec{a} + \vec{b} = (4\hat{i} + 3\hat{k}) + (-2\hat{i} + \hat{j} + 5\hat{k}) = 2\hat{i} + \hat{j} + 8\hat{k}$

(iii) $\vec{a} - \vec{b} = (4\hat{i} + 3\hat{k}) - (-2\hat{i} + \hat{j} + 5\hat{k}) = 6\hat{i} - \hat{j} - 2\hat{k}$

(iv) $3\vec{b} = 3(-2\hat{i} + \hat{j} + 5\hat{k}) = -6\hat{i} + 3\hat{j} + 15\hat{k}$

(v) $2\vec{a} + 5\vec{b} = 2(4\hat{i} + 3\hat{k}) + 5(-2\hat{i} + \hat{j} + 5\hat{k})$
 $= (8\hat{i} + 6\hat{k}) + (-10\hat{i} + 5\hat{j} + 25\hat{k}) = -2\hat{i} + 5\hat{j} + 31\hat{k}$

Ex. 11. What is the distance from the point (2, 3, 4) to (i) the XY plane? (ii) the X-axis? (iii) origin (iv) point (-2, 7, 3).

Solution :

(a) The distance from (2, 3, 4) to the XY plane is $|z| = 4$ units.

(b) The distance from (2, 3, 4) to the X-axis is $\sqrt{y^2 + z^2} = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$ units.

(c) The distance from $(x, y, z) \equiv (2, 3, 4)$ to origin (0, 0, 0) is

$$\sqrt{x^2 + y^2 + z^2} = \sqrt{2^2 + 3^2 + 4^2} = \sqrt{4 + 9 + 16} = \sqrt{29} \text{ units.}$$

(d) The distance from (2, 3, 4) to (-2, 7, 3) is

$$\sqrt{(2+2)^2 + (3-7)^2 + (4-3)^2} = \sqrt{16+16+9} = \sqrt{41} \text{ units.}$$

Ex. 12. Prove that the line segment joining the midpoints of two sides of a triangle is parallel to and half of the third side.

Solution : Let the triangle be ABC. If M and N are the midpoints of AB and AC respectively, then

$$\vec{AM} = \frac{1}{2} \vec{AB} \text{ and } \vec{AN} = \frac{1}{2} \vec{AC}. \text{ Thus by triangle law}$$

$$\vec{AN} = \vec{AM} + \vec{MN}$$

$$\therefore \vec{MN} = \vec{AN} - \vec{AM} = \frac{\vec{AC} - \vec{AB}}{2} = \frac{\vec{BC}}{2}$$

Thus, \vec{MN} is parallel to and half as long as \vec{BC} .




Fig 5.32

Ex. 13. In quadrilateral ABCD, M and N are the mid-points of the diagonals AC and BD respectively.

Prove that $\vec{AB} + \vec{AD} + \vec{CB} + \vec{CD} = 4\vec{MN}$

Solution : $\vec{AB} = \vec{AM} + \vec{MN} + \vec{NB}$... (1)

$\vec{AD} = \vec{AM} + \vec{MN} + \vec{ND}$... (2)

$\vec{CB} = \vec{CM} + \vec{MN} + \vec{NB}$... (3)

$\vec{CD} = \vec{CM} + \vec{MN} + \vec{ND}$... (4)




Fig 5.33

Add (1), (2), (3), (4) to get

$$\begin{aligned} \overline{AB} + \overline{AD} + \overline{CB} + \overline{CD} &= 2\overline{AM} + 2\overline{CM} + 4\overline{MN} + 2\overline{NB} + 2\overline{ND} \\ &= 2(\overline{AM} + \overline{CM}) + 4\overline{MN} + 2(\overline{NB} + \overline{ND}) \\ &= 2(\overline{AM} - \overline{AM}) + 4\overline{MN} + 2(\overline{NB} - \overline{NB}) \quad (\because \overline{MC} = \overline{AM} \text{ and } \overline{DN} = \overline{NB}) \\ &= 2 \times (\overline{0}) + 4\overline{MN} + 2 \times (\overline{0}) \\ &= 4\overline{MN} \end{aligned}$$

Ex. 14. Express $-\hat{i} - 3\hat{j} + 4\hat{k}$ as the linear combination of the vectors $2\hat{i} + \hat{j} - 4\hat{k}$, $2\hat{i} - \hat{j} + 3\hat{k}$ and $3\hat{i} + \hat{j} - 2\hat{k}$.

Solution : Let $\vec{r} = -\hat{i} - 3\hat{j} + 4\hat{k}$, $\vec{a} = 2\hat{i} + \hat{j} - 4\hat{k}$, $\vec{b} = 2\hat{i} - \hat{j} + 3\hat{k}$, $\vec{c} = 3\hat{i} + \hat{j} - 2\hat{k}$

Consider $\vec{r} = x\vec{a} + y\vec{b} + z\vec{c}$

$$-\hat{i} - 3\hat{j} + 4\hat{k} = x(2\hat{i} + \hat{j} - 4\hat{k}) + y(2\hat{i} - \hat{j} + 3\hat{k}) + z(3\hat{i} + \hat{j} - 2\hat{k})$$

$$-\hat{i} - 3\hat{j} + 4\hat{k} = (2x + 2y + 3z)\hat{i} + (x - y + z)\hat{j} + (-4x + 3y - 2z)\hat{k}$$

By equality of vectors, we get $-1 = 2x + 2y + 3z$, $-3 = x - y + z$, $4 = -4x + 3y - 2z$,

Using Cramer's rule we get, $x = 2$, $y = 2$, $z = 3$. Therefore $\vec{r} = 2\vec{a} + 2\vec{b} - 3\vec{c}$

Ex. 15. Show that the three points A(1, -2, 3), B(2, 3, -4) and C(0, -7, 10) are collinear.

Solution : If \vec{a} , \vec{b} and \vec{c} are the position vectors of the points A, B and C respectively, then

$$\vec{a} = \hat{i} - 2\hat{j} + 3\hat{k}$$

$$\vec{b} = 2\hat{i} + 3\hat{j} - 4\hat{k}$$

$$\vec{c} = 0\hat{i} - 7\hat{j} + 10\hat{k}$$

$$\overline{AB} = \vec{b} - \vec{a} = \hat{i} + 5\hat{j} - 7\hat{k} \quad \dots(1) \quad \text{and}$$

$$\overline{AC} = \vec{c} - \vec{a}$$

$$= -\hat{i} - 5\hat{j} + 7\hat{k}$$

$$= (-1)[\hat{i} + 5\hat{j} - 7\hat{k}]$$

$$\overline{AC} = (-1)\overline{AB} \quad \dots \text{from}(1)$$

That is, \overline{AC} is a scalar multiple of \overline{AB} . Therefore, they are parallel. But point A is in common. Hence, the points A, B and C are collinear.

Ex. 16. Show that the vectors $4\hat{i} + 13\hat{j} - 18\hat{k}$, $\hat{i} - 2\hat{j} + 3\hat{k}$ and $2\hat{i} + 3\hat{j} - 4\hat{k}$ are coplanar.

Solution : Let, $\vec{a} = 4\hat{i} + 13\hat{j} - 18\hat{k}$, $\vec{b} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\vec{c} = 2\hat{i} + 3\hat{j} - 4\hat{k}$

Consider $\vec{a} = m\vec{b} + n\vec{c}$

$$\therefore 4\hat{i} + 13\hat{j} - 18\hat{k} = m(\hat{i} - 2\hat{j} + 3\hat{k}) + n(2\hat{i} + 3\hat{j} - 4\hat{k})$$

$$\therefore 4\hat{i} + 13\hat{j} - 18\hat{k} = (m + 2n)\hat{i} + (-2m + 3n)\hat{j} + (3m - 4n)\hat{k}$$

By equality of two vectors, we have

$$m + 2n = 4 \quad \dots(1)$$

$$-2m + 3n = 13 \quad \dots(2)$$

$$3m - 4n = -18 \quad \dots(3)$$

Solving (1) and (2) we get, $\therefore m = -2$, $n = 3$

These values of m and n satisfy equation (3) also.

$$\therefore \vec{a} = -2\vec{b} + 3\vec{c}$$

Therefore, \vec{a} is a linear combination of \vec{b} and \vec{c} . Hence, \vec{a} , \vec{b} and \vec{c} are coplanar.



Exercise 5.1

- The vector \vec{a} is directed due north and $|\vec{a}| = 24$. The vector \vec{b} is directed due west and $|\vec{b}| = 7$. Find $|\vec{a} + \vec{b}|$.
- In the triangle PQR, $\vec{PQ} = 2\vec{a}$ and $\vec{QR} = 2\vec{b}$. The mid-point of PR is M. Find following vectors in terms of \vec{a} and \vec{b} .
(i) \vec{PR} (ii) \vec{PM} (iii) \vec{QM}
- OABCDE is a regular hexagon. The points A and B have position vectors \vec{a} and \vec{b} respectively, referred to the origin O. Find, in terms of \vec{a} and \vec{b} the position vectors of C, D and E.
- If ABCDEF is a regular hexagon, show that $\vec{AB} + \vec{AC} + \vec{AD} + \vec{AE} + \vec{AF} = 6\vec{AO}$, where O is the center of the hexagon.
- Check whether the vectors $2\hat{i} + 2\hat{j} + 3\hat{k}$, $-3\hat{i} + 3\hat{j} + 2\hat{k}$ and $3\hat{i} + 4\hat{k}$ form a triangle or not.
- In the figure 5.34 express \vec{c} and \vec{d} in terms of \vec{a} and \vec{b} .
Find a vector in the direction of $\vec{a} = \hat{i} - 2\hat{j}$ that has magnitude 7 units.
- Find the distance from (4, -2, 6) to each of the following :
(a) The XY-plane (b) The YZ-plane
(c) The XZ-plane (d) The X-axis
(e) The Y-axis (f) The Z-axis




Fig 5.34

8. Find the coordinates of the point which is located :
 - (a) Three units behind the YZ-plane, four units to the right of the XZ-plane and five units above the XY-plane.
 - (b) In the YZ-plane, one unit to the right of the XZ-plane and six units above the XY-plane.
9. Find the area of the triangle with vertices (1, 1, 0), (1, 0, 1) and (0, 1, 1).
10. If $\overline{AB} = 2\hat{i} - 4\hat{j} + 7\hat{k}$ and initial point $A \equiv (1, 5, 0)$. Find the terminal point B.
11. Show that the following points are collinear :
 - (i) A (3, 2, -4), B (9, 8, -10), C (-2, -3, 1).
 - (ii) P (4, 5, 2), Q (3, 2, 4), R (5, 8, 0).
12. If the vectors $2\hat{i} - q\hat{j} + 3\hat{k}$ and $4\hat{i} - 5\hat{j} + 6\hat{k}$ are collinear, then find the value of q .
13. Are the four points A(1, -1, 1), B(-1, 1, 1), C(1, 1, 1) and D(2, -3, 4) coplanar? Justify your answer.
14. Express $-\hat{i} - 3\hat{j} + 4\hat{k}$ as linear combination of the vectors $2\hat{i} + \hat{j} - 4\hat{k}$, $2\hat{i} - \hat{j} + 3\hat{k}$ and $3\hat{i} + \hat{j} - 2\hat{k}$.

5.2.1 Section Formula :

Theorem 5 : (Section formula for internal division) Let $A(\bar{a})$ and $B(\bar{b})$ be any two points in the space and $R(\bar{r})$ be a point on the line segment AB dividing it internally in the ratio $m : n$.

Then $\bar{r} = \frac{m\bar{b} + n\bar{a}}{m + n}$

Proof : As R is a point on the line segment AB (A-R-B) and \overline{AR} and \overline{RB} are in same direction.

$$\frac{AR}{RB} = \frac{m}{n}, \text{ so } n(AR) = m(RB)$$

As $m(\overline{RB})$ and $n(\overline{AR})$ have same direction and magnitude,

$$\therefore m(\overline{RB}) = n(\overline{AR})$$

$$\therefore m(\overline{OB} - \overline{OR}) = n(\overline{OR} - \overline{OA})$$

$$\therefore m(\bar{b} - \bar{r}) = n(\bar{r} - \bar{a})$$

$$\therefore m\bar{b} - m\bar{r} = n\bar{r} - n\bar{a}$$

$$\therefore m\bar{b} + n\bar{a} = m\bar{r} + n\bar{r} = (m + n)\bar{r}$$

$$\therefore \bar{r} = \frac{m\bar{b} + n\bar{a}}{m + n}$$

Note :

1. If $A \equiv (a_1, a_2, a_3)$, $B \equiv (b_1, b_2, b_3)$ and $R \equiv (r_1, r_2, r_3)$ divides the segment AB in the ratio $m : n$,

$$\text{then } \bar{r}_i = \frac{m\bar{b}_i + n\bar{a}_i}{m + n}, i = 1, 2, 3.$$




Fig 5.35

2. **(Midpoint formula)** If $R(\bar{r})$ is the mid-point of the line segment AB then $m = n$ so $m : n = 1 : 1$,

$$\therefore \bar{r} = \frac{1\bar{b} + 1\bar{a}}{1+1} \text{ that is } \bar{r} = \frac{\bar{a} + \bar{b}}{2}.$$

Theorem 6 : (Section formula for external division) Let $A(\bar{a})$ and $B(\bar{b})$ be any two points in the space and $R(\bar{r})$ be the third point on the line AB dividing the segment AB externally in the ratio $m : n$. Then $\bar{r} = \frac{m\bar{b} - n\bar{a}}{m - n}$.

Proof : As the point R divides line segment AB externally, we have either A-B-R or R-A-B.

Assume that A-B-R and $\overline{AR} : \overline{BR} = m : n$

$$\therefore \frac{AR}{BR} = \frac{m}{n} \text{ so } n(AR) = m(BR)$$

As $n(\overline{AR})$ and $m(\overline{BR})$ have same magnitude and direction,

$$\therefore n(\overline{AR}) = m(\overline{BR})$$

$$\therefore n(\bar{r} - \bar{a}) = m(\bar{r} - \bar{b})$$

$$\therefore n\bar{r} - n\bar{a} = m\bar{r} - m\bar{b}$$

$$\therefore m\bar{b} - n\bar{a} = m\bar{r} - n\bar{r} = (m - n)\bar{r}$$

$$\therefore \bar{r} = \frac{m\bar{b} - n\bar{a}}{m - n}$$




Fig 5.36

Note : 1) Whenever the ratio in which point R divides the join of two points A and B is required, it is convenient to take the ratio as $k : 1$.

Then, $\bar{r} = \frac{k\bar{b} + \bar{a}}{k + 1}$, if division is internal,

$$\bar{r} = \frac{k\bar{b} - \bar{a}}{k - 1}, \text{ if division is external.}$$




Fig 5.37

2. In ΔABC , centroid G divides the medians internally in ratio $2 : 1$ and is given by (see fig. 5.37)

$$\bar{g} = \frac{\bar{a} + \bar{b} + \bar{c}}{3} \text{ (Verify).}$$

3. In tetrahedron ABCD centroid G divides the line joining the vertex of tetrahedron to centroid of opposite triangle in the ratio $3 : 1$ and is given by

$$\bar{g} = \frac{\bar{a} + \bar{b} + \bar{c} + \bar{d}}{4}.$$




Fig 5.38



Solved Examples

Ex.1. Find the co-ordinates of the point which divides the line segment joining the points A(2, -6, 8) and B(-1, 3, -4). (i) Internally in the ratio 1 : 3. (ii) Externally in the ratio 1 : 3.

Solution : If \vec{a} and \vec{b} are position vectors of the points A and B respectively, then

$$\vec{a} = 2\hat{i} - 6\hat{j} + 8\hat{k} \text{ and } \vec{b} = -\hat{i} + 3\hat{j} - 4\hat{k}.$$

Suppose R(\vec{r}) is the point which divides the line segment joining the points A(\vec{a}) and B(\vec{b}) internally in the ratio 1 : 3 then,

$$\begin{aligned} \vec{r} &= \frac{1(\vec{b}) + 3(\vec{a})}{1+3} = \frac{-1(\hat{i} + 3\hat{j} - 4\hat{k}) + 3(2\hat{i} - 6\hat{j} + 8\hat{k})}{4} \\ \therefore \vec{r} &= \frac{5\hat{i} - 15\hat{j} + 20\hat{k}}{4} \end{aligned}$$

\therefore The coordinates of the point R are $\left(\frac{5}{4}, \frac{-15}{4}, 5\right)$.

Suppose S(\vec{s}) is the point which divides the line joining the points A(\vec{a}) and B(\vec{b}) externally in the ratio 1 : 3 then,

$$\begin{aligned} \vec{s} &= \frac{1\vec{b} - 3\vec{a}}{1-3} = \frac{(-\hat{i} + 3\hat{j} - 4\hat{k}) - 3(2\hat{i} - 6\hat{j} + 8\hat{k})}{-2} \\ \therefore \vec{s} &= \frac{-7\hat{i} + 21\hat{j} - 28\hat{k}}{-2} \end{aligned}$$

\therefore The coordinates of the point S are $\left(\frac{7}{2}, \frac{-21}{2}, 14\right)$.

Ex. 2. If the three points A(3, 2, p), B(q, 8, -10), C(-2, -3, 1) are collinear then find

(i) the ratio in which the point C divides the line segment AB, (ii) the values of p and q.

Solution : Let $\vec{a} = 3\hat{i} + 2\hat{j} + p\hat{k}$, $\vec{b} = q\hat{i} + 8\hat{j} - 10\hat{k}$ and $\vec{c} = -2\hat{i} - 3\hat{j} + \hat{k}$.

Suppose the point C divides the line segment AB in the ratio t : 1,

then by section formula, $\vec{c} = \frac{t\vec{b} + 1\vec{a}}{t+1}$.

$$\begin{aligned} \therefore -2\hat{i} - 3\hat{j} + \hat{k} &= \frac{t(q\hat{i} + 8\hat{j} - 10\hat{k}) + 1(3\hat{i} + 2\hat{j} + p\hat{k})}{t+1} \\ \therefore (-2\hat{i} - 3\hat{j} + \hat{k})(t+1) &= (tq+3)\hat{i} + (8t+2)\hat{j} + (-10t+p)\hat{k} \\ \therefore -2(t+1)\hat{i} - 3(t+1)\hat{j} + (t+1)\hat{k} &= (tq+3)\hat{i} + (8t+2)\hat{j} + (-10t+p)\hat{k} \end{aligned}$$

Using equality of two vectors $-2(t+1) = tq+3 \quad \dots (1)$

$$-3(t+1) = 8t+2 \quad \dots (2)$$

$$t+1 = -10t+p \quad \dots (3)$$

Now equation (2) gives $t = -\frac{5}{11}$

Put $t = -\frac{5}{11}$ in equation (1) and equation (3), we get $q = 9$ and $p = -4$.

The negative sign of t suggests that the point C divides the line segment AB externally in the ratio 5:11.

Ex.3. If A(5, 1, p), B(1, q, p) and C(1, -2, 3) are vertices of triangle and $G\left(r, -\frac{4}{3}, \frac{1}{3}\right)$ is its centroid, then find the values of p, q and r.

Solution : Let $\vec{a} = 5\hat{i} + \hat{j} + p\hat{k}$, $\vec{b} = \hat{i} + q\hat{j} + p\hat{k}$

$$\vec{c} = \hat{i} - 2\hat{j} + 3\hat{k} \quad \text{and} \quad \vec{g} = r\hat{i} - \frac{4}{3}\hat{j} + \frac{1}{3}\hat{k}$$

By centroid formula we have $\vec{g} = \frac{\vec{a} + \vec{b} + \vec{c}}{3}$

$$\therefore 3\vec{g} = \vec{a} + \vec{b} + \vec{c}$$

$$\therefore 3\left(r\hat{i} - \frac{4}{3}\hat{j} + \frac{1}{3}\hat{k}\right) = (5\hat{i} + \hat{j} + p\hat{k}) + (\hat{i} + q\hat{j} + p\hat{k}) + (\hat{i} - 2\hat{j} + 3\hat{k})$$

$$\therefore (3r)\hat{i} - 4\hat{j} + \hat{k} = 7\hat{i} + (q-1)\hat{j} + (2p+3)\hat{k}$$

$$\therefore 3r = 7, -4 = q - 1, 1 = 2p + 3$$

$$\therefore r = \frac{7}{3}, q = -3, p = -1$$

Ex.4. If \vec{a} , \vec{b} , \vec{c} are the position vectors of the points A, B, C respectively and $5\vec{a} - 3\vec{b} - 2\vec{c} = \vec{0}$, then find the ratio in which the point C divides the line segment BA.

Solution : As $5\vec{a} - 3\vec{b} - 2\vec{c} = \vec{0}$

$$\therefore 2\vec{c} = 5\vec{a} - 3\vec{b}$$

$$\therefore \vec{c} = \frac{5\vec{a} - 3\vec{b}}{2}$$

$$\therefore \vec{c} = \frac{5\vec{a} - 3\vec{b}}{5-3}$$

\therefore This shows that the point C divides BA externally in the ratio 5 : 3.

Ex.5. Prove that the medians of a triangle are concurrent.

Solution : Let A, B and C be vertices of a triangle. Let D, E and F be the mid-points of the sides BC, AC and AB respectively. Let \vec{a} , \vec{b} , \vec{c} , \vec{d} , \vec{e} and \vec{f} be position vectors of points A, B, C, D, E and F respectively.

Therefore, by mid-point formula,

$$\therefore \vec{d} = \frac{\vec{b} + \vec{c}}{2}, \vec{e} = \frac{\vec{a} + \vec{c}}{2} \quad \text{and} \quad \vec{f} = \frac{\vec{a} + \vec{b}}{2}$$

$$\therefore 2\vec{d} = \vec{b} + \vec{c}, 2\vec{e} = \vec{a} + \vec{c} \quad \text{and} \quad 2\vec{f} = \vec{a} + \vec{b}$$




Fig 5.39

$$\therefore 2\bar{d} + \bar{a} = \bar{a} + \bar{b} + \bar{c}, \text{ similarly } 2\bar{e} + \bar{b} = 2\bar{f} + \bar{c} = \bar{a} + \bar{b} + \bar{c}$$

$$\therefore \frac{2\bar{d} + \bar{a}}{3} = \frac{2\bar{e} + \bar{b}}{3} = \frac{2\bar{f} + \bar{c}}{3} = \frac{\bar{a} + \bar{b} + \bar{c}}{3} = \bar{g} \text{ (say)}$$

Then we have $\bar{g} = \frac{\bar{a} + \bar{b} + \bar{c}}{3} = \frac{(2)\bar{d} + (1)\bar{a}}{2+1} = \frac{(2)\bar{e} + (1)\bar{b}}{2+1} = \frac{(2)\bar{f} + (1)\bar{c}}{2+1}$

If G is the point whose position vector is \bar{g} , then from the above equation it is clear that the point G lies on the medians AD, BE, CF and it divides each of the medians AD, BE, CF internally in the ratio 2 : 1.

Therefore, three medians are concurrent.

Ex. 6. Prove that the angle bisectors of a triangle are concurrent.

Solution :

Let A, B and C be vertices of a triangle. Let AD, BE and CF be the angle bisectors of the triangle ABD. Let \bar{a} , \bar{b} , \bar{c} , \bar{d} , \bar{e} and \bar{f} be the position vectors of the points A, B, C, D, E and F respectively. Also $AB = z$ $BC = x$ $AC = y$. Now, the angle bisector AD meets the side BC at the point D. Therefore, the point D divides the line segment BC internally in the ratio $AB : AC$, that is $z : y$.

Hence, by section formula for internal division, we have $\bar{d} = \frac{z\bar{c} + y\bar{b}}{z + y}$

Similarly, we get

$$\bar{e} = \frac{x\bar{a} + z\bar{c}}{x + z} \quad \text{and} \quad \bar{f} = \frac{y\bar{b} + x\bar{a}}{y + x}$$

As $\bar{d} = \frac{z\bar{c} + y\bar{b}}{z + y}$

$$\therefore (z + y)\bar{d} = z\bar{c} + y\bar{b}$$

i.e. $(z + y)\bar{d} + x\bar{a} = x\bar{a} + y\bar{b} + z\bar{c}$

similarly $(x + z)\bar{e} + y\bar{b} = x\bar{a} + y\bar{b} + z\bar{c}$

and $(x + y)\bar{f} + z\bar{c} = x\bar{a} + y\bar{b} + z\bar{c}$

$$\therefore \frac{(z + y)\bar{d} + x\bar{a}}{x + y + z} = \frac{(x + z)\bar{e} + y\bar{b}}{x + y + z} = \frac{(x + y)\bar{f} + z\bar{c}}{x + y + z} = \frac{x\bar{a} + y\bar{b} + z\bar{c}}{x + y + z} = \bar{h} \text{ (say)}$$

Then we have

$$\bar{h} = \frac{(y + z)\bar{d} + x\bar{a}}{(y + z) + x} = \frac{(x + z)\bar{e} + y\bar{b}}{(x + z) + y} = \frac{(x + y)\bar{f} + z\bar{c}}{(x + y) + z}$$

That is point H(\bar{h}) divides AD in the ratio $(y + z) : x$, BE in the ratio $(x + z) : y$ and CF in the ratio $(x + y) : z$.

This shows that the point H is the point of concurrence of the angle bisectors AD, BE and CF of the triangle ABC, Thus, the angle bisectors of a triangle are concurrent and H is called incentre of the triangle ABC.




Fig 5.40

Ex. 7. Using vector method, find the incentre of the triangle whose vertices are A(0, 3, 0), B(0, 0, 4) and C(0, 3, 4).

Solution :

$$\text{Let } \vec{a} = 3\hat{j}, \vec{b} = 4\hat{k} \text{ and } \vec{c} = 3\hat{j} + 4\hat{k}$$

$$\therefore \vec{AB} = \vec{b} - \vec{a} = -3\hat{j} + 4\hat{k}, \vec{AC} = \vec{c} - \vec{a} = 4\hat{k}, \vec{BC} = \vec{c} - \vec{b} = 3\hat{j}$$

$$\therefore |\vec{AB}| = 5, |\vec{AC}| = 4, |\vec{BC}| = 3$$

If H (\vec{h}) is the incentre of triangle ABC then,

$$\therefore \vec{h} = \frac{|\vec{BC}|\vec{a} + |\vec{AC}|\vec{b} + |\vec{AB}|\vec{c}}{|\vec{BC}| + |\vec{AC}| + |\vec{AB}|}$$

$$\therefore \vec{h} = \frac{3(3\hat{j}) + 4(4\hat{k}) + 5(3\hat{j} + 4\hat{k})}{3 + 4 + 5}$$

$$\therefore = \frac{9\hat{j} + 16\hat{k} + 15\hat{j} + 20\hat{k}}{12}$$

$$\therefore \vec{h} = \frac{24\hat{j} + 36\hat{k}}{12}$$

$$\therefore \vec{h} = 2\hat{j} + 3\hat{k}$$

And H \equiv (0, 2, 3)

Note : In ΔABC ,

1) P.V. of Centroid is given by $\frac{\vec{a} + \vec{b} + \vec{c}}{3}$

2) P.V. of Incentre is given by $\frac{|\vec{AB}|\vec{c} + |\vec{BC}|\vec{a} + |\vec{AC}|\vec{b}}{|\vec{AB}| + |\vec{BC}| + |\vec{AC}|}$

3) P.V. of Orthocentre is given by $\frac{\tan A\vec{a} + \tan B\vec{b} + \tan C\vec{c}}{\tan A + \tan B + \tan C}$ (Verify)

Ex. 8. If $4\hat{i} + 7\hat{j} + 8\hat{k}$, $2\hat{i} + 3\hat{j} + 4\hat{k}$ and $2\hat{i} + 5\hat{j} + 7\hat{k}$ are the position vectors of the vertices A, B and C respectively of triangle ABC. Find the position vector of the point in which the bisector of $\angle A$ meets BC.

Solution : $\vec{a} = 4\hat{i} + 7\hat{j} + 8\hat{k}$, $\vec{b} = 2\hat{i} + 3\hat{j} + 4\hat{k}$, $\vec{c} = 2\hat{i} + 5\hat{j} + 7\hat{k}$

$$\vec{AC} = \vec{c} - \vec{a} = -2\hat{i} - 2\hat{j} - \hat{k}$$

$$\vec{AB} = \vec{b} - \vec{a} = -2\hat{i} - 4\hat{j} - 4\hat{k}$$

$$\therefore |\vec{AB}| = \sqrt{4 + 16 + 16} = 6 \text{ units}$$

$$\therefore |\vec{AC}| = \sqrt{4 + 4 + 1} = 3 \text{ units}$$

Let D be the point where angle bisector of $\angle A$ meets BC.

D divides BC in the ratio AB : AC




Fig 5.41

$$\begin{aligned}
 \text{i.e. } \bar{d} &= \frac{|\overline{AB}| \bar{c} + |\overline{AC}| \bar{b}}{|\overline{AB}| + |\overline{AC}|} \\
 &= \frac{6(2\hat{i} + 5\hat{j} + 7\hat{k}) + 3(2\hat{i} + 3\hat{j} + 4\hat{k})}{6 + 3} \\
 &= \frac{(12\hat{i} + 30\hat{j} + 42\hat{k}) + (6\hat{i} + 9\hat{j} + 12\hat{k})}{9} \\
 &= \frac{18\hat{i} + 39\hat{j} + 54\hat{k}}{9} \\
 \therefore \bar{d} &= 2\hat{i} + \frac{13}{3}\hat{j} + 6\hat{k}
 \end{aligned}$$

Ex.9. If $G(a, 2, -1)$ is the centroid of the triangle with vertices $P(1, 2, 3)$, $Q(3, b, -4)$ and $R(5, 1, c)$ then find the values of a, b and c .

Solution : As $G(\bar{g})$ is centroid of ΔPQR $\bar{g} = \frac{\bar{p} + \bar{q} + \bar{r}}{3}$

$$a\hat{i} + 2\hat{j} - \hat{k} = \frac{(\hat{i} + 3\hat{j} + 2\hat{k}) + (3\hat{i} + b\hat{j} - 4\hat{k}) + (5\hat{i} + \hat{j} + c\hat{k})}{3} = \frac{(1+3+5)\hat{i} + (3+b+1)\hat{j} + (2-4+c)\hat{k}}{3}$$

by equality of vectors $a = \frac{1+3+5}{3} = \frac{9}{3} = 3 \quad \therefore a = 3$

$$\therefore 2 = \frac{3+b+1}{3} \quad \therefore 6 = 4+b \quad \therefore b = 2$$

$$\therefore -1 = \frac{2-4+c}{3} \quad \therefore -3 = -2+c \quad \therefore c = -1$$

Ex. 10. Find the centroid of tetrahedron with vertices $A(3, -5, 7)$, $B(5, 4, 2)$, $C(7, -7, -3)$, $D(1, 0, 2)$?

Solution : Let $\bar{a} = 3\hat{i} - 5\hat{j} + 7\hat{k}$, $\bar{b} = 5\hat{i} + 4\hat{j} + 2\hat{k}$, $\bar{c} = 7\hat{i} - 7\hat{j} - 3\hat{k}$, $\bar{d} = \hat{i} + 2\hat{k}$ be position vectors of vertices A, B, C & D.

By centroid formula, centroid $G(\bar{g})$ is given by

$$\begin{aligned}
 \bar{g} &= \frac{\bar{a} + \bar{b} + \bar{c} + \bar{d}}{4} \\
 &= \frac{(3\hat{i} - 5\hat{j} + 7\hat{k}) + (5\hat{i} + 4\hat{j} + 2\hat{k}) + (7\hat{i} - 7\hat{j} - 3\hat{k}) + (\hat{i} + 2\hat{k})}{4} \\
 &= \frac{(3+5+7+1)\hat{i} + (-5+4-7+0)\hat{j} + (7+2-3+2)\hat{k}}{4} = \frac{16\hat{i} - 8\hat{j} + 8\hat{k}}{4} \\
 &= 4\hat{i} - 2\hat{j} + 2\hat{k}
 \end{aligned}$$

Therefore, centroid of tetrahedron is $G \equiv (4, -2, 2)$.

Ex. 11. Find the ratio in which point P divides AB and CD where A(2, -3, 4), B(0, 5, 2), C(-1, 5, 3) and D (2, -1, 3). Also, find its coordinates.

Solution : Let point P divides AB in ratio $m : 1$ and CD in ratio $n : 1$.

By section formula,

$$P \equiv \left(\frac{2}{m+1}, \frac{5m-3}{m+1}, \frac{2m+4}{m+1} \right) \equiv \left(\frac{2n-1}{n+1}, \frac{n+5}{n+1}, \frac{3n+3}{n+1} \right)$$

Equating z-coordinates

$$\begin{aligned} \therefore \frac{2m+4}{m+1} &= \frac{3n+3}{n+1} \\ \frac{2m+4}{m+1} &= \frac{3(n+1)}{(n+1)} \\ 2m+4 &= 3(m+1) \\ 2m+4 &= 3m+3 \\ 1 &= m \end{aligned}$$

Also, by equating x-coordinates

$$\begin{aligned} \frac{2}{m+1} &= \frac{2n-1}{n+1} \\ \frac{2}{1+1} &= \frac{2n-1}{n+1} \quad (m=1) \\ n+1 &= 2n-1 \\ 2 &= n \end{aligned}$$

P divides AB in ratio $m : 1$ i.e. 1 : 1 and CD in the ratio $n : 1$ i.e. 2 : 1.

$$P \equiv \left(\frac{2}{1+1}, \frac{5-3}{1+1}, \frac{2+4}{1+1} \right) \equiv (1, 1, 3).$$




Fig 5.42

Ex. 12. In a triangle ABC, D and E are points on BC and AC respectively, such that $BD = 2 DC$ and $AE = 3 EC$. Let P be the point of intersection of AD and BE. Find BP/PF using vector methods.

Solution : Let \vec{a} , \vec{b} , \vec{c} be the position vectors of A, B and C respectively with respect to some origin.

D divides BC in the ratio 2 : 1 and E divides AC in the ratio 3 : 1.

$$\therefore \vec{d} = \frac{\vec{b} + 2\vec{c}}{3} \quad \vec{e} = \frac{\vec{a} + 3\vec{c}}{4}$$

Let point of intersection P of AD and BE divides BE

in the ratio $k : 1$ and AD in the ratio $m : 1$, then position




Fig 5.43

vectors of P in these two cases are $\frac{\vec{b} + k \left(\frac{\vec{a} + 3\vec{c}}{4} \right)}{k+1}$ and $\frac{\vec{a} + m \left(\frac{\vec{b} + 2\vec{c}}{3} \right)}{m+1}$ and respectively.

Equating the position vectors of P we get,

$$\frac{k}{4(k+1)}\bar{a} + \frac{1}{k+1}\bar{b} + \frac{3k}{4(k+1)}\bar{c} = \frac{1}{m+1}\bar{a} + \frac{m}{3(m+1)}\bar{b} + \frac{2m}{3(m+1)}\bar{c}$$

$$\therefore \frac{k}{4(k+1)} = \frac{1}{m+1} \quad \dots (1)$$

$$\frac{1}{k+1} = \frac{m}{3(m+1)} \quad \dots (2)$$

$$\frac{3k}{4(k+1)} = \frac{2m}{3(m+1)} \quad \dots (3)$$

Dividing (3) by (2) we get,

$$\frac{3k}{4} = 2 \text{ i.e. } k = \frac{8}{3} \quad \text{therefore } \frac{BP}{PF} = k : 1 = 8 : 3$$



Exercise 5.2

- Find the position vector of point R which divides the line joining the points P and Q whose position vectors are $2\hat{i} - \hat{j} + 3\hat{k}$ and $-5\hat{i} + 2\hat{j} - 5\hat{k}$ in the ratio 3 : 2 (i) internally (ii) externally.
- Find the position vector of mid-point M joining the points L (7, -6, 12) and N (5, 4, -2).
- If the points A(3, 0, p), B(-1, q, 3) and C(-3, 3, 0) are collinear, then find
 - The ratio in which the point C divides the line segment AB.
 - The values of p and q.
- The position vector of points A and B are $6\bar{a} + 2\bar{b}$ and $\bar{a} - 3\bar{b}$. If the point C divides AB in the ratio 3 : 2 then show that the position vector of C is $3\bar{a} - \bar{b}$.
- Prove that the line segments joining mid-point of adjacent sides of a quadrilateral form a parallelogram.
- D and E divide sides BC and CA of a triangle ABC in the ratio 2 : 3 respectively. Find the position vector of the point of intersection of AD and BE and the ratio in which this point divides AD and BE.
- Prove that a quadrilateral is a parallelogram if and only if its diagonals bisect each other.
- Prove that the median of a trapezium is parallel to the parallel sides of the trapezium and its length is half the sum of parallel sides.
- If two of the vertices of the triangle are A(3, 1, 4) and B(-4, 5, -3) and the centroid of a triangle is G(-1, 2, 1), then find the coordinates of the third vertex C of the triangle.
- In ΔOAB , E is the mid-point of OB and D is the point on AB such that $AD : DB = 2 : 1$. If OD and AE intersect at P, then determine the ratio OP : PD using vector methods.
- If the centroid of a tetrahedron OABC is (1, 2, -1) where $A = (a, 2, 3)$, $B = (1, b, 2)$, $C = (2, 1, c)$ respectively, find the distance of P (a, b, c) from the origin.
- Find the centroid of tetrahedron with vertices K(5, -7, 0), L(1, 5, 3), M(4, -6, 3), N(6, -4, 2) ?

5.3 Product of vectors :

The product of two vectors is defined in two different ways. One form of product results in a scalar quantity while other form gives a vector quantity. Let us study these products and interpret them geometrically.

Angle between two vectors :

When two non zero vectors \vec{a} and \vec{b} are placed such that their initial points coincide, they form an angle θ of measure $0 \leq \theta \leq \pi$.

Angle between \vec{a} and \vec{b} is also denoted as $\widehat{\vec{a} \vec{b}}$.
The angle between the collinear vectors is 0 if they point in the same direction and π if they are in opposite directions.




Fig 5.44

5.3.1 Scalar product of two vectors :

The scalar product of two non-zero vectors \vec{a} and \vec{b} is denoted by $\vec{a} \cdot \vec{b}$, and is defined as $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$, where θ is the angle between \vec{a} and \vec{b} .

$\vec{a} \cdot \vec{b}$ is a real number, that is, a scalar. For this reason, the dot product is also called a scalar product.

Note :

- 1) If either $\vec{a} = 0$ or $\vec{b} = 0$ then θ is not defined and in this case, we define $\vec{a} \cdot \vec{b} = 0$.
- 2) If $\theta = 0$, then $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos 0 = |\vec{a}| |\vec{b}|$. In particular, $\vec{a} \cdot \vec{a} = |\vec{a}|^2$, as $\theta = 0$.
- 3) If $\theta = \pi$, then $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \pi = -|\vec{a}| |\vec{b}|$.
- 4) If \vec{a} and \vec{b} are perpendicular or orthogonal then $\theta = \pi/2$

Conversely if $\vec{a} \cdot \vec{b} = 0$ then either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$ or $\theta = \pi/2$.

$$\text{Also, } \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta = |\vec{b}| |\vec{a}| \cos \theta = \vec{b} \cdot \vec{a}.$$

- 5) Dot product is distributive over vector addition. If \vec{a} , \vec{b} , \vec{c} are any three vectors, then $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$.
- 6) If \vec{a} and \vec{b} are vectors and m, n are scalars, then
 - (i) $(m\vec{a}) \cdot (n\vec{b}) = mn(\vec{a} \cdot \vec{b})$
 - (ii) $(m\vec{a}) \cdot \vec{b} = m(\vec{a} \cdot \vec{b}) = \vec{a} \cdot (m\vec{b})$
- 7) $|\vec{a} \cdot \vec{b}| \leq |\vec{a}| |\vec{b}|$ (This is known as Cauchy Schwartz Inequality).

5.3.2 Finding angle between two vectors :

Angle θ , ($0 \leq \theta \leq \pi$) between two non-zero vectors \vec{a} and \vec{b} is given by $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$, that is

$$\theta = \cos^{-1} \left(\frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \right)$$

Note :

- 1) If $0 \leq \theta < \frac{\pi}{2}$, then $\cos \theta = \frac{\bar{a} \cdot \bar{b}}{|\bar{a}| |\bar{b}|} > 0$, that is $\bar{a} \cdot \bar{b} > 0$.
- 2) If $\theta = \frac{\pi}{2}$, then $\cos \theta = \frac{\bar{a} \cdot \bar{b}}{|\bar{a}| |\bar{b}|} = 0$, that is $\bar{a} \cdot \bar{b} = 0$.
- 3) If $\frac{\pi}{2} < \theta \leq \pi$, then $\cos \theta = \frac{\bar{a} \cdot \bar{b}}{|\bar{a}| |\bar{b}|} < 0$, that is $\bar{a} \cdot \bar{b} < 0$.
- 4) In particular scalar product of \hat{i} , \hat{j} , \hat{k} vectors are
 (i) $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$ and (ii) $\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0$.




Fig 5.45

The scalar product of vectors $\bar{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ and $\bar{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

$$\bar{a} \cdot \bar{b} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) \cdot (b_1\hat{i} + b_2\hat{j} + b_3\hat{k}) = a_1b_1 + a_2b_2 + a_3b_3$$

$$\cos \theta = \frac{\bar{a} \cdot \bar{b}}{|\bar{a}| |\bar{b}|} = \frac{a_1b_1 + a_2b_2 + a_3b_3}{\sqrt{a_1^2 + a_2^2 + a_3^2} \sqrt{b_1^2 + b_2^2 + b_3^2}}$$

5.3.3 Projections :

\overline{PQ} and \overline{PR} represent the vectors \bar{a} and \bar{b} with same initial point P. If M is the foot of perpendicular from R to the line containing \overline{PQ} then $|\overline{PS}|$ is called the scalar projection of \bar{b} on \bar{a} . We can think of it as a shadow of \bar{b} on \bar{a} , when sun is overhead.

$$\text{Scalar Projection of } \bar{b} \text{ on } \bar{a} = |\overline{PS}| = |\bar{b}| \cos \theta = \frac{|\bar{a}| |\bar{b}| \cos \theta}{|\bar{a}|}$$

$$\text{Scalar Projection of } \bar{b} \text{ on } \bar{a} = \frac{\bar{a} \cdot \bar{b}}{|\bar{a}|}$$

$$\text{Vector Projection of } \bar{b} \text{ on } \bar{a} = \overline{PS}$$

$$= |\overline{PS}| \hat{a}, \text{ where } \hat{a} \text{ is unit vector along } \bar{a}$$

$$= \left(\frac{\bar{a} \cdot \bar{b}}{|\bar{a}|} \right) \hat{a}$$

$$= \left(\frac{\bar{a} \cdot \bar{b}}{|\bar{a}|} \right) \frac{\bar{a}}{|\bar{a}|}$$

$$\text{Vector Projection of } \bar{b} \text{ on } \bar{a} = \left(\frac{\bar{a} \cdot \bar{b}}{|\bar{a}|^2} \right) \bar{a}$$




Fig 5.46

5.3.4 Direction Angles and Direction Cosines :

The direction angles of a non-zero vector \bar{a} are the angles $\alpha, \beta,$ and γ ($\in [0, \pi]$) that \bar{a} makes with the positive X-, Y- and Z-axes respectively. These angles completely determine the direction of the vector \bar{a} .

The cosines of these direction angles, that is $\cos \alpha, \cos \beta, \cos \gamma$ are called the direction cosines (abbreviated as d.c.s) of vector \bar{a}

As α is angle between \hat{i} (unit vector) along X-axis and $\bar{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ then.

$$\cos \alpha = \frac{\bar{a} \cdot \hat{i}}{|\bar{a}| |\hat{i}|} = \frac{(a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) \cdot \hat{i}}{|\bar{a}| (1)} = \frac{a_1}{|\bar{a}|}, \text{ Similarly } \cos \beta = \frac{a_2}{|\bar{a}|} \text{ and } \cos \gamma = \frac{a_3}{|\bar{a}|},$$

where $|\bar{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$.

By squaring and adding, we get $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \frac{a_1^2}{|\bar{a}|^2} + \frac{a_2^2}{|\bar{a}|^2} + \frac{a_3^2}{|\bar{a}|^2}$.

As $|\bar{a}|^2 = a_1^2 + a_2^2 + a_3^2$

$\therefore \boxed{\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1}$

Also, $\bar{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$
 $= |\bar{a}| \cos \alpha \hat{i} + |\bar{a}| \cos \beta \hat{j} + |\bar{a}| \cos \gamma \hat{k}$
 $= |\bar{a}| (\cos \alpha \hat{i} + \cos \beta \hat{j} + \cos \gamma \hat{k})$

That is, $\frac{\bar{a}}{|\bar{a}|} = \cos \alpha \hat{i} + \cos \beta \hat{j} + \cos \gamma \hat{k} = \hat{a}$

Which means that the direction cosines of \bar{a} , are components of the unit vector in the direction of \bar{a}




Fig 5.47

Direction cosines (d.c.s) of any line along a vector \bar{a} has same direction cosines as that of \bar{a} .

Direction cosines are generally denoted by l, m, n , where $l = \cos \alpha, m = \cos \beta, n = \cos \gamma$.

As the unit vectors along X, Y- and Z- axes are $\hat{i}, \hat{j}, \hat{k}$. Then \hat{i} makes the direction angles $0, \frac{\pi}{2}, \frac{\pi}{2}$ so its direction cosines are $\cos 0, \cos \frac{\pi}{2}, \cos \frac{\pi}{2}$ that is $1, 0, 0$. Similarly direction cosines of Y- and Z- axes are $0, 1, 0$ and $0, 0, 1$ respectively.

Let \overline{OL} and $\overline{OL'}$ be the vectors in the direction of line LL' . If $\alpha, \beta,$ and γ are direction angles of \overline{OL} then the direction angles of $\overline{OL'}$ are $\pi - \alpha, \pi - \beta,$ and $\pi - \gamma$. Therefore, direction cosines of \overline{OL} are $\cos \alpha, \cos \beta, \cos \gamma$ i.e. l, m, n whereas direction cosines of $\overline{OL'}$ are $\cos(\pi - \alpha), \cos(\pi - \beta)$ and $\cos(\pi - \gamma)$. i.e. $-\cos \alpha, -\cos \beta$ and $-\cos \gamma$. i.e. $-l, -m, -n$. Therefore direction cosines of line LL'

are same as that of vectors \overline{OL} or $\overline{OL'}$ in the direction of line LL' . i.e. either l, m, n or $-l, -m, -n$.
As $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$ so $l^2 + m^2 + n^2 = 1$.

Direction ratios :

Any 3 numbers which are proportional to direction cosines of the line are called the direction ratios (abbreviated as *d.r.s*) of the line. Generally the direction ratios are denoted by a, b, c .

If l, m, n are the direction cosines and a, b, c are direction ratios then

$$a = \lambda l, b = \lambda m, c = \lambda n, \text{ for some } \lambda \in \mathbb{R}.$$

For Example.: If direction cosines of the line are $0, \frac{1}{2}, \frac{\sqrt{3}}{2}$ then $0, 1, \sqrt{3}$ or $0, \sqrt{3}, 3$ or $0, 2\sqrt{3}, 6$ are also direction ratios of the same line.




Fig 5.48

Note : A line has infinitely many direction ratios but unique direction cosines.

Relation between direction ratios and direction cosines :

Let a, b, c be direction ratios and l, m, n be direction cosines of a line.

By definition of d.r.s, $\frac{l}{a} = \frac{m}{b} = \frac{n}{c} = \lambda$ i.e. $l = \lambda a, m = \lambda b, n = \lambda c$.

But $l^2 + m^2 + n^2 = 1$

$$\therefore (\lambda a)^2 + (\lambda b)^2 + (\lambda c)^2 = 1$$

$$\therefore \lambda^2(a^2 + b^2 + c^2) = 1$$

$$\therefore \lambda = \pm \frac{1}{\sqrt{a^2 + b^2 + c^2}}$$

$$\therefore l = \pm \frac{a}{\sqrt{a^2 + b^2 + c^2}}, m = \pm \frac{b}{\sqrt{a^2 + b^2 + c^2}} \text{ and } n = \pm \frac{c}{\sqrt{a^2 + b^2 + c^2}}.$$

Note that direction cosines are similar to the definition of unit vector, that is if $\vec{x} = a\hat{i} + b\hat{j} + c\hat{k}$ be any vector (d.r.s) then $\hat{x} = \pm \frac{\vec{x}}{|\vec{x}|} = \pm \frac{a\hat{i} + b\hat{j} + c\hat{k}}{\sqrt{a^2 + b^2 + c^2}}$ is unit vector (d.c.s) along \vec{x} .



Solved Examples

Ex.1. Find $\vec{a} \cdot \vec{b}$ if $|\vec{a}| = 3$, $|\vec{b}| = \sqrt{6}$, the angle between \vec{a} and \vec{b} is 45° .

Solution : $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta = (3)(\sqrt{6}) \cos 45^\circ = 3\sqrt{6} \left(\frac{\sqrt{2}}{2}\right) = \frac{3}{2} \cdot 2\sqrt{3} = 3\sqrt{3}$

Ex.2. If $\vec{a} = 3\hat{i} + 4\hat{j} - 5\hat{k}$ and $\vec{b} = 3\hat{i} - 4\hat{j} - 5\hat{k}$

- find $\vec{a} \cdot \vec{b}$
- the angle between \vec{a} and \vec{b} .
- the scalar projection of \vec{a} in the direction of \vec{b} .
- the vector projection of \vec{b} along \vec{a} .

Solution : Here $\vec{a} = 3\hat{i} + 4\hat{j} - 5\hat{k}$ and $\vec{b} = 3\hat{i} - 4\hat{j} - 5\hat{k}$

i) $\vec{a} \cdot \vec{b} = (3)(3) + (4)(-4) + (-5)(-5) = 9 - 16 + 25 = 18$

ii) $|\vec{a}| = \sqrt{9+16+25} = \sqrt{50}$, $|\vec{b}| = \sqrt{9+16+25} = \sqrt{50}$

The angle between \vec{a} and \vec{b} is $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{18}{50} \therefore \theta = \cos^{-1} \left(\frac{18}{50} \right)$

iii) The scalar projectio of \vec{a} in the direction of \vec{b} is $\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{18}{\sqrt{50}} = \frac{18}{5\sqrt{2}}$.

iv) The vector projection of \vec{b} along \vec{a} is $\frac{(\vec{a} \cdot \vec{b}) \vec{a}}{|\vec{a}|^2} = \frac{18}{50} (3\hat{i} + 4\hat{j} - 5\hat{k}) = \frac{9}{25} (3\hat{i} + 4\hat{j} - 5\hat{k})$.

Ex.3. Find the value of a for which the vectors $3\hat{i} + 2\hat{j} + 9\hat{k}$ and $\hat{i} + a\hat{j} + 3\hat{k}$ are

- perpendicular
- parallel

Solution : Let $\vec{p} = 3\hat{i} + 2\hat{j} + 9\hat{k}$ and $\vec{q} = \hat{i} + a\hat{j} + 3\hat{k}$

(i) The two vectors are perpendicular if $\vec{p} \cdot \vec{q} = 0$ i.e. $(3\hat{i} + 2\hat{j} + 9\hat{k}) \cdot (\hat{i} + a\hat{j} + 3\hat{k}) = 0$

i.e. $3(1) + 2(a) + 9(3) = 0$. i.e. $2a + 30 = 0$ or $a = -15$.

(ii) The two vectors are parallel if $\frac{3}{1} = \frac{2}{a} = \frac{9}{3}$ i.e. $3a = 2$ i.e. $a = \frac{2}{3}$.

Ex.4. If $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$ and $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$ find the angle between the vectors $2\vec{a} + \vec{b}$ and $\vec{a} + 2\vec{b}$.

Solution : $2\vec{a} + \vec{b} = 2(\hat{i} + 2\hat{j} - 3\hat{k}) + (3\hat{i} - \hat{j} + 2\hat{k}) = 5\hat{i} + 3\hat{j} - 4\hat{k} = \vec{m}$ (say)

$\vec{a} + 2\vec{b} = (\hat{i} + 2\hat{j} - 3\hat{k}) + 2(3\hat{i} - \hat{j} + 2\hat{k}) = 7\hat{i} + \hat{k} = \vec{n}$ (say)

$\therefore \vec{m} \cdot \vec{n} = (2\vec{a} + \vec{b}) \cdot (\vec{a} + 2\vec{b}) = (5\hat{i} + 3\hat{j} - 4\hat{k}) \cdot (7\hat{i} + \hat{k})$

$= (5)(7) + (3)(0) + (-4)(1) = 31$

$$|\vec{m}| = \sqrt{(5)^2 + (3)^2 + (-4)^2} = \sqrt{50}$$

$$|\vec{n}| = \sqrt{(7)^2 + (0)^2 + (1)^2} = \sqrt{50}$$

If θ is the angle between \vec{m} and \vec{n} then

$$\cos \theta = \frac{\vec{m} \cdot \vec{n}}{|\vec{m}| |\vec{n}|} = \frac{31}{\sqrt{50} \times \sqrt{50}} = \frac{31}{50}$$

$$\therefore \theta = \cos^{-1} \left(\frac{31}{50} \right)$$

Ex 5 : If a line makes angle 90° , 60° and 30° with the positive direction of X, Y and Z axes respectively, find its direction cosines.

Solution : Let the d.c.s. of the lines be l, m, n then $l = \cos 90^\circ = 0$, $m = \cos 60^\circ = \frac{1}{2}$
 $n = \cos 30^\circ = \frac{\sqrt{3}}{2}$. Therefore, l, m, n are $0, \frac{1}{2}, \frac{\sqrt{3}}{2}$

Ex. 6 : Find the vector projection of \vec{PQ} on \vec{AB} where P, Q, A, B are the points $(-2, 1, 3)$, $(3, 2, 5)$ $(4, -3, 5)$ and $(7, -5, -1)$ respectively.

Solution : Let the position vectors of P, Q, A, B are $\vec{p}, \vec{q}, \vec{a}, \vec{b}$ respectively

$$\vec{p} = -2\hat{i} + \hat{j} + 3\hat{k}, \vec{q} = 3\hat{i} + 2\hat{j} + 5\hat{k}$$

$$\vec{a} = 4\hat{i} - 3\hat{j} + 5\hat{k}, \vec{b} = 7\hat{i} - 5\hat{j} - \hat{k}$$

$$\therefore \vec{PQ} = \vec{q} - \vec{p} = (3\hat{i} + 2\hat{j} + 5\hat{k}) - (-2\hat{i} + \hat{j} + 3\hat{k}) = 5\hat{i} + \hat{j} + 2\hat{k}$$

$$\text{and } \vec{AB} = \vec{b} - \vec{a} = (7\hat{i} - 5\hat{j} - \hat{k}) - (4\hat{i} - 3\hat{j} + 5\hat{k}) = 3\hat{i} - 2\hat{j} - 6\hat{k}$$

\therefore Vector Projection of \vec{PQ} on \vec{AB}

$$= \frac{\vec{PQ} \cdot \vec{AB}}{|\vec{AB}|^2} \vec{AB} = \frac{(5)(3) + (1)(-2) + (2)(-6)}{(3)^2 + (-2)^2 + (-6)^2} \vec{AB}$$

$$= \frac{1}{49} (3\hat{i} - 2\hat{j} - 6\hat{k}) = \frac{3}{49} \hat{i} - \frac{2}{49} \hat{j} - \frac{6}{49} \hat{k}$$

Ex. 7 : Find the values of λ for which the angle between the vectors

$$\vec{a} = 2\lambda^2 \hat{i} + 4\lambda \hat{j} + \hat{k} \text{ and } \vec{b} = 7\hat{i} - 2\hat{j} + \lambda \hat{k}$$

Solution : If θ is the angle between \vec{a} and \vec{b} , then $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$

If θ is obtuse then $\cos \theta < 0$

$$\therefore \frac{\bar{a} \cdot \bar{b}}{|\bar{a}| |\bar{b}|} < 0 \text{ i.e. } \bar{a} \cdot \bar{b} < 0 \quad \left[\because |\bar{a}| |\bar{b}| > 0 \right]$$

$$\therefore (2\lambda^2 \hat{i} + 4\lambda \hat{j} + \hat{k}) \cdot (7\hat{i} - 2\hat{j} + \lambda \hat{k}) < 0$$

$$\therefore 14\lambda^2 - 8\lambda + \lambda < 0 \text{ i.e. } 14\lambda^2 - 7\lambda < 0$$

$$\therefore 7\lambda(2\lambda - 1) < 0 \text{ i.e. } \lambda \left(\lambda - \frac{1}{2} \right) < 0 \text{ i.e. } 0 < \lambda < \frac{1}{2}$$

Thus the angle between \bar{a} and \bar{b} is obtuse if $0 < \lambda < \frac{1}{2}$

Ex. 8 Find the direction cosines of the vector $2\hat{i} + 2\hat{j} - \hat{k}$

Solutions : Let $|\bar{a}| = 2\hat{i} + 2\hat{j} - \hat{k}$

$$\therefore |\bar{a}| = \sqrt{(2)^2 + (2)^2 + (-1)^2} = \sqrt{9} = 3$$

$$\therefore \hat{a} = \frac{\bar{a}}{|\bar{a}|} = \frac{2\hat{i} + 2\hat{j} - \hat{k}}{3} = \frac{2}{3}\hat{i} + \frac{2}{3}\hat{j} - \frac{1}{3}\hat{k}$$

\therefore The direction cosines of \bar{a} are $\frac{2}{3}, \frac{2}{3}, -\frac{1}{3}$

Ex. 9 : Find the position vector of a point P such that AB is inclined to X axis at 45° and to Y axis at 60° and $OP = 12$ units.

Solution : We have $l = \cos 45^\circ = \frac{1}{\sqrt{2}}, m = \cos 60^\circ = \frac{1}{2}, n = \cos \gamma$

Now $l^2 + m^2 + n^2 = 1$

$$\therefore \frac{1}{2} + \frac{1}{4} + \cos^2 \gamma = 1, \text{ i.e. } \cos^2 \gamma = \frac{1}{4}, \text{ i.e. } n = \cos \gamma = \pm \frac{1}{2}$$

$$\text{Now } \vec{r} = |\vec{r}| \hat{r} = |\vec{r}| (l\hat{i} + m\hat{j} + n\hat{k}) = 12 \left(\frac{1}{\sqrt{12}}\hat{i} + \frac{1}{2}\hat{j} \pm \frac{1}{2}\hat{k} \right)$$

Hence $\vec{r} = 6\sqrt{2}\hat{i} + 6\hat{j} \pm 6\hat{k}$

Ex. 10 A line makes angles of measure 45° and 60° with the positive direction of the Y and Z axes respectively. Find the angle made by the line with the positive directions of the X-axis.

Let α, β, γ be the angles made by the line with positive direction of X, Y and Z axes respectively. Given $\beta = 45^\circ$ and $\gamma = 60^\circ$.

Now $\cos \beta = \cos 45^\circ$ and $\cos \gamma = \cos 60^\circ = \frac{1}{2}$

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

$$\therefore \cos^2 \alpha + \frac{1}{2} + \frac{1}{4} = 1$$

$$\therefore \cos^2 \alpha = \frac{1}{4}$$

$$\therefore \cos \alpha = \pm \frac{1}{2}$$

$\therefore \alpha = 60^\circ$ or 120° There are two lines satisfying given conditions. Their direction angles are $45^\circ, 60^\circ, 60^\circ$ and $45^\circ, 60^\circ, 120^\circ$

Ex. 11. A line passes through the points $(6, -7, -1)$ and $(2, -3, 1)$. Find the direction ratios and the direction cosines of the line so that the angle α is acute.

Solution : Let $A(6, -7, -1)$ and $B(2, -3, 1)$ be the given points. So $\vec{a} = 6\hat{i} - 7\hat{j} - \hat{k}, \vec{b} = 2\hat{i} - 3\hat{j} + \hat{k}$

$$\vec{AB} = \vec{b} - \vec{a} = (2 - 6)\hat{i} + (-3 + 7)\hat{j} + (1 + 1)\hat{k} = -4\hat{i} + 4\hat{j} + 2\hat{k}$$

the direction ratios of \vec{AB} are $-4, 4, 2$.

Let the direction cosines of \vec{AB} be $-4k, 4k, 2k$. Then

$$(-4k)^2 + (4k)^2 + (2k)^2 = 1$$

i.e. $16k^2 + 16k^2 + 4k^2 = 1$ i.e. $36k^2 = 1$ i.e. $k = \pm \frac{1}{6}$

Since the line AB is so directed that the angle α which it makes with the x -Axis is acute,

$$\therefore \cos \alpha = -4k > 0$$

$$\therefore \text{As } k < 0 \quad \therefore k = -\frac{1}{6}$$

$$\therefore \text{the direction cosines of } \vec{AB} \text{ are } -4\left(-\frac{1}{6}\right), 4\left(-\frac{1}{6}\right), 2\left(-\frac{1}{6}\right) \text{ i.e. } \frac{2}{3}, \frac{-2}{3}, \frac{-1}{3}$$

Ex. 12 Prove that the altitudes of a triangle are concurrent.

Solution : Let A, B and C be the vertices of a triangle

Let AD, BE and CF be the altitudes of the triangle ABC , therefore $AD \perp BC, BE \perp AC, CF \perp AB$.

Let $\vec{a}, \vec{b}, \vec{c}, \vec{d}, \vec{e}, \vec{f}$ be the position vectors of A, B, C, D, E, F respectively. Let P be the point of intersection of the altitudes AD and BE with \vec{p} as the position vector.

$$\text{Therefore, } \vec{AP} \perp \vec{BC}, \vec{BP} \perp \vec{AC} \quad \dots\dots(1)$$

To show that the altitudes AD, BE and CF are concurrent, it is sufficient to show that the altitude CF passes through the point P . We will have to prove that \vec{CF} and \vec{CP} are collinear vectors. This can be achieved by showing $\vec{CP} \perp \vec{AB}$




Fig. 5.49

Now from (1) we have

$$\begin{aligned} \overline{AP} \perp \overline{BC} \quad & \text{and} \quad \overline{BP} \perp \overline{AC} \\ \overline{AP} \perp \overline{BC} = 0 \quad & \text{and} \quad \overline{BP} \perp \overline{AC} = 0 \\ \therefore (\overline{p} - \overline{a}) \cdot (\overline{c} - \overline{b}) = 0 \quad & \text{and} \quad (\overline{p} - \overline{b}) \cdot (\overline{c} - \overline{a}) = 0 \\ \therefore \overline{p} \cdot \overline{c} - \overline{p} \cdot \overline{b} - \overline{a} \cdot \overline{c} + \overline{a} \cdot \overline{b} = 0 \quad & \dots (2) \\ \overline{p} \cdot \overline{c} - \overline{p} \cdot \overline{a} - \overline{b} \cdot \overline{c} + \overline{b} \cdot \overline{a} = 0 \quad & \dots (3) \end{aligned}$$

Therefore, subtracting equation (2) from equation (3), we get

$$\begin{aligned} -\overline{p} \cdot \overline{a} + \overline{p} \cdot \overline{b} - \overline{b} \cdot \overline{c} + \overline{a} \cdot \overline{c} &= 0 \quad (\text{Since } \overline{a} \cdot \overline{b} = \overline{b} \cdot \overline{a}) \\ \therefore \overline{p}(\overline{b} - \overline{a}) - \overline{c}(\overline{b} - \overline{a}) &= 0 \\ \therefore (\overline{p} - \overline{c}) \cdot (\overline{b} - \overline{a}) &= 0 \\ \therefore \overline{CP} \cdot \overline{AB} &= 0 \\ \therefore \overline{CP} \perp \overline{AB} \end{aligned}$$

Hence the proof.



Exercise 5.3

- Find two unit vectors each of which is perpendicular to both \overline{u} and \overline{v} , where $\overline{u} = 2\hat{i} + \hat{j} - 2\hat{k}$, $\overline{v} = \hat{i} + 2\hat{j} - 2\hat{k}$
- If \overline{a} and \overline{b} are two vectors perpendicular to each other, prove that $(\overline{a} + \overline{b})^2 = (\overline{a} - \overline{b})^2$
- Find the values of c so that for all real x the vectors $x\hat{i} - 6\hat{j} + 3\hat{k}$ and $x\hat{i} + 2\hat{j} + 2cx\hat{k}$ make an obtuse angle.
- Show that the sum of the length of projections of $p\hat{i} + q\hat{j} + r\hat{k}$ on the coordinate axes, where $p = 2$, $q = 3$ and $r = 4$, is 9.
- Suppose that all sides of a quadrilateral are equal in length and opposite sides are parallel. Use vector methods to show that the diagonals are perpendicular.
- Determine whether \overline{a} and \overline{b} are orthogonal, parallel or neither.
 - $\overline{a} = -9\hat{i} + 6\hat{j} + 15\hat{k}$, $\overline{b} = 6\hat{i} - 4\hat{j} - 10\hat{k}$
 - $\overline{a} = 2\hat{i} + 3\hat{j} - \hat{k}$, $\overline{b} = 5\hat{i} - 2\hat{j} + 4\hat{k}$
 - $\overline{a} = -\frac{3}{5}\hat{i} + \frac{1}{2}\hat{j} + \frac{1}{3}\hat{k}$, $\overline{b} = 5\hat{i} + 4\hat{j} + 3\hat{k}$
 - $\overline{a} = 4\hat{i} - \hat{j} + 6\hat{k}$, $\overline{b} = 5\hat{i} - 2\hat{j} + 4\hat{k}$
- Find the angle P of the triangle whose vertices are P(0, -1, -2), Q(3, 1, 4) and R(5, 7, 1).
- If \hat{p} , \hat{q} and \hat{r} are unit vectors, find i) $\hat{p} \cdot \hat{q}$ ii) $\hat{p} \cdot \hat{r}$ (see fig.5.50)
- Prove by vector method that the angle subtended on semicircle is a right angle.
- If a vector has direction angles 45° and 60° find the third direction angle.
- If a line makes angles 90° , 135° , 45° with the X, Y and Z axes respectively, then find its direction cosines.




Fig.5.50

12. If a line has the direction ratios, 4, -12, 18 then find its direction cosines.
13. The direction ratios of \overline{AB} are -2, 2, 1. If A = (4, 1, 5) and $l(AB) = 6$ units, find B.
14. Find the angle between the lines whose direction cosines l, m, n satisfy the equations $5l+m+3n = 0$ and $5mn - 2nl + 6lm = 0$.

5.4.1 Vector Product of two vectors

In a plane, to describe how a line is tilting we used the notions of slope and angle of inclination. In space, we need to know how plane is tilting. We get this by multiplying two vectors in the plane together to get the third vector perpendicular to the plane. Third vector tell us inclination of the plane. The product we use for finding the third vector is called vector product.

Let \vec{a} and \vec{b} be two nonzero vectors in space. If \vec{a} and \vec{b} are not collinear, they determine a plane. We choose a unit vector \hat{n} perpendicular to the plane by the right-hand rule. Which means \hat{n} points in the way, right thumb points when our fingers curl through the angle from \vec{a} to \vec{b} (See fig 5.51). Then we define a new vector $\vec{a} \times \vec{b}$ as $\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{n}$

The vector product is also called as the cross product of two vectors.

Remarks :

- (i) $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$ as $|\hat{n}| = 1$
- (ii) $\vec{a} \times \vec{b}$ is perpendicular vector to the plane of \vec{a} and \vec{b} .
- (iii) The unit vector

$$\hat{n} \text{ along } \vec{a} \times \vec{b} \text{ is given by } \hat{n} = \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$$

iv) If \vec{a} and \vec{b} are any two coplanar (but non collinear) vectors then any vector \vec{c} in the space can be given by $\vec{c} = x\vec{a} + y\vec{b} + z(\vec{a} \times \vec{b})$ This is because $\vec{a} \times \vec{b}$ is perpendicular to both \vec{a} and \vec{b} and thus \vec{a}, \vec{b} and $\vec{a} \times \vec{b}$ span the whole space.

v) If $\vec{a}, \vec{b}, \hat{n}$ form a right handed triplet, then $\vec{b}, \vec{a}, -\hat{n}$ also form a right handed triplet and

$$\vec{b} \times \vec{a} = |\vec{a}| |\vec{b}| \sin \theta (-\hat{n}) = -|\vec{a}| |\vec{b}| \sin \theta (\hat{n}) = -\vec{a} \times \vec{b}. \text{ Thus vector product is anticommutative.}$$

vi) If \vec{a} and \vec{b} are non zero vectors such that \vec{a} is parallel to \vec{b} .

$$\therefore \theta = 0 \text{ i.e } \sin \theta = 0 \text{ i.e } \vec{a} \times \vec{b} = \vec{0}$$

Conversely if $\vec{a} \times \vec{b} = \vec{0}$, then either $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$ or $\sin \theta = 0$ that is $\theta = 0$.

Thus the cross product of two non zero vectors is zero only when \vec{a} and \vec{b} are collinear. In particular




Fig.5.51

if $\vec{a} = k\vec{b}$ then $\vec{a} \times \vec{b} = k\vec{b} \times \vec{b} = k(\vec{0}) = \vec{0}$

(4) If $\vec{a}, \vec{b}, \vec{c}$ are any three vectors, then

(i) $\vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c}$ Left distributive law

(ii) $(\vec{b} + \vec{c}) \times \vec{a} = \vec{b} \times \vec{a} + \vec{c} \times \vec{a}$ Right distributive law

viii) If \vec{a} and \vec{b} are any two vectors and m, n are two scalars then

(i) $m(\vec{a} \times \vec{b}) = (m\vec{a}) \times \vec{b} = \vec{a} \times (m\vec{b})$

(ii) $m\vec{a} \times n\vec{b} = mn(\vec{a} \times \vec{b})$

ix) If $\hat{i}, \hat{j}, \hat{k}$ are the unit vectors along the co-ordinate axes then

(i) $\hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = \vec{0}$

$\hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i}, \hat{k} \times \hat{i} = \hat{j}$

(ii) $\hat{j} \times \hat{i} = -\hat{k}, \hat{k} \times \hat{j} = -\hat{i}, \hat{i} \times \hat{k} = -\hat{j}$

(Since $\hat{i}, \hat{j}, \hat{k}$ form a right handed triplet)




Fig.5.52

x) If $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ and $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ are two vectors in space then

$$\vec{a} \times \vec{b} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) \times (b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$$

This is given using determinant by $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$

Angle between two vectors: Let θ be the angle between \vec{a} and \vec{b} (so $0 \leq \theta < \pi$),

then $|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$, so $\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|}$

Geometrical meaning of vector product of \vec{a} and \vec{b} :

If \vec{a} and \vec{b} are represented by directed line segments with the same initial point, then they determine a parallelogram with base $|\vec{a}|$, height $|\vec{b}| \sin \theta$ and area of parallelogram

$A = (\text{Base}) (\text{Height}) = |\vec{a}| (|\vec{b}| \sin \theta) = |\vec{a} \times \vec{b}|$




Fig.5.53

Ex. 1 Find the cross product $\vec{a} \times \vec{b}$ and verify that it is orthogonal (perpendicular) to both \vec{a} and \vec{b}



Solved Examples

$$(i) \quad \bar{a} = \hat{i} + \hat{j} - \hat{k}, \quad \bar{b} = 2\hat{i} + 4\hat{j} + 6\hat{k} \qquad (ii) \quad \bar{a} = \hat{i} + 3\hat{j} - 2\hat{k} \quad \bar{b} = -\hat{i} + 5\hat{k}$$

Solution : (i) $\bar{a} \times \bar{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & -1 \\ 2 & 4 & 6 \end{vmatrix} = \begin{vmatrix} 1 & -1 \\ 4 & 6 \end{vmatrix} \hat{i} - \begin{vmatrix} 1 & -1 \\ 2 & 6 \end{vmatrix} \hat{j} + \begin{vmatrix} 1 & 1 \\ 2 & 4 \end{vmatrix} \hat{k}$

$$= [6 - (-4)]\hat{i} - [6 - (-2)]\hat{j} + (4 - 2)\hat{k} = 10\hat{i} - 8\hat{j} + 2\hat{k}$$

Now $\bar{a} \times \bar{b} \cdot \bar{a} = (10\hat{i} - 8\hat{j} + 2\hat{k}) \cdot (\hat{i} + \hat{j} - \hat{k}) = 10 - 8 - 2 = 0$ and

$$\bar{a} \times \bar{b} \cdot \bar{b} = (10\hat{i} - 8\hat{j} + 2\hat{k}) \cdot (2\hat{i} + 4\hat{j} - 6\hat{k}) = 20 - 32 + 12 = 0$$

so $\bar{a} \times \bar{b}$ is orthogonal to both \bar{a} and \bar{b} .

$$(ii) \quad \bar{a} \times \bar{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 3 & -2 \\ -1 & 0 & 5 \end{vmatrix} = \begin{vmatrix} 3 & -2 \\ 0 & 5 \end{vmatrix} \hat{i} - \begin{vmatrix} 1 & -2 \\ -1 & 5 \end{vmatrix} \hat{j} + \begin{vmatrix} 1 & 3 \\ -1 & 0 \end{vmatrix} \hat{k}$$

$$= (15 - 0)\hat{i} - (5 - 2)\hat{j} + [0 - (-3)]\hat{k} = 15\hat{i} - 3\hat{j} + 3\hat{k}$$

Now $(\bar{a} \times \bar{b}) \cdot \bar{a} = (15\hat{i} - 3\hat{j} + 3\hat{k}) \cdot (\hat{i} + 3\hat{j} - 2\hat{k}) = 15 - 9 - 6 = 0,$

and $(\bar{a} \times \bar{b}) \cdot \bar{b} = (15\hat{i} - 3\hat{j} + 3\hat{k}) \cdot (-\hat{i} + 5\hat{k}) = 15 + 0 + 15 = 0,$ $\bar{a} \times \bar{b}$ is orthogonal to both \bar{a} and \bar{b}

Ex.2: Find all vectors of magnitude $10\sqrt{3}$ that are perpendicular to the plane of $\hat{i} + 2\hat{j} + \hat{k}$ and $-\hat{i} + 3\hat{j} + 4\hat{k}$

Solution : Let $\bar{a} = \hat{i} + 2\hat{j} + \hat{k}$ and $\bar{b} = -\hat{i} + 3\hat{j} + 4\hat{k}$. Then

$$\bar{a} \times \bar{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & 1 \\ -1 & 3 & 4 \end{vmatrix} = \hat{i}(8 - 3) - \hat{j}(4 + 1) + \hat{k}(3 + 2) = 5\hat{i} - 5\hat{j} + 5\hat{k} = \bar{m} \text{ (say)}$$

$$\therefore |\bar{a} \times \bar{b}| = \sqrt{(5)^2 + (-5)^2 + (5)^2} = \sqrt{3(5)^2} = 5\sqrt{3} = |\bar{m}|$$

Therefore, unit vector perpendicular to the plane of \vec{a} and \vec{b} is given by

$$\hat{m} = \frac{\vec{m}}{|\vec{m}|} = \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} = \frac{5\hat{i} - 5\hat{j} + 5\hat{k}}{5\sqrt{3}} = \frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}}$$

Hence, vectors of magnitude of $10\sqrt{3}$ that are perpendicular to plane of \vec{a} and \vec{b} are

$$\pm 10\sqrt{3} \hat{m} = \pm 10\sqrt{3} \left(\frac{\hat{i} - \hat{j} + \hat{k}}{\sqrt{3}} \right), \text{ i.e. } \pm 10(\hat{i} - \hat{j} + \hat{k}).$$

Ex.3: If $\vec{u} + \vec{v} + \vec{w} = \vec{0}$, show that $\vec{u} \times \vec{v} = \vec{v} \times \vec{w} = \vec{w} \times \vec{u}$.

Solution : Suppose that $\vec{u} + \vec{v} + \vec{w} = \vec{0}$. Then

$$\begin{aligned} (\vec{u} + \vec{v} + \vec{w}) \times \vec{v} &= \vec{0} \times \vec{v} \\ \vec{u} \times \vec{v} + \vec{v} \times \vec{v} + \vec{w} \times \vec{v} &= \vec{0}. \end{aligned}$$

But $\vec{v} \times \vec{v} = \vec{0}$

Thus $\vec{u} \times \vec{v} + \vec{w} \times \vec{v} = \vec{0}$.

Thus $\vec{u} \times \vec{v} = -\vec{w} \times \vec{v} = \vec{v} \times \vec{w}$.

Similarly, we have $\vec{v} \times \vec{w} = \vec{w} \times \vec{u}$.

Ex 5. If $\vec{a} = 3\hat{i} - \hat{j} + 2\hat{k}$, $\vec{b} = 2\hat{i} + \hat{j} - \hat{k}$, $\vec{c} = \hat{i} - 2\hat{j} + 2\hat{k}$, find $(\vec{a} \times \vec{b}) \times \vec{c}$ and $\vec{a} \times (\vec{b} \times \vec{c})$ and hence show that $(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$.

Solution :

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -1 & 2 \\ 2 & 1 & -1 \end{vmatrix} = -\hat{i} + 7\hat{j} + 5\hat{k}$$

$$\begin{aligned} (\vec{a} \times \vec{b}) \times \vec{c} &= (-\hat{i} + 7\hat{j} + 5\hat{k}) \times (\hat{i} - 2\hat{j} + 2\hat{k}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 7 & 5 \\ 1 & -2 & 2 \end{vmatrix} \\ &= 24\hat{i} + 7\hat{j} - 5\hat{k} \quad \dots(1) \end{aligned}$$

Now, $\vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -1 \\ 1 & -2 & 2 \end{vmatrix} = -5\hat{j} - 5\hat{k}$

$$\vec{a} \times (\vec{b} \times \vec{c}) = (3\hat{i} - \hat{j} + 2\hat{k}) \times (-5\hat{j} - 5\hat{k})$$

$$\vec{a} \times (\vec{b} \times \vec{c}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -1 & 2 \\ 0 & -5 & -5 \end{vmatrix} = 15(\hat{i} + \hat{j} - \hat{k}) \quad \dots(2)$$

from (1) and (2), we conclude that

$$(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$$

Ex. 6 Find the area of the triangle with vertices (1,2,0), (1,0,2), and (0,3,1).

Solution : If A = (1, 2, 0), B = (1, 0, 2) and C = (0, 3, 1), then $\overline{AB} = -2j + 2k$, $\overline{AC} = -i + j + k$ and

the area of triangle ABC is $\frac{1}{2} |\overline{AB} \times \overline{AC}|$ and $\overline{AB} \times \overline{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & -2 & 2 \\ -1 & 1 & 1 \end{vmatrix} = -4\hat{i} - 2\hat{j} - 2\hat{k}$

$$\frac{|\overline{AB} \times \overline{AC}|}{2} = \frac{|-4i - 2j - 2k|}{2} = \frac{\sqrt{16+4+4}}{2} = \frac{\sqrt{24}}{2} = \frac{2\sqrt{6}}{2} = \sqrt{6} \text{ sq. units}$$

Ex. 7 Find the area of the parallelogram with vertices K(1, 2, 3), L(1, 3, 6), M(3, 8, 6) and N(3, 7,3)

Solution : The parallelogram is determined by the vectors $\overline{KL} = \hat{j} + 3\hat{k}$ and $\overline{KN} = 2\hat{i} + 5\hat{j}$, so the area of parallelogram KLMN is

$$|\overline{KL} \times \overline{KN}| = \left\| \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & 3 \\ 2 & 5 & 0 \end{vmatrix} \right\| = |(-15)\hat{i} - (-6)\hat{j} + (-2)\hat{k}| = |-15\hat{i} + 6\hat{j} - 2\hat{k}| = \sqrt{265} \text{ square units}$$

Ex. 8 Find $|\vec{u} \times \vec{v}|$ if




Fig 5.54

Solution : i) We have $|\vec{u} \times \vec{v}| = |\vec{u}| |\vec{v}| \sin \theta = (4)(5) \sin 45^\circ = 20 \frac{1}{\sqrt{2}} = 10\sqrt{2}$




Fig 5.55

ii) If we sketch \vec{u} and \vec{v} starting from the same initial point, we see that the angle between them is 60° . we have $|\vec{u} \times \vec{v}| = |\vec{u}| |\vec{v}| \sin \theta = (12)(16) \sin 60^\circ = 192 \frac{\sqrt{3}}{2} = 96\sqrt{3}$.

Ex. 9 Show that $(\vec{a} - \vec{b}) \times (\vec{a} + \vec{b}) = 2(\vec{a} \times \vec{b})$

Solution : Using distributive property $(\vec{a} - \vec{b}) \times (\vec{a} + \vec{b}) = \vec{a} \times \vec{a} + \vec{a} \times \vec{b} - \vec{b} \times \vec{a} - \vec{b} \times \vec{b}$ ($\because \vec{a} \times \vec{b} = -\vec{b} \times \vec{a}$)
 $= \vec{0} + \vec{a} \times \vec{b} + \vec{a} \times \vec{b} - \vec{0}$ ($\because \vec{a} \times \vec{a} = \vec{0}$)
 $= 2(\vec{a} \times \vec{b})$

Ex. 10 Show that the three points with position vectors $3\hat{j} - 2\hat{j} + 4\hat{k}$, $\hat{i} + \hat{j} + \hat{k}$ and $-\hat{i} + 4\hat{j} - 2\hat{k}$ respectively are collinear.

Solution : Let A, B, C be the given three points.

$\vec{a} = 3\hat{j} - 2\hat{j} + 4\hat{k}$, $\vec{b} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = -\hat{i} + 4\hat{j} - 2\hat{k}$ To show that points A, B, C are collinear.

Now $\vec{AB} = \vec{b} - \vec{a} = -2\hat{i} + 3\hat{j} - 3\hat{k}$, $\vec{AC} = \vec{c} - \vec{a} = -4\hat{i} + 6\hat{j} - 6\hat{k}$.

$$\begin{aligned} \vec{AB} \times \vec{AC} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -2 & 3 & -3 \\ -4 & 6 & -6 \end{vmatrix} \\ &= (-18 + 18)\hat{i} - (12 - 12)\hat{j} + (-12 + 12)\hat{k} \\ &= 0\hat{i} - 0\hat{j} + 0\hat{k} \\ &= \vec{0} \end{aligned}$$

Vectors \vec{AB} and \vec{AC} are collinear, but the point A is common, therefore points A, B, C are collinear.

Ex. 11 Find a unit vector perpendicular to \vec{PQ} and \vec{PR} where P \equiv (2,2,0), Q \equiv (0,3,5) and R \equiv (5,0,3). Also find the sine of angle between \vec{PQ} and \vec{PR}

Solution : $\vec{PQ} = \vec{q} - \vec{p} = -2\hat{i} + \hat{j} + 5\hat{k}$

and $\vec{PR} = \vec{r} - \vec{p} = 3\hat{i} - 2\hat{j} + 3\hat{k}$

Now $|\vec{PQ}| = \sqrt{4 + 1 + 25} = \sqrt{30}$

and $|\vec{PR}| = \sqrt{9 + 4 + 9} = \sqrt{22}$

$$\therefore \vec{PQ} \times \vec{PR} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -2 & 1 & 5 \\ 3 & -2 & 3 \end{vmatrix}$$

$$= (3+10)\hat{i} - (-6-15)\hat{j} + (4-3)\hat{k}$$

$$= 13\hat{i} + 21\hat{j} + \hat{k}$$

$$\therefore |\overline{PQ} \times \overline{PR}| = \sqrt{169 + 441 + 1} = \sqrt{611}$$

If \hat{n} is a unit vector perpendicular to \overline{PQ} and \overline{PR} , then

$$\hat{n} = \frac{\overline{PQ} \times \overline{PR}}{|\overline{PQ} \times \overline{PR}|} = \frac{13\hat{i} + 21\hat{j} + \hat{k}}{\sqrt{611}}$$

If θ is the angle between \overline{PQ} and \overline{PR} then $\sin \theta = \frac{|\overline{PQ} \times \overline{PR}|}{|\overline{PQ}| |\overline{PR}|} = \frac{\sqrt{611}}{\sqrt{30} \sqrt{22}}$

Ex. 12 If $|\vec{a}| = 5$, $|\vec{b}| = 13$ and $|\vec{a} \times \vec{b}| = 25$, find $\vec{a} \cdot \vec{b}$.

Solution : Given $|\vec{a} \times \vec{b}| = 25$

$$\therefore |\vec{a}| \cdot |\vec{b}| \sin \theta = 25 \quad (\theta \text{ is the angle between } \vec{a} \text{ and } \vec{b})$$

$$5 \times 13 \sin \theta = 25$$

$$\sin \theta = \frac{25}{5 \times 13} = \frac{5}{13}$$

$$\therefore \cos \theta = \sqrt{1 - \sin^2 \theta} = \sqrt{1 - \frac{25}{169}} = \pm \frac{12}{13}$$

$$\therefore \vec{a} \cdot \vec{b} = |\vec{a}| \cdot |\vec{b}| \cos \theta$$

$$= 5 \times 13 \times \left(\pm \frac{12}{13} \right)$$

$$= \pm 60$$

Thus $\vec{a} \cdot \vec{b} = 60$ if $0 < \theta < \pi/2$ and

$$\vec{a} \cdot \vec{b} = -60 \text{ if } \pi/2 < \theta < \pi.$$

Ex. 13.: Direction ratios of two lines satisfy the relation $2a-b+2c = 0$ and $ab+bc+ca = 0$. Show that the lines are perpendicular.

Solution : Given equations are $2a-b+2c = 0$ i.e $b = 2a+2c$ (I) and $ab+bc+ca = 0$(II)

Put $b = 2a+2c$ in equation (II), we get

$$a(2a+2c) + (2a+2c)c + ca = 0$$

$$2a^2+2ac+2ac+2c^2+ac = 0$$

$$2a^2+5ac+2c^2 = 0$$

$$\therefore (2a+c)(a+2c) = 0$$

Case I : i.e. $2a+c = 0 \therefore 2a = -c \dots(III)$

Using this equation (I) becomes $b = -c + 2c = c$ i.e. $b = c \dots(IV)$ from (III) and (IV) we get,

$$\frac{a}{-\frac{1}{2}} = \frac{b}{1} = \frac{c}{1} \text{ Direction ratios of 1st line are i.e. } -\frac{1}{2}, 1, 1 \text{ i.e. } -1, 2, 2 = \bar{p} \text{ (say)}$$

Case II: i.e. $a + 2c=0, \therefore a = -2c \dots(V)$

Using this equation (I) becomes

$$b = 2(-2c) + 2c = -2c \text{ i.e. } b = -2c \dots(VI)$$

From (V) and (VI), we get

$$\frac{a}{-2} = \frac{b}{-2} = \frac{c}{1}$$

\therefore Direction ratios of second line are $-2, -2, 1$ i.e. $2, 2, -1 = \bar{q}$ (Say)

$$\text{Now } \bar{p} \cdot \bar{q} = (-1\hat{i} + 2\hat{j} + 2\hat{k}) \cdot (2\hat{i} + 2\hat{j} - \hat{k}) = -2 + 4 - 2 = 0$$

\therefore The lines are perpendicular.

Ex. 13 Find the direction cosines of the line which is perpendicular to the lines with direction ratios $-1, 2, 2$ and $0, 2, 1$.

Solution : Given $-1, 2, 2$ and $0, 2, 1$ be direction ratios of lines L_1 and L_2 .

Let l, m, n be direction cosines of line L. As line L is perpendicular to lines L_1 and L_2 .

$$\text{Then } -l+2m+2n = 0 \quad \text{and} \quad 2m+n = 0$$

$$2m+n = 0$$

$$\therefore 2m = -n$$

$$\therefore \frac{m}{-1} = \frac{n}{2}$$

$$\therefore \frac{m}{-1} = \frac{n}{2} \quad \dots(I) \text{ and}$$

$$-l+2m+2n = 0 \quad \text{becomes}$$

$$-l-n+2n = 0$$

$$-l+n = 0$$

$$l = n$$

$$\frac{l}{1} = \frac{n}{1} \quad \text{i.e.} \quad \frac{l}{2} = \frac{n}{2} \quad \dots(II)$$

$$\text{from (I) and (II) } \frac{l}{2} = \frac{m}{-1} = \frac{n}{2}$$

The direction ratios of line L are $2, -1, 2$ and the direction cosines of line L are $\frac{2}{3}, \frac{-1}{3}, \frac{2}{3}$.

Ex. 15 If M is the foot of the perpendicular drawn from A(4, 3, 2) on the line joining the points B(2, 4, 1) and C (4, 5, 3), find the coordinates of M.

Let the point M divides BC internally in the ratio k:1

$$\therefore M \equiv \left(\frac{4k+2}{k+1}, \frac{5k+4}{k+1}, \frac{3k+1}{k+1} \right) \dots \text{(I)}$$

\therefore Direction ratios of AM are

$$\frac{4k+2}{k+1} - 4, \frac{5k+4}{k+1} - 3, \frac{3k+1}{k+1} - 2 = \bar{p} \text{ (say)}$$

i.e. $\frac{-2}{k+1}, \frac{2k+1}{k+1}, \frac{k-1}{k+1}$ and direction ratios of BC are 4-2, 5-4, 3-1 i.e. 2, 1, 2 = \bar{q} (say)

since AM is perpendicular to BC, $\bar{p} \cdot \bar{q} = 0$

$$\text{i.e. } 2 \frac{(-2)}{k+1} + 1 \frac{(2k+1)}{k+1} + 2 \frac{(k-1)}{k+1} = 0$$

$$\text{i.e. } -4 + 2k + 1 + 2k - 2 = 0$$

$$\therefore 4k - 5 = 0$$

$$k = \frac{5}{4}$$

\therefore

from (I) $M \equiv \left(\frac{28}{9}, \frac{41}{9}, \frac{19}{9} \right)$



Exercise 5.4

- If $\bar{a} = 2\hat{i} + 3\hat{j} - \hat{k}$, $\bar{b} = \hat{i} - 4\hat{j} + 2\hat{k}$ find $(\bar{a} + \bar{b}) \times (\bar{a} - \bar{b})$
- Find a unit vector perpendicular to the vectors $\hat{j} + 2\hat{k}$ and $\hat{i} + \hat{j}$.
- If $\bar{a} \cdot \bar{b} = \sqrt{3}$ and $\bar{a} \times \bar{b} = 2\hat{i} + \hat{j} + 2\hat{k}$, find the angle between \bar{a} and \bar{b} .
- If $\bar{a} = 2\hat{i} + \hat{j} - 3\hat{k}$ and $\bar{b} = \hat{i} - 2\hat{j} + \hat{k}$, find a vector of magnitude 5 perpendicular to both \bar{a} and \bar{b} .
- Find i) $\bar{u} \cdot \bar{v}$ if $|\bar{u}| = 2, |\bar{v}| = 5, |\bar{u} \times \bar{v}| = 8$ ii) $|\bar{u} \times \bar{v}|$ if $|\bar{u}| = 10, |\bar{v}| = 2, \bar{u} \cdot \bar{v} = 12$
- Prove that $2(\bar{a} - \bar{b}) \times 2(\bar{a} + \bar{b}) = 4(\bar{a} \times \bar{b})$
- If $\bar{a} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\bar{b} = 4\hat{i} - 3\hat{j} + \hat{k}$ and $\bar{c} = \hat{i} - \hat{j} + 2\hat{k}$ verify that $\bar{a} \times (\bar{b} + \bar{c}) = \bar{a} \times \bar{b} + \bar{a} \times \bar{c}$
- Find the area of the parallelogram whose adjacent sides are the vectors $\bar{a} = 2\hat{i} - 2\hat{j} + \hat{k}$ and $\bar{b} = \hat{i} - 3\hat{j} - 3\hat{k}$.
- Show that vector area of a quadrilateral ABCD is $\frac{1}{2} (\overline{AC} \times \overline{BD})$, where AC and BD are its diagonals.

10. Find the area of parallelogram whose diagonals are determined by the vectors $\vec{a} = 3\hat{i} - \hat{j} - 2\hat{k}$, and $\vec{b} = -\hat{i} + 3\hat{j} - 3\hat{k}$
11. If $\vec{a}, \vec{b}, \vec{c}$ and \vec{d} are four distinct vectors such that $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$ and $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$, prove that $\vec{a} - \vec{d}$ is parallel to $\vec{b} - \vec{c}$
12. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{j} - \hat{k}$, find a vector \vec{b} satisfying $\vec{a} \times \vec{b} = \vec{c}$ and $\vec{a} \cdot \vec{b} = 3$
13. Find \vec{a} , if $\vec{a} \times \hat{i} + 2\vec{a} - 5\hat{j} = 0$.
14. If $|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$ and $\vec{a} \cdot \vec{b} < 0$, then find the angle between \vec{a} and \vec{b}
15. Prove by vector method that $\sin(\alpha + \beta) = \sin\alpha \cos\beta + \cos\alpha \sin\beta$.
16. Find the direction ratios of a vector perpendicular to the two lines whose direction ratios are
 - (i) $-2, 1, -1$ and $-3, -4, 1$
 - (ii) $1, 3, 2$ and $-1, 1, 2$
17. Prove that two vectors whose direction cosines are given by relations $al + bm + cn = 0$ and $fmn + gnl + hlm = 0$ are perpendicular if $\frac{f}{a} + \frac{g}{b} + \frac{h}{c} = 0$
18. If A(1, 2, 3) and B(4, 5, 6) are two points, then find the foot of the perpendicular from the point B to the line joining the origin and point A.

5.5.1 Scalar Triple Product :

We define the scalar triple product of three vectors $\vec{a}, \vec{b}, \vec{c}$ (Order is important) which is denoted by

$[\vec{a} \ \vec{b} \ \vec{c}]$ and is defined as $\vec{a} \cdot (\vec{b} \times \vec{c})$

For $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$, $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ and $\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$

$$[\vec{a} \ \vec{b} \ \vec{c}] = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

Scalar triple product is also called as box product.

Properties of scalar triple product:

Using the properties of determinant, we get following properties of scalar triple product.

- (1) A cyclic change of vectors $\vec{a}, \vec{b}, \vec{c}$ in a scalar triple product does not change its value

i.e. $[\vec{a} \ \vec{b} \ \vec{c}] = [\vec{c} \ \vec{a} \ \vec{b}] = [\vec{b} \ \vec{c} \ \vec{a}]$

This follows as a cyclic change is equivalent to interchanging a pair of rows in the determinant two times.

(2) A single interchange of vectors in a scalar triple product changes the sign of its value.

$$\text{i.e. } [\bar{a} \bar{b} \bar{c}] = -[\bar{b} \bar{a} \bar{c}] = -[\bar{c} \bar{b} \bar{a}] = -[\bar{a} \bar{c} \bar{b}]$$

This follows as interchange of any 2 rows changes the value of determinant by sign only.

(3) If a row of determinant can be expressed as a linear combination of other rows then the determinant is zero. Using this fact we get following properties.

The scalar triple product of vectors is zero if any one of the following is true.

- (i) One of the vectors is a zero vector.
- (ii) Any two vectors are collinear.
- (iii) The three vectors are coplanar.

(4) An interchange of 'dot' and 'cross' in a scalar triple product does not change its value

$$\text{i.e. } \bar{a} \cdot (\bar{b} \times \bar{c}) = (\bar{a} \times \bar{b}) \cdot \bar{c}$$

This is followed by property (1) and the commutativity of dot product.

Theorem 9 : The volume of parallelopiped with coterminus edges as \bar{a}, \bar{b} and \bar{c} is $[\bar{a} \bar{b} \bar{c}]$

Proof : Let $\overline{OA} = \bar{a}, \overline{OB} = \bar{b}$ and $\overline{OC} = \bar{c}$ be coterminus edges of parallelopiped.

Let AP be the height of the parallelopiped.

Volume of Parallelopiped = (Area of base parallelogram OBDC) (Height AP)

$$\text{But AP} = \text{Scalar Projection of } \bar{a} \text{ on } (\bar{b} \times \bar{c}) = \frac{(\bar{b} \times \bar{c}) \cdot \bar{a}}{|\bar{b} \times \bar{c}|} \left(\because \text{scalar projection of } \bar{p} \text{ on } \bar{q} \text{ is } \frac{\bar{p} \cdot \bar{q}}{q} \right)$$

$$\text{and area of parallelogram OBDC} = |\bar{b} \times \bar{c}|$$

$$\text{volume of parallelopiped} = \frac{(\bar{b} \times \bar{c}) \cdot \bar{a}}{|\bar{b} \times \bar{c}|} |\bar{b} \times \bar{c}|$$

$$= \bar{a} \cdot (\bar{b} \times \bar{c}) = [\bar{a} \bar{b} \bar{c}]$$




Fig. 5.56

Theorem 10. The volume of a tetrahedron with coterminus

edges \bar{a}, \bar{b} and \bar{c} is $\frac{1}{6} [\bar{a} \bar{b} \bar{c}]$.

Proof : Let $\overline{OA} = \bar{a}, \overline{OB} = \bar{b}$ and $\overline{OC} = \bar{c}$ be coterminus edges of tetrahedron OABC.

Let AP be the height of tetrahedron

Volume of tetrahedron = $\frac{1}{3}$ (Area of base ΔOCB) (Height AP)

$$\text{But AP} = \text{Scalar Projection of } \bar{a} \text{ on } (\bar{b} \times \bar{c}) = \frac{(\bar{b} \times \bar{c}) \cdot \bar{a}}{|\bar{b} \times \bar{c}|} \left(\because \text{scalar projection of } \bar{p} \text{ on } \bar{q} \text{ is } \frac{\bar{p} \cdot \bar{q}}{q} \right)$$




Fig. 5.57

$$\text{Area of } \triangle OBC = \frac{1}{2} |\vec{b} \times \vec{c}|$$

$$\begin{aligned} \text{Volume of tetrahedron} &= \frac{1}{3} \times \frac{1}{2} |\vec{b} \times \vec{c}| \frac{(\vec{b} \times \vec{c}) \cdot \vec{a}}{|\vec{b} \times \vec{c}|} \\ &= \frac{1}{6} [(\vec{b} \times \vec{c}) \cdot \vec{a}] = \frac{1}{6} [\vec{a} \cdot \vec{b} \times \vec{c}] \end{aligned}$$

5.5.2 : Vector triple product :

For vectors \vec{a} , \vec{b} and \vec{c} in the space, we define the vector triple product without proof as

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c}) \vec{b} - (\vec{a} \cdot \vec{b}) \vec{c}.$$




Fig. 5.58

Properties of vector triple product

- 1) $\vec{a} \times (\vec{b} \times \vec{c}) = -(\vec{b} \times \vec{c}) \times \vec{a} (\because \vec{p} \times \vec{q} = -\vec{q} \times \vec{p})$
- 2) $(\vec{a} \times \vec{b}) \times \vec{c} = (\vec{a} \cdot \vec{c}) \vec{b} - (\vec{b} \cdot \vec{c}) \vec{a}$
- 3) $\vec{a} \times (\vec{b} \times \vec{c}) \neq (\vec{a} \times \vec{b}) \times \vec{c}$
- 4) $\hat{i} \times (\hat{j} \times \hat{k}) = \vec{0}$
- 5) $\vec{a} \times (\vec{b} \times \vec{c})$ is linear combination of \vec{b} and \vec{c} , hence it is coplanar with \vec{b} and \vec{c} .



Solved Examples

Ex. 1 Find the volume of the parallelepiped determined by the vectors \vec{a} , \vec{b} and \vec{c}

- (i) $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}, \vec{b} = -\hat{i} + \hat{j} + 2\hat{k}, \vec{c} = 2\hat{i} + \hat{j} + 4\hat{k}$
- (ii) $\vec{a} = \hat{i} + \hat{j}, \vec{b} = \hat{j} + \hat{k}, \vec{c} = \hat{i} + \hat{j} + \hat{k}$

Solution (i) The volume of the parallelepiped determined by \vec{a} , \vec{b} and \vec{c} is the magnitude of their scalar triple product $\vec{a} \cdot (\vec{b} \times \vec{c})$

$$\text{and } \vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} 1 & 2 & 3 \\ -1 & 1 & 2 \\ 2 & 1 & 4 \end{vmatrix} = 1 \begin{vmatrix} 1 & 2 \\ 1 & 4 \end{vmatrix} - 2 \begin{vmatrix} -1 & 2 \\ 2 & 4 \end{vmatrix} + 3 \begin{vmatrix} -1 & 1 \\ 2 & 1 \end{vmatrix} = 1(4-2) - 2(-4-4) + 3(-1-2) = 9.$$

Thus the volume of the parallelepiped is 9 cubic units.

$$(ii) \vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{vmatrix} = 1 \begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} + 0 \begin{vmatrix} 0 & 1 \\ 1 & 1 \end{vmatrix} = 0 + 1 + 0 = 1.$$

So the volume of the parallelepiped is 1 cubic unit.

Ex. 2. Find the scalar triple product $[\vec{u} \ \vec{v} \ \vec{w}]$ and verify that the vectors $\vec{u} = \hat{i} + 5\hat{j} - 2\hat{k}, \vec{v} = 3\hat{i} - \hat{j}$ and $\vec{w} = 5\hat{i} + 9\hat{j} - 4\hat{k}$ are coplanar.

Solution :

$$\vec{u} \cdot (\vec{v} \times \vec{w}) = \begin{vmatrix} 1 & 5 & -2 \\ 3 & -1 & 0 \\ 5 & 9 & -4 \end{vmatrix} = 1 \begin{vmatrix} -1 & 0 \\ 9 & -4 \end{vmatrix} - 5 \begin{vmatrix} 3 & 0 \\ 5 & -4 \end{vmatrix} + (-2) \begin{vmatrix} 3 & -1 \\ 5 & 9 \end{vmatrix} = 4 + 60 - 64 = 0$$

i.e. volume of the parallelepiped is 0 and thus these three vectors are coplanar.

Ex. 3. Find the vector which is orthogonal to the vector $3\hat{i} + 2\hat{j} + 6\hat{k}$ and is co-planar with the vectors $2\hat{i} + \hat{j} + \hat{k}$ and $\hat{i} - \hat{j} + \hat{k}$

Solution : Let, $\vec{a} = 3\hat{i} + 2\hat{j} + 6\hat{k}$, $\vec{b} = 2\hat{i} + \hat{j} + \hat{k}$, $\vec{c} = \hat{i} - \hat{j} + \hat{k}$

Then by definition, a vector orthogonal to \vec{a} and co-planar to \vec{b} and \vec{c} is given by

$$\begin{aligned} \vec{a} \times (\vec{b} \times \vec{c}) &= (\vec{a} \cdot \vec{c})\vec{b} - (\vec{a} \cdot \vec{b})\vec{c} \\ &= 7(2\hat{i} + \hat{j} + \hat{k}) - 14(\hat{i} + \hat{j} + \hat{k}) = 21\hat{j} - 7\hat{k} \end{aligned}$$

Ex. 4. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{a} \cdot \vec{b} = 1$ and $\vec{a} \times \vec{b} = \hat{j} - \hat{k}$ then prove that $\vec{b} = \hat{i}$.

Solution : As $(\vec{a} \times \vec{b}) \times \vec{a} = (\vec{a} \cdot \vec{a})\vec{b} - (\vec{a} \cdot \vec{b})\vec{a}$

$$\vec{a} \cdot \vec{a} = |\vec{a}|^2 = 1+1+1 = 3 \text{ and } \vec{a} \cdot \vec{b} = 1$$

$$\therefore (\hat{j} - \hat{k}) \times (\hat{i} + \hat{j} + \hat{k}) = 3(\vec{b}) - (\hat{i} + \hat{j} + \hat{k})$$

$$\text{i.e. } \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 0 & 1 & -1 \\ 1 & 1 & 1 \end{vmatrix} = 3(\vec{b}) - (\hat{i} + \hat{j} + \hat{k})$$

$$(2\hat{i} - \hat{j} - \hat{k}) + (\hat{i} + \hat{j} + \hat{k}) = 3\vec{b}$$

$$\text{i.e. } \hat{i} = \vec{b}$$

Ex. 5. Prove that : $\vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b}) = \vec{0}$

Solution : $\vec{a} \times (\vec{b} \times \vec{c}) + \vec{b} \times (\vec{c} \times \vec{a}) + \vec{c} \times (\vec{a} \times \vec{b})$

$$[(\vec{a} \cdot \vec{c})\vec{b}(\vec{a} \cdot \vec{b})\vec{c}] + [(\vec{b} \cdot \vec{a})\vec{c}(\vec{b} \cdot \vec{c})\vec{a}] + [(\vec{c} \cdot \vec{b})\vec{a}(\vec{c} \cdot \vec{a})\vec{b}]$$

$$(\vec{a} \cdot \vec{c})\vec{b}(\vec{a} \cdot \vec{b})\vec{c} + (\vec{a} \cdot \vec{b})\vec{c}(\vec{b} \cdot \vec{c})\vec{a} + (\vec{b} \cdot \vec{c})\vec{a}(\vec{a} \cdot \vec{c})\vec{b} = \vec{0}$$

Ex. 6. Show that the points $A(2,-1,0)$, $B(-3,0,4)$, $C(-1,-1,4)$ and $D(0,-5,2)$ are non coplanar.

Solution: Let $\vec{a} = 2\hat{i} - \hat{j}$, $\vec{b} = -3\hat{i} + 4\hat{k}$, $\vec{c} = -\hat{i} - \hat{j} + 4\hat{k}$, $\vec{d} = \hat{i} + \hat{j} + 4\hat{k}$, $\vec{d} = 5\hat{i} + 2\hat{k}$

$$\vec{AB} = \vec{b} - \vec{a} = -5\hat{i} + \hat{j} + 4\hat{k}$$

$$\vec{AC} = \vec{c} - \vec{a} = -3\hat{i} + 4\hat{k}$$

$$\vec{AD} = \vec{d} - \vec{a} = -2\hat{i} - 4\hat{j} + 2\hat{k}$$

$$\begin{aligned} \text{Consider : } (\overline{AB}) \cdot (\overline{AC}) \times (\overline{AD}) &= \begin{vmatrix} -5 & 1 & 4 \\ -3 & 0 & 4 \\ -2 & -4 & 2 \end{vmatrix} \\ &= -5[0 + 16] - 1[-6 + 8] + 4[12] \\ &= -80 - 2 + 48 \\ &= 34 \neq 0 \end{aligned}$$

Therefore, the points A, B, C, D are non-coplanar.

Ex. 7: If $\bar{a}, \bar{b}, \bar{c}$ are non coplanar vectors, then show that the four points $2\bar{a} + \bar{b}, \bar{a} + 2\bar{b} + \bar{c}, 4\bar{a} - 2\bar{b} - \bar{c}$ and $3\bar{a} + 4\bar{b} - 5\bar{c}$ are coplanar.

Solution : Let 4 points be, $P(\bar{p}), Q(\bar{q}), R(\bar{r})$ and $S(\bar{s})$

$$\begin{aligned} \bar{p} &= 2\bar{a} + \bar{b}, \bar{q} = \bar{a} + 2\bar{b} + \bar{c}, \\ \bar{r} &= 4\bar{a} - 2\bar{b} - \bar{c}, \bar{s} = 3\bar{a} + 4\bar{b} - 5\bar{c} \end{aligned}$$

Let us form 3 coinitial vectors

$$\overline{PQ} = \bar{q} - \bar{p} = (\bar{a} + 2\bar{b} + \bar{c}) - (2\bar{a} + \bar{b}) = -\bar{a} + \bar{b} + \bar{c}$$

$$\overline{PR} = \bar{r} - \bar{p} = (4\bar{a} - 2\bar{b} - \bar{c}) - (2\bar{a} + \bar{b}) = 2\bar{a} - 3\bar{b} - \bar{c}$$

$$\overline{PS} = \bar{s} - \bar{p} = (3\bar{a} + 4\bar{b} - 5\bar{c}) - (2\bar{a} + \bar{b}) = \bar{a} + 3\bar{b} - 5\bar{c}$$

If P, Q, R, S , are coplanar then these three vectors $\overline{PQ}, \overline{PR}$ and \overline{PS} are also coplanar this is possible only if $\overline{PQ} \cdot (\overline{PR} \times \overline{PS}) = 0$

$$\overline{PQ} \cdot (\overline{PR} \times \overline{PS}) = \begin{vmatrix} -1 & 1 & 1 \\ 2 & -3 & -1 \\ 1 & 3 & -5 \end{vmatrix} = -1(15+3) - 1(-10+1) + 1(6+3) = -18 + 9 + 9 = 0$$



Exercise 5.5

- Find $\bar{a} \cdot (\bar{b} \times \bar{c})$, if $\bar{a} = 3\hat{i} - \hat{j} + 4\hat{k}, \bar{b} = 2\hat{i} + 3\hat{j} - \hat{k}$ and $\bar{c} = -5\hat{i} + 2\hat{j} + 3\hat{k}$
- If the vectors $3\hat{i} + 5\hat{k}, 4\hat{i} + 2\hat{j} - 3\hat{k}$ and $3\hat{i} + \hat{j} + 4\hat{k}$ are to co-terminus edges of the parallelo piped, then find the volume of the parallelopiped.
- If the vectors $-3\hat{i} + 4\hat{j} - 2\hat{k}, \hat{i} + 2\hat{k}$ and $\hat{i} - p\hat{j}$ are coplanar, then find the value of p .

4. Prove that :

$$(i) [\bar{a} \bar{b} + \bar{c} \bar{a} + \bar{b} + \bar{c}] = 0 \quad (ii) (\bar{a} + 2\bar{b} - \bar{c}) [(\bar{a} - \bar{b}) \times \bar{a} - \bar{b} - \bar{c}] = 3[\bar{a} - \bar{b} - \bar{c}]$$

5. If, $\bar{c} = 3\bar{a} - 2\bar{b}$. then prove that $[\bar{a} \bar{b} \bar{c}] = 0$

6. If $u = \hat{i} - 2\hat{j} + \hat{k}$, $\bar{r} = 3\hat{i} + \hat{k}$ and $w = \hat{j}$, \hat{k} are given vectors, then find

$$(i) [\bar{u} + \bar{w}] \cdot [(\bar{w} \times \bar{r}) \times (\bar{r} \times \bar{w})]$$

7. Find the volume of a tetrahedron whose vertices are $A(-1, 2, 3)$, $B(3, -2, 1)$, $C(2, 1, 3)$ and $D(-1, -2, 4)$.

8. If $\bar{a} = \hat{i} + 2\hat{j} + 3$, $\bar{b} = 3\hat{i} + 2\hat{j} +$ and $c = 2\hat{i} + \hat{j} + 3$ then verify that

$$\bar{a} \times (\bar{b} \times \bar{c}) = (\bar{a} \cdot \bar{c})\bar{b} - (\bar{a} \cdot \bar{b})\bar{c}$$

9. If, $\bar{a} = \hat{i} - 2\hat{j}$, $\bar{b} = \hat{i} + 2\hat{j}$ and $\bar{c} = 2\hat{i} + \hat{j} - 2$ then find

$$(i) \bar{a} \times (\bar{b} \times \bar{c}) \quad (ii) (\bar{a} \times \bar{b}) \times \bar{c} \text{ Are the results same? Justify.}$$

10. Show that $\bar{a} \times (\bar{b} \times \bar{c}) + \bar{b} \times (\bar{c} \times \bar{a}) + \bar{c} \times (\bar{a} \times \bar{b}) = 0$



Let's remember!

- If two vectors \vec{a} and \vec{b} are represented by the two adjacent sides of a parallelogram then the diagonal of the parallelogram represents $\vec{a} + \vec{b}$.
- If two vectors \vec{a} and \vec{b} are represented by the two adjacent sides of a triangle so that the initial point of \vec{b} coincides with the terminal point of \vec{a} , then the vector $\vec{a} + \vec{b}$ is represented by the third side.
- Unit vector along \vec{a} is denoted by \hat{a} and is given by $\hat{a} = \frac{\vec{a}}{|\vec{a}|}$
- The vector \vec{OP} where the origin O as initial point and P terminal point, is the position vector (P, V) of the point P with respect to O. $\vec{OP} = \vec{p}$.
- $\vec{AB} = \vec{b} - \vec{a}$
- Unit vector along positive X-axis, Y-axis and Z-axis denoted by \hat{i} , \hat{j} , \hat{k} respectively.
- If P \equiv (x, y, z) is any point in space and O is the origin then.

$$OP = x\hat{i} + y\hat{j} + z\hat{k} \text{ and } |\vec{OP}| = \sqrt{x^2 + y^2 + z^2}$$

- Two non zero vectors \vec{a} and \vec{b} are said to be collinear if $\vec{a} = k\vec{b}$ ($k \neq 0$)
- Three non zero vectors \vec{a} , \vec{b} and \vec{c} are said to be coplanar if $\vec{a} = m\vec{b} + n\vec{c}$ ($m, n \neq 0$)
- Section formula for the internal division : If R(\vec{r}) divides the line segment joining the points

$$A(\vec{a}) \text{ and } B(\vec{b}) \text{ internally in the ratio } m: n \text{ then } \vec{r} = \frac{m\vec{b} + n\vec{a}}{m + n}$$

- Section formula for the external division.

If R(\vec{r}) is any point on the line AB such that points A(\vec{a}), B(\vec{b}), R(\vec{r}) are collinear (i.e. A-B-R or R-A-B) and $\frac{AR}{BR} = \frac{m}{n}$, and where m, n are scalars then $\vec{r} = \frac{m\vec{b} - n\vec{a}}{m - n}$.

- Mid Point Formula : If M(\vec{m}) is the mid-point of the ine segment joining the points A(\vec{a}) and B(\vec{b}) then $\vec{m} = \frac{(\vec{a} + \vec{b})}{2}$.

- Centroid Formula : If G(\vec{g}) is the centroid of the triangle whose vertices are the point A(\vec{a}), B(\vec{b}), C(\vec{c}) then $(\vec{g}) = \frac{\vec{a} + \vec{b} + \vec{c}}{3}$.

- If H(\vec{h}) is incenter of ΔABC then $\vec{h} = \frac{|\vec{BC}|\vec{a} + |\vec{AC}|\vec{b} + |\vec{AB}|\vec{c}}{|\vec{BC}| + |\vec{AC}| + |\vec{AB}|}$

- If G(\vec{g}) is centroid of tetrahedron whose vertices are A(\vec{a}), B(\vec{b}), C(\vec{c}) and D(\vec{d}) then

$$\bar{g} = \frac{\bar{a} + \bar{b} + \bar{c} + \bar{d}}{4}$$

- The scalar product of two non-zero vectors \bar{a} and \bar{b} denoted by $\bar{a} \cdot \bar{b}$, is given by

$$\bar{a} \cdot \bar{b} = |\bar{a}| |\bar{b}| \cos \theta$$

Where θ is the angle between \bar{a} and \bar{b} , $0 \leq \theta \leq \pi$

- If \bar{a} is perpendicular to \bar{b} then $\bar{a} \cdot \bar{b} = 0$

$$\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1 \quad \hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0$$

- The angle θ between two non-zero vectors \bar{a} and \bar{b} is given by $\cos \theta = \frac{\bar{a} \cdot \bar{b}}{|\bar{a}| |\bar{b}|}$

- If $\bar{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$ and $\bar{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$ then $\bar{a} \cdot \bar{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$

- Scalar projection of a vector \bar{a} on vector \bar{b} is given $\frac{\bar{a} \cdot \bar{b}}{|\bar{b}|}$

- Vector projection of \bar{a} on \bar{b} is given by $\left(\frac{\bar{a} \cdot \bar{b}}{|\bar{b}|} \right) (\hat{b})$

- The vector product or cross product of two non-zero vectors \bar{a} and \bar{b} , denoted by $\bar{a} \times \bar{b}$ is given by $\bar{a} \times \bar{b} = |\bar{a}| |\bar{b}| \sin \theta \hat{n}$

where θ is the angle between \bar{a} and \bar{b} , $0 \leq \theta \leq \pi$ and \hat{n} is a unit vector perpendicular to both \bar{a} and \bar{b} .

- $\bar{a} \times \bar{b} = \bar{0}$ if and only if \bar{a} and \bar{b} are collinear.

- $\bar{a} \times \bar{b} = -\bar{b} \times \bar{a}$

- The angle between two vectors \bar{a} and \bar{b} may be given as $\frac{|\bar{a} \times \bar{b}|}{|\bar{a}| |\bar{b}|}$

- The $\hat{i} \times \hat{j} = \hat{k}$, $\hat{j} \times \hat{k} = \hat{i}$, $\hat{k} \times \hat{i} = \hat{j}$ also

$$\hat{i} \times \hat{i} = \bar{0}, \quad \hat{j} \times \hat{j} = \bar{0}, \quad \hat{k} \times \hat{k} = \bar{0}$$

- For a plane containing two vectors \bar{a} and \bar{b} the two perpendicular directions are given by $\pm (\bar{a} \times \bar{b})$.

- If $\bar{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$ and $\bar{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$ then $\bar{a} \times \bar{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$

- For \bar{a} and \bar{b} represent the adjacent sides of a parallelogram then its area is given by $|\bar{a} \times \bar{b}|$.

- For \vec{a} and \vec{b} represent the adjacent sides of a triangle then its area is given by $\frac{1}{2} |\vec{a} \times \vec{b}|$.
- If \vec{OP} makes angles α, β, γ with coordinate axes, then α, β, γ are known as direction angles $\cos \alpha, \cos \beta, \cos \gamma$ are known as direction cosines.

Thus we have d.c.s of \vec{OP} are $l = \cos \alpha, m = \cos \beta, n = \cos \gamma$. i.e. l, m, n

\therefore Direction cosines of \vec{PO} are $-l, -m, -n$

- If l, m, n are direction cosines of a vector \vec{r} ,

Then (i) $l^2 + m^2 + n^2 = 1$

(ii) $\vec{r} = |\vec{r}|(l\hat{i} + m\hat{j} + n\hat{k})$

(iii) $\hat{r} = l\hat{i} + m\hat{j} + n\hat{k}$

- If l, m, n are direction cosines of a vector \vec{r} and a, b, c are three real numbers such that $\frac{l}{a} = \frac{m}{b} = \frac{n}{c}$ then a, b, c are called as direction ratios of vector \vec{r} and

$$l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}, m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}, n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$$

- Scalar Triple Product (or Box Product): The dot product of \vec{a} and $\vec{b} \times \vec{c}$ is called the scalar triple product of \vec{a}, \vec{b} and \vec{c} . It is denoted by $\vec{a} \cdot (\vec{b} \times \vec{c})$ or $[\vec{a} \vec{b} \vec{c}]$.
- $[\vec{a} \vec{b} \vec{c}] = [\vec{b} \vec{c} \vec{a}] = [\vec{c} \vec{a} \vec{b}]$.
- $[\vec{a} \vec{b} \vec{c}] = -[\vec{b} \vec{a} \vec{c}] = -[\vec{a} \vec{c} \vec{b}] = -[\vec{c} \vec{b} \vec{a}]$
- Scalar Triple product is zero, if at least one of the vectors is a zero vector or any two vectors are collinear or all vectors are coplanar.
- Four points A(\vec{a}), B(\vec{b}), C(\vec{c}) and D(\vec{d}) are coplanar if and only if $\vec{AB} \cdot (\vec{AC} \times \vec{AD}) = 0$
- The volume of the parallelepiped whose coterminus edges are represented by the vectors \vec{a}, \vec{b} and \vec{c} is given by $|\vec{a} \cdot (\vec{b} \times \vec{c})|$.
- The volume of the tetrahedron whose coterminus edges are given by \vec{a}, \vec{b} and \vec{c} is $\frac{1}{6} [\vec{a} \vec{b} \vec{c}]$

Miscellaneous Exercise 5

1) Select the correct option from the given alternatives :

1) If $|\vec{a}| = 2, |\vec{b}| = 3, |\vec{c}| = 4$ then $[\vec{a} + \vec{b} \quad \vec{b} + \vec{c} \quad \vec{c} - \vec{a}]$ is equal to

- A) 24 B) -24 C) 0 D) 48

- 2) If $|\vec{a}| = 3$, $|\vec{b}| = 4$, then the value of λ for which $\vec{a} + \lambda \vec{b}$ is perpendicular to $\vec{a} - \lambda \vec{b}$, is
 A) $\frac{9}{16}$ B) $\frac{3}{4}$ C) $\frac{3}{2}$ D) $\frac{4}{3}$
- 3) If sum of two unit vectors is itself a unit vector, then the magnitude of their difference is
 A) $\sqrt{2}$ B) $\sqrt{3}$ C) 1 D) 2
- 4) If $|\vec{a}| = 3$, $|\vec{b}| = 5$, $|\vec{c}| = 7$ and $\vec{a} + \vec{b} + \vec{c} = 0$, then the angle between \vec{a} and \vec{b} is
 A) $\frac{\pi}{2}$ B) $\frac{\pi}{4}$ C) $\frac{\pi}{4}$ D) $\frac{\pi}{6}$
- 5) The volume of tetrahedron whose vertices are $(1, -6, 10)$, $(-1, -3, 7)$, $(5, -1, \lambda)$ and $(7, -4, 7)$ is 11 cu. units then the value of λ is
 A) 7 B) $\frac{\pi}{3}$ C) 1 D) 5
- 6) If α, β, γ are direction angles of a line and $\alpha = 60^\circ$, $\beta = 45^\circ$, the $\gamma =$
 A) 30° or 90° B) 45° or 60° C) 90° or 30° D) 60° or 120°
- 7) The distance of the point $(3, 4, 5)$ from Y- axis is
 A) 3 B) 5 C) $\sqrt{34}$ D) $\sqrt{41}$
- 8) The line joining the points $(-2, 1, -8)$ and (a, b, c) is parallel to the line whose direction ratios are 6, 2, 3. The value of a, b, c are
 A) 4, 3, -5 B) $1, 2, \frac{-13}{2}$ C) 10, 5, -2 D) 3, 5, 11
- 9) If $\cos \alpha, \cos \beta, \cos \gamma$ are the direction cosines of a line then the value of $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma$ is
 A) 1 B) 2 C) 3 D) 4
- 10) If l, m, n are direction cosines of a line then $l\hat{i} + m\hat{j} + n\hat{k}$ is
 A) null vector B) the unit vector along the line
 C) any vector along the line D) a vector perpendicular to the line
- 11) If $|\vec{a}| = 3$ and $-1 \leq k \leq 2$, then $|k\vec{a}|$ lies in the interval
 A) $[0, 6]$ B) $[-3, 6]$ C) $[3, 6]$ D) $[1, 2]$
- 12) Let α, β, γ be distinct real numbers. The points with position vectors $\alpha\hat{i} + \beta\hat{j} + \gamma\hat{k}$,
 $\beta\hat{i} + \gamma\hat{j} + \alpha\hat{k}$, $\gamma\hat{i} + \alpha\hat{j} + \beta\hat{k}$
 A) are collinear B) form an equilateral triangle
 C) form a scalene triangle D) form a right angled triangle

- 13) Let \vec{p} and \vec{q} be the position vectors of P and Q respectively, with respect to O and $|\vec{p}| = p$, $|\vec{q}| = q$. The points R and S divide PQ internally and externally in the ratio 2 : 3 respectively. If OR and OS are perpendicular then.
- A) $9p^2 = 4q^2$ B) $4p^2 = 9q^2$ C) $9p = 4q$ D) $4p = 9q$
- 14) The 2 vectors $\hat{j} + \hat{k}$ and $3\hat{i} - \hat{j} + 4\hat{k}$ represents the two sides AB and AC, respectively of a ΔABC . The length of the median through A is
- A) $\frac{\sqrt{34}}{2}$ B) $\frac{\sqrt{48}}{2}$ C) $\sqrt{18}$ D) None of these
- 15) If \vec{a} and \vec{b} are unit vectors, then what is the angle between \vec{a} and \vec{b} for $\sqrt{3}\vec{a} - \vec{b}$ to be a unit vector ?
- A) 30° B) 45° C) 60° D) 90°
- 16) If θ be the angle between any two vectors \vec{a} and \vec{b} , then $|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$, when θ is equal to
- A) 0 B) $\frac{\pi}{4}$ C) $\frac{\pi}{2}$ D) π
- 17) The value of $\hat{i} \cdot (\hat{j} \times \hat{k}) + \hat{j} \cdot (\hat{i} \times \hat{k}) + \hat{k} \cdot (\hat{i} \times \hat{j})$
- A) 0 B) -1 C) 1 D) 3
- 18) Let a, b, c be distinct non-negative numbers. If the vectors $a\hat{i} + a\hat{j} + c\hat{k}$, $\hat{i} + \hat{k}$ and $c\hat{i} + c\hat{j} + b\hat{k}$ lie in a plane, then c is
- A) The arithmetic mean of a and b B) The geometric mean of a and b
C) The harmonic man of a and b D) 0
- 19) Let $\vec{a} = \hat{i} - \hat{j}$, $\vec{b} = \hat{j} - \hat{k}$, $\vec{c} = \hat{k} - \hat{i}$. If \vec{d} is a unit vector such that $\vec{a} \cdot \vec{d} = 0 = [\vec{b} \ \vec{c} \ \vec{d}]$, then \vec{d} equals.
- A) $\pm \frac{\hat{i} + \hat{j} - 2\hat{k}}{\sqrt{6}}$ B) $\pm \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$ C) $\pm \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$ D) $\pm \hat{k}$
- 20) If \vec{a} , \vec{b} , \vec{c} are non coplanar unit vectors such that $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{(\vec{b} + \vec{c})}{\sqrt{2}}$ then the angle between \vec{a} and \vec{b} is
- A) $\frac{3\pi}{4}$ B) $\frac{\pi}{4}$ C) $\frac{\pi}{2}$ D) π

II Answer the following :

- 1) ABCD is a trapezium with AB parallel to DC and $DC = 3AB$. M is the mid-point of DC, $\overline{AB} = \vec{p}$ and $\overline{BC} = \vec{q}$. Find in terms of \vec{p} and \vec{q} .
 i) \overline{AM} ii) \overline{BD} iii) \overline{MB} iv) \overline{DA}
- 2) The points A, B and C have position vectors \vec{a} , \vec{b} and \vec{c} respectively. The point P is midpoint of AB. Find in terms of \vec{a} , \vec{b} and \vec{c} the vector \overline{PC}
- 3) In a pentagon ABCDE
 Show that $\overline{AB} + \overline{AE} + \overline{BC} + \overline{DC} + \overline{ED} = 2\overline{AC}$
- 4) If in parallelogram ABCD, diagonal vectors are $\overline{AC} = 2\hat{i} + 3\hat{j} + 4\hat{k}$ and $\overline{BD} = -6\hat{i} + 7\hat{j} - 2\hat{k}$, then find the adjacent side vectors \overline{AB} and \overline{AD}
- 5) If two sides of a triangle are $\hat{i} + 2\hat{j}$ and $\hat{i} + \hat{k}$, then find the length of the third side.
- 6) If $|\vec{a}| = |\vec{b}| = 1$, $\vec{a} \cdot \vec{b} = 0$ and $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ then find $|\vec{c}|$
- 7) Find the lengths of the sides of the triangle and also determine the type of a triangle.
 i) A(2, -1, 0), B(4, 1, 1), C(4, -5, 4) ii) L(3, -2, -3), M(7, 0, 1), N(1, 2, 1)
- 8) Find the component form of if \vec{a} if
 i) It lies in YZ plane and makes 60° with positive Y-axis and $|\vec{a}| = 4$
 ii) It lies in XZ plane and makes 45° with positive Z-axis and $|\vec{a}| = 10$
- 9) Two sides of a parallelogram are $3\hat{i} + 4\hat{j} - 5\hat{k}$ and $-2\hat{j} + 7\hat{k}$. Find the unit vectors parallel to the diagonals.
- 10) If D, E, F are the mid-points of the sides BC, CA, AB of a triangle ABC, prove that

$$\overline{AD} + \overline{BE} + \overline{CF} = \vec{0}$$




Fig.5.59

- 11) Find the unit vectors that are parallel to the tangent line to the parabola $y = x^2$ at the point (2, 4)
- 12) Express the vector $\hat{i} + 4\hat{j} - 4\hat{k}$ as a linear combination of the vectors $2\hat{i} - \hat{j} + 3\hat{k}$, $\hat{i} - 2\hat{j} + 4\hat{k}$ and $-\hat{i} + 3\hat{j} - 5\hat{k}$
- 13) If $\overline{OA} = \vec{a}$ and $\overline{OB} = \vec{b}$ then show that the vector along the angle bisector of angle AOB is given by
$$\vec{d} = \lambda \left(\frac{\vec{a}}{|\vec{a}|} + \frac{\vec{b}}{|\vec{b}|} \right)$$

- 14) The position vectors of three consecutive vertices of a parallelogram are $\hat{i} + \hat{j} + \hat{k}$, $\hat{i} + 3\hat{j} + 5\hat{k}$ and $7\hat{i} + 9\hat{j} + 11\hat{k}$. Find the position vector of the fourth vertex.
- 15) A point P with P.V. $\frac{-14\hat{i} + 39\hat{j} + 28\hat{k}}{5}$ divides the line joining A(-1, 6, 5) and B in the ratio 3:2 then find the point B.
- 16) Prove that the sum of the three vectors determined by the medians of a triangle directed from the vertices is zero.
- 17) ABCD is a parallelogram E, F are the mid points of BC and CD respectively. AE, AF meet the diagonal BD at Q and P respectively. Show that P and Q trisect DB.
- 18) If $\triangle ABC$ is a triangle whose orthocenter is P and the circumcenter is Q, then prove that $\overline{PA} + \overline{PC} + \overline{PB} = 2 \overline{PQ}$
- 19) If P is orthocenter, Q is circumcenter and G is centroid of a triangle ABC, then prove that $\overline{QP} = 3 \overline{QG}$
- 20) In a triangle OAB, E is the midpoint of BO and D is a point on AB such that AD: DB = 2:1. If OD and AE intersect at P, determine the ratio OP:PD using vector methods.
- 21) Dot-product of a vector with vectors $3\hat{i} - 5\hat{k}$, $2\hat{i} + 7\hat{j}$ and $\hat{i} + \hat{j} + \hat{k}$ are respectively -1, 6 and 5. Find the vector.
- 22) If \vec{a} , \vec{b} , \vec{c} are unit vectors such that $\vec{a} + \vec{b} + \vec{c} = 0$, then find the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$
- 23) If a parallelogram is constructed on the vectors $\vec{a} = 3\vec{p} - \vec{q}$, $\vec{b} = \vec{p} + 3\vec{q}$ and $|\vec{p}| = |\vec{q}| = 2$ and angle between \vec{p} and \vec{q} is $\pi/3$ show that the ratio of the lengths of the sides is $\sqrt{7} : \sqrt{13}$
- 24) Express the vector $\vec{a} = 5\hat{i} - 2\hat{j} + 5\hat{k}$ as a sum of two vectors such that one is parallel to the vector $\vec{b} = 3\hat{i} + \hat{k}$ and other is perpendicular to \vec{b} .
- 25) Find two unit vectors each of which makes equal angles with \vec{u} , \vec{v} and \vec{w} .
 $\vec{u} = 2\hat{i} + \hat{j} - 2\hat{k}$, $\vec{v} = \hat{i} + 2\hat{j} - 2\hat{k}$ and $\vec{w} = 2\hat{i} - 2\hat{j} + \hat{k}$
- 26) Find the acute angles between the curves at their points of intersection. $y = x^2$, $y = x^3$
- 27) Find the direction cosines and direction angles of the vector.
 i) $2\hat{i} + \hat{j} + 2\hat{k}$ (ii) $(1/2)\hat{i} + \hat{j} + \hat{k}$
- 28) Let $\vec{b} = 4\hat{i} + 3\hat{j}$ and \vec{c} be two vectors perpendicular to each other in the XY-plane. Find vectors in the same plane having projection 1 and 2 along \vec{b} and \vec{c} , respectively, are given by.

- 29) Show that no line in space can make angle $\pi/6$ and $\pi/4$ with X- axis and Y-axis.
- 30) Find the angle between the lines whose direction cosines are given by the equation $6mn-2nl+5lm=0, 3l+m+5n=0$
- 31) If Q is the foot of the perpendicular from P(2,4,3) on the line joining the points A(1,2,4) and B(3,4,5), find coordinates of Q.
- 32) Show that the area of a triangle ABC, the position vectors of whose vertices are a, b and c is $\frac{1}{2}[\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}]$
- 33) Find a unit vector perpendicular to the plane containing the point (a, 0, 0), (0, b, 0), and (0, 0, c). What is the area of the triangle with these vertices?
- 34) State whether each expression is meaningful. If not, explain why? If so, state whether it is a vector or a scalar.

- | | |
|--|---|
| (a) $\vec{a} \cdot (\vec{b} \times \vec{c})$ | (b) $\vec{a} \times (\vec{b} \cdot \vec{c})$ |
| (c) $\vec{a} \times (\vec{b} \times \vec{c})$ | (d) $\vec{a} \cdot (\vec{b} \cdot \vec{c})$ |
| (e) $(\vec{a} \cdot \vec{b}) \times (\vec{c} \cdot \vec{d})$ | (f) $(\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d})$ |
| (g) $(\vec{a} \cdot \vec{b}) \cdot \vec{c}$ | (h) $(\vec{a} \cdot \vec{b}) \vec{c}$ |
| (i) $(\vec{a})(\vec{b} \cdot \vec{c})$ | (j) $\vec{a} \cdot (\vec{b} + \vec{c})$ |
| (k) $\vec{a} \cdot \vec{b} + \vec{c}$ | (l) $ \vec{a} \cdot (\vec{b} + \vec{c})$ |

35. Show that, for any vectors $\vec{a}, \vec{b}, \vec{c}$
 $(\vec{a} + \vec{b} + \vec{c}) \times \vec{c} + (\vec{a} + \vec{b} + \vec{c}) \times \vec{b} + (\vec{b} + \vec{c}) \times \vec{a} = 2\vec{a} \times \vec{c}$
36. Suppose that $\vec{a} = 0$.
- (a) If $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$ then is $\vec{b} = \vec{c}$?
- (b) If $\vec{a} \times \vec{b} = \vec{a} \times \vec{c}$ then is $\vec{b} = \vec{c}$?
- (c) If $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$ and $\vec{a} \times \vec{b} = \vec{a} \times \vec{c}$ then is $\vec{b} = \vec{c}$?
37. If A(3, 2, -1), B(-2, 2, -3), C(3, 5, -2), D(-2, 5, -4) then (i) verify that the points are the vertices of a parallelogram and (ii) find its area.

38. Let A, B, C, D be any four points in space. Prove that

$$\left| \overline{AB} \times \overline{CD} + \overline{BC} \times \overline{AD} + \overline{CA} \times \overline{BD} \right| = 4 \text{ (area of } \triangle ABC)$$
39. Let $\hat{a}, \hat{b}, \hat{c}$ be unit vectors such that $\hat{a} \cdot \hat{b} = \hat{a} \cdot \hat{c} = 0$ and the angle between \hat{b} and \hat{c} be $\pi/6$.
 Prove that $\hat{a} = \pm 2(\hat{b} \times \hat{c})$
40. Find the value of 'a' so that the volume of parallelepiped formed by $\hat{i} + \hat{j} + \hat{k} + a\hat{k}$ and $a\hat{j} + \hat{k}$ becomes minimum.
41. Find the volume of the parallelepiped spanned by the diagonals of the three faces of a cube of side a that meet at one vertex of the cube.
42. If $\bar{a}, \bar{b}, \bar{c}$ are three non-coplanar vectors, then show that

$$\frac{\bar{a} \cdot (\bar{b} \times \bar{c})}{(\bar{c} \times \bar{a}) \cdot \bar{b}} + \frac{\bar{b} \cdot (\bar{a} \times \bar{c})}{(\bar{c} \times \bar{a}) \cdot \bar{b}} = 0$$
43. Prove that $(\bar{a} \times \bar{b}) \cdot (\bar{c} \times \bar{d}) \left| \begin{array}{cc} \bar{a} \cdot \bar{c} & \bar{b} \cdot \bar{c} \\ \bar{a} \cdot \bar{d} & \bar{b} \cdot \bar{d} \end{array} \right|$
44. Find the volume of a parallelepiped whose coterminus edges are represented by the vector $\hat{j} + \hat{k}$, $\hat{i} + \hat{k}$ and $\hat{i} + \hat{j}$. Also find volume of tetrahedron having these coterminus edges.
45. Using properties of scalar triple product, prove that $[\bar{a} + \bar{b} \quad \bar{b} + \bar{c} \quad \bar{c} + \bar{a}] = 2[\bar{a} \quad \bar{b} \quad \bar{c}]$.
- 46) If four points A(\bar{a}), B(\bar{b}), C(\bar{c}) and D(\bar{d}) are coplanar then show that

$$[\bar{a} \quad \bar{b} \quad \bar{d}] + [\bar{b} \quad \bar{c} \quad \bar{d}] + [\bar{c} \quad \bar{a} \quad \bar{d}] = [\bar{a} \quad \bar{b} \quad \bar{c}]$$
- 47) If \bar{a}, \bar{b} and \bar{c} are three non coplanar vectors, then $(\bar{a} + \bar{b} + \bar{c}) \cdot [(\bar{a} + \bar{b}) \times (\bar{a} + \bar{c})] = -[\bar{a} \quad \bar{b} \quad \bar{c}]$
- 48) If in a tetrahedron, edges in each of the two pairs of opposite edges are perpendicular, then show that the edges in the third pair are also perpendicular.



6

Line and Plane



Let's Study

6.1 Vector and Cartesian equations of a line.

6.1.1 Passing through a point and parallel to a vector.

6.1.2 Passing through two points.

6.2 Distance of a point from a line.**6.3 Skew lines**

6.3.1 Distance between skew lines

6.3.2 Distance between parallel lines.

6.4 Equations of Plane:

6.4.1 Passing through a point and perpendicular to a vector.

6.4.2 Passing through a point and parallel to two vectors.

6.4.3 Passing through three non-collinear points.

6.4.4 In normal form.

6.4.5 Passing through the intersection of two planes.

6.5 Angle between planes:

6.5.1 Angle between two planes.

6.5.2 Angle between a line and a plane.

6.6 Coplanarity of two lines.**6.7 Distance of a point from a plane.**

Let's recall.

A line in space is completely determined by a point on it and its direction. Two points on a line determine the direction of the line. Let us derive equations of lines in different forms and discuss parallel lines.

6.1 Vector and Cartesian equations of a line:

Line in space is a locus. Points on line have position vectors. Position vector of a point determines the position of the point in space. In this topic position vector of a variable point on line will be denoted by \vec{r} .

6.1.1 Equation of a line passing through a given point and parallel to given vector:

Theorem 6.1 :

The vector equation of the line passing through $A(\vec{a})$ and parallel to vector \vec{b} is $\vec{r} = \vec{a} + \lambda \vec{b}$.

Proof:

Let L be the line which passes through $A(\vec{a})$ and parallel to vector \vec{b} .

Let $P(\vec{r})$ be a variable point on the line L .

$\therefore \vec{AP}$ is parallel to \vec{b} .

$\therefore \vec{AP} = \lambda \vec{b}$, where λ is a scalar.

$\therefore \vec{OP} - \vec{OA} = \lambda \vec{b}$

$\therefore \vec{r} - \vec{a} = \lambda \vec{b}$

$\therefore \vec{r} = \vec{a} + \lambda \vec{b}$




Fig. 6.1

is the required vector equation of the line.

Remark: Each real value of λ corresponds to a point on line L and conversely each point on L determines unique value of λ . There is one to one correspondence between points on L and values of λ . Here λ is called a parameter and equation $\vec{r} = \vec{a} + \lambda \vec{b}$ is called the **parametric form of vector equation** of line.

Activity: Write position vectors of any three points on the line $\vec{r} = \vec{a} + \lambda \vec{b}$.

Remark: The equation of line passing through $A(\vec{a})$ and parallel to vector \vec{b} can also be expressed as $(\vec{r} - \vec{a}) \times \vec{b} = \vec{0}$. This equation is called the non-parametric form of vector equation of line.

Theorem 6.2 :

The Cartesian equations of the line passing through $A(x_1, y_1, z_1)$ and having direction ratios a, b, c are $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$.

Proof: Let L be the line which passes through $A(x_1, y_1, z_1)$ and has direction ratios a, b, c .

Let $P(x, y, z)$ be a variable point on the line L other than A .

\therefore Direction ratios of L are $x - x_1, y - y_1, z - z_1$.

But direction ratios of line L are a, b, c .

$\therefore \frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}$ are the required Cartesian equations of the line.

In Cartesian form line cannot be represented by a single equation.

Remark :

- If $\vec{b} = a_1 \hat{i} + b_1 \hat{j} + c_1 \hat{k}$ then a_1, b_1, c_1 are direction ratios of the line and conversely if a_1, b_1, c_1 are direction ratios of a line then $\vec{b} = a_1 \hat{i} + b_1 \hat{j} + c_1 \hat{k}$ is parallel to the line.
- The equations $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c} = \lambda$ are called the symmetric form of Cartesian equations of line.
- The equations $x = x_1 + \lambda a, y = y_1 + \lambda b, z = z_1 + \lambda c$ are called parametric form of the Cartesian equations of line.

- The co-ordinates of variable point P on the line are $(x_1 + \lambda a, y_1 + \lambda b, z_1 + \lambda c)$.
- Corresponding to each real value of λ there is one point on the line and conversely corresponding to each point on the line there is unique real value of λ .
- Whenever we write the equations of line in symmetric form, it is assumed that, none of a, b, c is zero. If atleast one of them is zero then we write the equations of line in parametric form and not in symmetric form.
- In place of direction ratios a, b, c if we take direction cosines l, m, n then the co-ordinates of the variable point P will be $(x_1 + \lambda l, y_1 + \lambda m, z_1 + \lambda n)$, A (x_1, y_1, z_1)

$$\begin{aligned} \therefore AP^2 &= (x_1 + \lambda l - x_1)^2 + (y_1 + \lambda m - y_1)^2 + (z_1 + \lambda n - z_1)^2 \\ &= (\lambda l)^2 + (\lambda m)^2 + (\lambda n)^2 = \lambda^2 \{ (l)^2 + (m)^2 + (n)^2 \} = \lambda^2 \end{aligned}$$

$$AP^2 = \lambda^2 \therefore AP = |\lambda| \text{ and } \lambda = \pm AP$$

Thus parameter of point on a line gives its distance from the base point of the line.

6.1.2 Equation of a line passing through given two points.

Theorem 6.3 : The equation of the line passing through A

(\bar{a}) and B (\bar{b}) is $\bar{r} = \bar{a} + \lambda(\bar{b} - \bar{a})$.

Proof: Let L be the line which passes through A (\bar{a}) and B (\bar{b}) .

Let P (\bar{r}) be a variable point on the line L other than A.

$\therefore \overline{AP}$ and $\lambda \overline{AB}$ are collinear.

$\therefore \overline{AP} = \lambda \overline{AB}$, where λ is a scalar .

$$\therefore \bar{r} - \bar{a} = \lambda(\bar{b} - \bar{a}) \quad \bar{r} = \bar{a} + \lambda(\bar{b} - \bar{a})$$

$\therefore \bar{r} = \bar{a} + \lambda(\bar{b} - \bar{a})$ is the required equation of the line.




Fig. 6.2

Remark: The equation of the line passing through A (\bar{a}) and B (\bar{b}) can also be expressed as

$$(\bar{r} - \bar{a}) \times (\bar{b} - \bar{a}) = \bar{0}.$$

Theorem 6.4 : The Cartesian equations of the line passing through A (x_1, y_1, z_1) and B (x_2, y_2, z_2) is

$$\frac{x - x_1}{x_2 - x_1} = \frac{y - y_1}{y_2 - y_1} = \frac{z - z_1}{z_2 - z_1}$$

Proof: Let L be the line which passes through A (x_1, y_1, z_1) and B (x_2, y_2, z_2)

Let P (x, y, z) be a variable point on the line L other than A.

\therefore Direction ratios of L are $x - x_1, y - y_1, z - z_1$

But direction ratios of line L are $x_2 - x_1, y_2 - y_1, z_2 - z_1$

$\therefore \frac{x - x_1}{x_2 - x_1} = \frac{y - y_1}{y_2 - y_1} = \frac{z - z_1}{z_2 - z_1}$ are the required Cartesian equations of the line.



Solved Examples

Ex.(1) Verify that point having position vector $4\hat{i} - 11\hat{j} + 2\hat{k}$ lies on the line

$$\bar{r} = (6\hat{i} - 4\hat{j} + 5\hat{k}) + \lambda(2\hat{i} + 7\hat{j} + 3\hat{k}).$$

Solution: Replacing \bar{r} by $4\hat{i} - 11\hat{j} + 2\hat{k}$ we get,

$$4\hat{i} - 11\hat{j} + 2\hat{k} = (6\hat{i} - 4\hat{j} + 5\hat{k}) + \lambda(2\hat{i} + 7\hat{j} + 3\hat{k})$$

$$\therefore 6 + 2\lambda = 4, -4 + 7\lambda = -11, 5 + 3\lambda = 2$$

From each of these equations we get the same value of λ .

\therefore Given point lies on the given line.

Alternative Method: Equation $\bar{r} = \bar{a} + \lambda\bar{b}$ can be written as $\bar{r} - \bar{a} = \lambda\bar{b}$

Thus point P(\bar{r}) lies on the line if and only if $\bar{r} - \bar{a}$ is a scalar multiple of \bar{b} .

$$\text{Equation of line is } \bar{r} = (6\hat{i} - 4\hat{j} + 5\hat{k}) + \lambda(2\hat{i} + 7\hat{j} + 3\hat{k}).$$

$$\therefore \bar{a} = 6\hat{i} - 4\hat{j} + 5\hat{k} \text{ and } \bar{b} = 2\hat{i} + 7\hat{j} + 3\hat{k}$$

The position vector of given point is $\bar{r} = 4\hat{i} - 11\hat{j} + 2\hat{k}$

$$\bar{r} - \bar{a} = (4\hat{i} - 11\hat{j} + 2\hat{k}) - (6\hat{i} - 4\hat{j} + 5\hat{k}) = -2\hat{i} - 7\hat{j} - 3\hat{k}$$

$$= -(2\hat{i} + 7\hat{j} + 3\hat{k})$$

$$= -1\bar{b}, \text{ a scalar multiple of } \bar{b}.$$

\therefore Given point lies on the given line.

Ex.(2) Find the vector equation of the line passing through the point having position vector

$$4\hat{i} - \hat{j} + 2\hat{k} \text{ and parallel to vector } -2\hat{i} - \hat{j} + \hat{k}$$

Solution: The equation of the line passing through A(\bar{a}) and parallel to vector \bar{b} is $\bar{r} = \bar{a} + \lambda\bar{b}$.

The equation of the line passing through $4\hat{i} - \hat{j} + 2\hat{k}$ and parallel to vector $-2\hat{i} - \hat{j} + \hat{k}$ is

$$\bar{r} = (4\hat{i} - \hat{j} + 2\hat{k}) + \lambda(-2\hat{i} - \hat{j} + \hat{k}).$$

Ex.(3) Find the vector equation of the line passing through the point having position vector

$$2\hat{i} + \hat{j} - 3\hat{k} \text{ and perpendicular to vectors } \hat{i} + \hat{j} + \hat{k} \text{ and } \hat{i} + 2\hat{j} - \hat{k}$$

Solution:

$$\text{Let } \bar{a} = 2\hat{i} + \hat{j} - 3\hat{k}, \bar{b} = \hat{i} + \hat{j} + \hat{k} \text{ and } \bar{c} = \hat{i} + 2\hat{j} - \hat{k}$$

We know that $\bar{b} \times \bar{c}$ is perpendicular to both \bar{b} and \bar{c} .

$\therefore \bar{b} \times \bar{c}$ is parallel to the required line.

$$\bar{b} \times \bar{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 2 & -1 \end{vmatrix} = -3\hat{i} + 2\hat{j} + \hat{k}$$

Thus required line passes through $\bar{a} = 2\hat{i} + \hat{j} - 3\hat{k}$ and parallel to $-3\hat{i} + 2\hat{j} + \hat{k}$.

\therefore Its equation is $\vec{r} = (2\hat{i} + \hat{j} - 3\hat{k}) + \lambda(-3\hat{i} + 2\hat{j} + \hat{k})$

Ex.(4) Find the vector equation of the line passing through $2\hat{i} + \hat{j} - \hat{k}$ and parallel to the line joining points $-\hat{i} + \hat{j} + 4\hat{k}$ and $\hat{i} + 2\hat{j} + 2\hat{k}$.

Solution :

Let A, B, C be points with position vectors $\vec{a} = 2\hat{i} + \hat{j} - \hat{k}$, $\vec{b} = -\hat{i} + \hat{j} + 4\hat{k}$ and $\vec{c} = \hat{i} + 2\hat{j} + 2\hat{k}$ respectively.

$$\vec{BC} = \vec{c} - \vec{b} = (\hat{i} + 2\hat{j} + 2\hat{k}) - (-\hat{i} + \hat{j} + 4\hat{k}) = 2\hat{i} + \hat{j} - 2\hat{k}$$

The required line passes through $2\hat{i} + \hat{j} - \hat{k}$ and is parallel to $2\hat{i} + \hat{j} - 2\hat{k}$

$$\text{Its equation is } \vec{r} = (2\hat{i} + \hat{j} - \hat{k}) + \lambda(2\hat{i} + \hat{j} - 2\hat{k})$$

Ex.(5) Find the vector equation of the line passing through A(1, 2, 3) and B(2, 3, 4).

Solution: Let position vectors of points A and B be \vec{a} and \vec{b} .

$$\therefore \vec{a} = \hat{i} + 2\hat{j} + 3\hat{k} \text{ and } \vec{b} = 2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$\therefore \vec{b} - \vec{a} = (2\hat{i} + 3\hat{j} + 4\hat{k}) - (\hat{i} + 2\hat{j} + 3\hat{k}) = (\hat{i} + \hat{j} + \hat{k})$$

The equation of the line passing through A (\vec{a}) and B (\vec{b}) is $\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$.

The equation of the required line is $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(\hat{i} + \hat{j} + \hat{k})$

Activity: Verify that position vector of B satisfies the above equation.

Ex.(6) Find the Cartesian equations of the line passing through A(1, 2, 3) and having direction ratios 2, 3, 7.

Solution:

The Cartesian equations of the line passing through A(x_1, y_1, z_1) and having direction ratios

$$a, b, c \text{ are } \frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c}.$$

Here (x_1, y_1, z_1) = (1, 2, 3) and direction ratios are 2, 3, 7.

$$\text{Required equation } \frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{7}.$$

Ex.(7) Find the Cartesian equations of the line passing through A(1, 2, 3) and B(2, 3, 4).

Solution: The Cartesian equations of the line passing through A(x_1, y_1, z_1) and B(x_2, y_2, z_2) are

$$\frac{x-x_1}{x_2-x_1} = \frac{y-y_1}{y_2-y_1} = \frac{z-z_1}{z_2-z_1}$$

Here (x_1, y_1, z_1) = (1, 2, 3) and (x_2, y_2, z_2) = (2, 3, 4).

$$\therefore \text{ Required Cartesian equations are } \frac{x-1}{2-1} = \frac{y-2}{3-2} = \frac{z-3}{4-3}.$$

$$\therefore \frac{x-1}{1} = \frac{y-2}{1} = \frac{z-3}{1}.$$

$$\therefore x-1 = y-2 = z-3.$$

Activity: Verify that co-ordinates of B satisfy the above equation.

Ex.(8) Find the Cartesian equations of the line passing through the point A(2, 1, -3) and perpendicular to vectors $\vec{b} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{c} = \hat{i} + 2\hat{j} - \hat{k}$

Solution: We know that $\vec{b} \times \vec{c}$ is perpendicular to both b and c .

$$\therefore \vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 2 & -1 \end{vmatrix} = -3\hat{i} + 2\hat{j} + \hat{k} \text{ is parallel to the required line.}$$

The direction ratios of the required line are -3, 2, 1 and it passes through A(2, 1, -3).

$$\therefore \text{Its Cartesian equations are } \frac{x-2}{-3} = \frac{y-1}{2} = \frac{z+3}{1}.$$

Ex.(9) Find the angle between lines $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(2\hat{i} - 2\hat{j} + \hat{k})$ and $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(\hat{i} + 2\hat{j} + \hat{k})$

Solution: Let \vec{b} and \vec{c} be vectors along given lines .

$$\therefore \vec{b} = 2\hat{i} - 2\hat{j} + \hat{k} \text{ and } \vec{c} = \hat{i} + 2\hat{j} + \hat{k}$$

Angle between lines is same as the angle between \vec{b} and \vec{c} .

The angle between \vec{b} and \vec{c} is given by,

$$\cos \theta = \frac{\vec{b} \cdot \vec{c}}{|\vec{b}| \cdot |\vec{c}|} = \frac{(2\hat{i} - 2\hat{j} + \hat{k}) \cdot (\hat{i} + 2\hat{j} + \hat{k})}{3 \times 3} = \frac{0}{9} = 0$$

$$\therefore \cos \theta = 0 \quad \therefore \theta = 90^\circ$$

Lines are perpendicular to each other.

Ex.(10) Show that lines $\vec{r} = (-\hat{i} - 3\hat{j} + 4\hat{k}) + \lambda(-10\hat{i} - \hat{j} + \hat{k})$ and

$$\vec{r} = (-10\hat{i} - \hat{j} + \hat{k}) + \mu(-\hat{i} - 3\hat{j} + 4\hat{k}) \text{ intersect each other.}$$

Find the position vector of their point of intersection.

Solution: The position vector of a variable point on the line $\vec{r} = (-\hat{i} - 3\hat{j} + 4\hat{k}) + \lambda(-10\hat{i} - \hat{j} + \hat{k})$ is $(-1 - 10\lambda)\hat{i} + (-3 - \lambda)\hat{j} + (4 + \lambda)\hat{k}$

The position vector of a variable point on the line

$$\vec{r} = (-10\hat{i} - \hat{j} + \hat{k}) + \mu(-\hat{i} - 3\hat{j} + 4\hat{k}) \text{ is } (-10 - 1\mu)\hat{i} + (-1 - 3\mu)\hat{j} + (1 + 4\mu)\hat{k}$$

Given lines intersect each other if there exist some values of λ and μ for which

$$(-1 - 10\lambda)\hat{i} + (-3 - \lambda)\hat{j} + (4 + \lambda)\hat{k} = (-10 - 1\mu)\hat{i} + (-1 - 3\mu)\hat{j} + (1 + 4\mu)\hat{k}$$

$$\therefore -1 - 10\lambda = -10 - 1\mu, -3 - \lambda = -1 - 3\mu \text{ and } 4 + \lambda = 1 + 4\mu$$

$$\therefore 10\lambda - \mu = 9, \lambda - 3\mu = -2 \text{ and } \lambda - 4\mu = -3 \quad \dots \dots \dots (1)$$

Given lines intersect each other if this system is consistent

$$\text{As } \begin{vmatrix} 10 & -1 & 9 \\ 1 & -3 & -2 \\ 1 & -4 & -3 \end{vmatrix} = 10(9 - 8) + 1(-3 + 2) + 9(-4 + 3) = 10 - 1 - 9 = 0$$

∴ The system (1) is consistent and lines intersect each other.

Solving any two equations in system (1), we get. $\lambda = 1, \mu = 1$

Substituting this value of λ in $(-1 - 10\lambda)\hat{i} + (-3 - \lambda)\hat{j} + (4 + \lambda)\hat{k}$ we get, $-11\hat{i} - 4\hat{j} + 5\hat{k}$

∴ The position vector of their point of intersection is $-11\hat{i} - 4\hat{j} + 5\hat{k}$.

Ex.(11) Find the co-ordinates of points on the line $\frac{x+1}{2} = \frac{y-2}{3} = \frac{z+3}{6}$, which are at 3 unit distance from the base point A(-1, 2, -3).

Solution: Let Q ($2\lambda - 1, 3\lambda + 2, 6\lambda - 3$) be a point on the line which is at 3 unit distance from the point A(-1,2,-3) ∴ AQ = 3

$$\therefore \sqrt{(2\lambda)^2 + (3\lambda)^2 + (6\lambda)^2} = 3 \quad \therefore (2\lambda)^2 + (3\lambda)^2 + (6\lambda)^2 = 9 \quad \therefore 49\lambda^2 = 9$$

$$\therefore \lambda = -\frac{3}{7} \text{ or } \frac{3}{7}$$

∴ There are two points on the line which are at a distance of 3 units from P.

Their co-ordinates are Q ($2\lambda - 1, 3\lambda + 2, 6\lambda - 3$)

Hence, the required points are

$$\left(-\frac{1}{7}, 3\frac{2}{7}, -\frac{3}{7}\right) \text{ and } \left(-1\frac{6}{7}, \frac{5}{7}, -5\frac{4}{7}\right).$$



Exercise 6.1

- (1) Find the vector equation of the line passing through the point having position vector $-2\hat{i} + \hat{j} + \hat{k}$ and parallel to vector $4\hat{i} - \hat{j} + 2\hat{k}$.
- (2) Find the vector equation of the line passing through points having position vectors $3\hat{i} + 4\hat{j} - 7\hat{k}$ and $6\hat{i} - \hat{j} + \hat{k}$.
- (3) Find the vector equation of line passing through the point having position vector $5\hat{i} + 4\hat{j} + 3\hat{k}$ and having direction ratios $-3, 4, 2$.
- (4) Find the vector equation of the line passing through the point having position vector $\hat{i} + 2\hat{j} + 3\hat{k}$ and perpendicular to vectors $\hat{i} + \hat{j} + \hat{k}$ and $2\hat{i} - \hat{j} + \hat{k}$.
- (5) Find the vector equation of the line passing through the point having position vector $-\hat{i} - \hat{j} + 2\hat{k}$ and parallel to the line $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(3\hat{i} + 2\hat{j} + \hat{k})$.
- (6) Find the Cartesian equations of the line passing through A(-1, 2, 1) and having direction ratios 2, 3, 1.
- (7) Find the Cartesian equations of the line passing through A(2, 2,1) and B(1, 3, 0).
- (8) A(-2, 3, 4), B(1, 1, 2) and C(4, -1, 0) are three points. Find the Cartesian equations of the line AB and show that points A, B, C are collinear.
- (9) Show that lines $\frac{x+1}{-10} = \frac{y+3}{-1} = \frac{z-4}{1}$ and $\frac{x+10}{-1} = \frac{y+1}{-3} = \frac{z-1}{4}$ intersect each other. Find the co-ordinates of their point of intersection.
- (10) A line passes through (3, -1, 2) and is perpendicular to lines $\vec{r} = (\hat{i} + \hat{j} - \hat{k}) + \lambda(2\hat{i} - 2\hat{j} + \hat{k})$ and $\vec{r} = (2\hat{i} + \hat{j} - 3\hat{k}) + \mu(\hat{i} - 2\hat{j} + 2\hat{k})$. Find its equation.

(11) Show that the line $\frac{x-2}{1} = \frac{y-4}{2} = \frac{z+4}{-2}$ passes through the origin.

6.2 Distance of a point from a line:

Theorem 6.5:

The distance of point $P(\bar{\alpha})$ from the line $\bar{r} = \bar{a} + \lambda \bar{b}$ is

$$\sqrt{|\bar{\alpha} - \bar{a}|^2 - \left[\frac{(\bar{\alpha} - \bar{a}) \cdot \bar{b}}{|\bar{b}|} \right]^2}$$




Fig. 6.3

Proof: The line $\bar{r} = \bar{a} + \lambda \bar{b}$ passes through $A(\bar{a})$.

Let M be the foot of the perpendicular drawn from P to the line

$$\bar{r} = \bar{a} + \lambda \bar{b}$$

$\therefore AM = |\overline{AM}|$ = the projection of \overline{AP} on the line.

= the projection of \overline{AP} on \bar{b} . (As line is parallel to \bar{b})

$$= \frac{\overline{AP} \cdot \bar{b}}{|\bar{b}|}$$

Now ΔAMP is a right angled triangle. $\therefore PM^2 = AP^2 - AM^2$

$$PM^2 = |\overline{AP}|^2 - |\overline{AM}|^2 = |\bar{\alpha} - \bar{a}|^2 - \left[\frac{(\bar{\alpha} - \bar{a}) \cdot \bar{b}}{|\bar{b}|} \right]^2$$

$$\therefore PM = \sqrt{|\bar{\alpha} - \bar{a}|^2 - \left[\frac{(\bar{\alpha} - \bar{a}) \cdot \bar{b}}{|\bar{b}|} \right]^2}$$

The distance of point $P(\bar{\alpha})$ from the line $\bar{r} = (\bar{a} + \lambda \bar{b})$ is $PM = \sqrt{|\bar{\alpha} - \bar{a}|^2 - \left[\frac{(\bar{\alpha} - \bar{a}) \cdot \bar{b}}{|\bar{b}|} \right]^2}$

Ex.(12) : Find the length of the perpendicular drawn from the point $P(3, 2, 1)$ to the line

$$\bar{r} = (7\hat{i} + 7\hat{j} + 6\hat{k}) + \lambda(-2\hat{i} + 2\hat{j} + 3\hat{k})$$

Solution:

The length of the perpendicular is same as the distance of P from the given line.

The distance of point $P(\bar{\alpha})$ from the line $\bar{r} = \bar{a} + \lambda \bar{b}$ is $\sqrt{|\bar{\alpha} - \bar{a}|^2 - \left[\frac{(\bar{\alpha} - \bar{a}) \cdot \bar{b}}{|\bar{b}|} \right]^2}$

Here $\vec{a} = 3\hat{i} + 2\hat{j} + \hat{k}$, $\vec{a}_2 = 7\hat{i} + 7\hat{j} + 6\hat{k}$, $\vec{b} = -2\hat{i} + 2\hat{j} + 3\hat{k}$

$\therefore \vec{a} - \vec{a}_2 = (3\hat{i} + 2\hat{j} + \hat{k}) - (7\hat{i} + 7\hat{j} + 6\hat{k}) = -4\hat{i} - 5\hat{j} - 5\hat{k}$

$|\vec{a} - \vec{a}_2| = \sqrt{(-4)^2 + (-5)^2 + (-5)^2} = \sqrt{16 + 25 + 25} = \sqrt{66}$

$(\vec{a} - \vec{a}_2) \cdot \vec{b} = (-4\hat{i} - 5\hat{j} - 5\hat{k}) \cdot (-2\hat{i} + 2\hat{j} + 3\hat{k}) = 8 - 10 - 15 = -17$

$|\vec{b}| = \sqrt{(-2)^2 + (2)^2 + (3)^2} = \sqrt{17}$

The require length =

$\sqrt{|\vec{a} - \vec{a}_2|^2 - \left[\frac{(\vec{a} - \vec{a}_2) \cdot \vec{b}}{|\vec{b}|} \right]^2} = \sqrt{66 - \left[\frac{-17}{\sqrt{17}} \right]^2} = \sqrt{66 - 17} = \sqrt{49} = 7 \text{ unit}$

Ex.(13) : Find the distance of the point P(0, 2, 3) from the line $\frac{x+3}{5} = \frac{y-1}{2} = \frac{z+4}{3}$.

Solution:

Let M be the foot of the perpendicular drawn from the point P(0, 2, 3) to the given line.

M lies on the line. Let co-ordinates of M be $(5\lambda - 3, 2\lambda + 1, 3\lambda - 4)$.

The direction ratios of PM are $(5\lambda - 3) - 0, (2\lambda + 1) - 2, (3\lambda - 4) - 3$
i.e. $5\lambda - 3, 2\lambda - 1, 3\lambda - 7$

The direction ratios of given line are 5, 2, 3 and

PM is perpendicular to the given line .

$\therefore 5(5\lambda - 3) + 2(2\lambda - 1) + 3(3\lambda - 7) = 0$

$\therefore \lambda = 1$

\therefore The co-ordinates of M are (2,3, -1).




Fig. 6.4

The distance of P from the line is $PM = \sqrt{(2-0)^2 + (3-2)^2 + (-1-3)^2} = \sqrt{21}$ unit

6.3 Skew lines:

If two lines in space intersect at a point then the shortest distance between them is zero. If two lines in space are parallel to each other then the shortest distance between them is the perpendicular distance between them.

A pair of lines in space which neither intersect each other nor are parallel to each other are called skew lines. Skew lines are non-coplanar. Lines in the same plane either intersect or are parallel to each other.

In the figure 6.5, line CP that goes diagonally across the plane CSPR and line SQ passes across the plane SAQP are skew lines. The shortest distance between skew lines is the length of the segment which is perpendicular to both the lines.




Fig. 6.5

6.3.1 Distance between skew lines :

Theorem 6.6: The distance between lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ is $\left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot \vec{b}_1 \times \vec{b}_2}{|\vec{b}_1 \times \vec{b}_2|} \right|$

Proof: Let L_1 and L_2 be the lines whose equations are $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$, $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ respectively. Let PQ be the segment which is perpendicular to both L_1 and L_2 .

To find the length of segment PQ.

Lines L_1 and L_2 pass through points $A(\vec{a}_1)$ and $B(\vec{a}_2)$ respectively.

Lines L_1 and L_2 are parallel to \vec{b}_1 and \vec{b}_2 respectively.

As PQ is perpendicular to both L_1 and L_2 , it is parallel to $\vec{b}_1 \times \vec{b}_2$

The unit vector along \vec{PQ} = unit vector along $\vec{b}_1 \times \vec{b}_2 = \hat{n}$ (say)

PQ = The projection of \vec{AB} on $\vec{PQ} = \vec{AB} \cdot \hat{n}$

$$PQ = \left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right|$$




Fig. 6.6

The shortest distance between lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ is $\left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right|$

Remark

- Two lines intersect each other if and only if the shortest distance between them is zero.
- Lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ intersect each other if and only if $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$

Lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$ and $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ intersect each other if and only if

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Ex.(14) Find the shortest distance between lines $\vec{r} = (2\hat{i} - \hat{j}) + \lambda(2\hat{i} + \hat{j} - 3\hat{k})$ and $\vec{r} = \hat{i} - \hat{j} + 2\hat{k} + \mu(2\hat{i} + \hat{j} - 5\hat{k})$

Solution: The shortest distance between lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ is $\left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right|$

Here $\vec{a}_1 = 2\hat{i} - \hat{j}$, $\vec{a}_2 = \hat{i} - \hat{j} + 2\hat{k}$, $\vec{b}_1 = 2\hat{i} + \hat{j} + 3\hat{k}$, $\vec{b}_2 = 2\hat{i} + \hat{j} - 5\hat{k}$

$$\vec{a}_2 - \vec{a}_1 = (\hat{i} - \hat{j} + 2\hat{k}) - (2\hat{i} - \hat{j}) = -\hat{i} + 2\hat{k}$$

$$\text{And } \bar{b}_1 \times \bar{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -3 \\ 2 & 1 & -5 \end{vmatrix} = -2\hat{i} + 4\hat{j}$$

$$\therefore |\bar{b}_1 \times \bar{b}_2| = \sqrt{4+16} = \sqrt{20} = 2\sqrt{5}$$

$$(\bar{a}_2 - \bar{a}_1) \cdot (\bar{b}_1 \times \bar{b}_2) = (-\hat{i} + 2\hat{k}) \cdot (-2\hat{i} + 4\hat{j}) = 2$$

$$\text{The required shortest distance } \left| \frac{(\bar{a}_2 - \bar{a}_1) \cdot (\bar{b}_1 \times \bar{b}_2)}{|\bar{b}_1 \times \bar{b}_2|} \right| = \left| \frac{2}{2\sqrt{5}} \right| = \frac{1}{\sqrt{5}} \text{ unit.}$$

Ex.(15) Find the shortest distance between lines $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ and $\frac{x-2}{3} = \frac{y-4}{4} = \frac{z-5}{5}$

Solution:

The vector equations of given lines are $\bar{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 4\hat{k})$
and $\bar{r} = (2\hat{i} + 4\hat{j} + 5\hat{k}) + \mu(3\hat{i} + 4\hat{j} + 5\hat{k})$

The shortest distance between lines $\bar{r} = \bar{a}_1 + \lambda\bar{b}_1$ and $\bar{r} = \bar{a}_2 + \lambda\bar{b}_2$ is $\left| \frac{(\bar{a}_2 - \bar{a}_1) \cdot (\bar{b}_1 \times \bar{b}_2)}{|\bar{b}_1 \times \bar{b}_2|} \right|$ and

Here $\bar{a}_1 = \hat{i} + 2\hat{j} + 3\hat{k}$, $\bar{a}_2 = 2\hat{i} + 4\hat{j} + 5\hat{k}$, $\bar{b}_1 = 2\hat{i} + 3\hat{j} + 4\hat{k}$, $\bar{b}_2 = 3\hat{i} + 4\hat{j} + 5\hat{k}$
 $\therefore \bar{a}_2 - \bar{a}_1 = (2\hat{i} + 4\hat{j} + 5\hat{k}) - (\hat{i} + 2\hat{j} + 3\hat{k}) = \hat{i} + 2\hat{j} + 2\hat{k}$

$$\text{And } \bar{b}_1 \times \bar{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{vmatrix} = -\hat{i} + 2\hat{j} - \hat{k}$$

$$|\bar{b}_1 \times \bar{b}_2| = \sqrt{1+4+1} = \sqrt{6}$$

$$(\bar{a}_2 - \bar{a}_1) \cdot (\bar{b}_1 \times \bar{b}_2) = (\hat{i} + 2\hat{j} + 2\hat{k}) \cdot (-\hat{i} + 2\hat{j} - \hat{k}) = 1$$

$$\therefore \text{The required shortest distance } \left| \frac{(\bar{a}_2 - \bar{a}_1) \cdot (\bar{b}_1 \times \bar{b}_2)}{|\bar{b}_1 \times \bar{b}_2|} \right| = \frac{1}{\sqrt{6}} \text{ unit.}$$

Ex.(16) Show that lines :

$$\vec{r} = (\hat{i} + \hat{j} - \hat{k}) + \lambda (2\hat{i} - 2\hat{j} + \hat{k}) \text{ and } \vec{r} = (4\hat{i} - 3\hat{j} + 2\hat{k}) + \mu (\hat{i} - 2\hat{j} + 2\hat{k})$$

intersect each other.

Solution: Lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ intersect each other if and only if

$$(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$$

Here $\vec{a}_1 = \hat{i} + \hat{j} - \hat{k}$, $\vec{a}_2 = 4\hat{i} - 3\hat{j} + 2\hat{k}$, $\vec{b}_1 = 2\hat{i} - 2\hat{j} + \hat{k}$, $\vec{b}_2 = \hat{i} - 2\hat{j} + 2\hat{k}$,
 $\vec{a}_2 - \vec{a}_1 = 3\hat{i} - 4\hat{j} + 3\hat{k}$

$$(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = \begin{vmatrix} 3 & -4 & 3 \\ 2 & -2 & 1 \\ 1 & -2 & 2 \end{vmatrix} = 3(-2) + 4(3) + 3(-2) = -6 + 12 - 6 = 0$$

Given lines intersect each other.

6.3.2 Distance between parallel lines:

Theorem 6.7: The distance between parallel lines $\vec{r} = \vec{a}_1 + \lambda \vec{b}$ and $\vec{r} = \vec{a}_2 + \lambda \vec{b}$ is $|(\vec{a}_2 - \vec{a}_1) \times \hat{b}|$

Proof: Let lines represented by $\vec{r} = \vec{a}_1 + \lambda \vec{b}$ and $\vec{r} = \vec{a}_2 + \lambda \vec{b}$ be l_1 and l_2

Lines L_1 and L_2 pass through $A(\vec{a}_1)$ and

$B(\vec{a}_2)$ respectively.

Let BM be perpendicular to L_1 . To find BM

$\triangle AMB$ is a right angle triangle. Let $m \angle BAM = \theta$

$$\sin \theta = \frac{BM}{AB}$$

$$\therefore BM = AB \sin \theta = AB \cdot 1 \sin \theta = AB \cdot |\hat{b}| \cdot \sin \theta$$

$$|AB \times \hat{b}| = |(\vec{a}_2 - \vec{a}_1) \times \hat{b}|$$

\therefore The distance between parallel lines

$$\vec{r} = \vec{a}_1 + \lambda \vec{b} \text{ and } \vec{r} = \vec{a}_2 + \lambda \vec{b} \text{ is given by}$$

$$d = BM = |(\vec{a}_2 - \vec{a}_1) \times \hat{b}|$$




Fig. 6.7

Ex.(17) Find the distance between parallel lines $\vec{r} = (2\hat{i} - \hat{j} + \hat{k}) + \lambda (2\hat{i} + \hat{j} - 2\hat{k})$ and $\vec{r} = (\hat{i} - \hat{j} + 2\hat{k}) + \mu (2\hat{i} + \hat{j} - 2\hat{k})$

Solution: The distance between parallel lines $\vec{r} = \vec{a}_1 + \lambda \vec{b}$ and $\vec{r} = \vec{a}_2 + \lambda \vec{b}$ given by

$$d = |(\vec{a}_2 - \vec{a}_1) \times \hat{b}|$$

Here $\vec{a}_1 = 2\hat{i} - \hat{j} + \hat{k}$, $\vec{a}_2 = \hat{i} - \hat{j} + 2\hat{k}$, $\vec{b} = 2\hat{i} + \hat{j} - 2\hat{k}$

$$\therefore \bar{a}_2 - \bar{a}_1 = (\hat{i} - \hat{j} + 2\hat{k}) - (2\hat{i} - \hat{j} + \hat{k}) = -\hat{i} + \hat{k} \quad \text{and} \quad \hat{b} = \frac{2\hat{i} + \hat{j} - 2\hat{k}}{3}$$

$$\therefore (\bar{a}_2 - \bar{a}_1) \times \hat{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 0 & 1 \\ \frac{2}{3} & \frac{1}{3} & -\frac{2}{3} \end{vmatrix} = \frac{1}{3} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 0 & 1 \\ 2 & 1 & -2 \end{vmatrix} = \frac{1}{3} \{-\hat{i} - \hat{k}\}$$

$$d = |(\bar{a}_2 - \bar{a}_1) \times \hat{b}| = \frac{\sqrt{2}}{3} \text{ unit}$$

Alternative Method:

The distance between parallel lines $\bar{r} = \bar{a}_1 + \lambda\bar{b}$ and $\bar{r} = \bar{a}_2 + \lambda\bar{b}$ is same as the distance of point A(\bar{a}_1) from the line $\bar{r} = \bar{a}_2 + \lambda\bar{b}$. This distance is given by

$$d = \sqrt{|\bar{a}_2 - \bar{a}_1|^2 - \left[\frac{(\bar{a}_2 - \bar{a}_1) \cdot \bar{b}}{|\bar{b}|} \right]^2}$$

$$\text{Now } \bar{a}_2 - \bar{a}_1 = (\hat{i} - \hat{j} + 2\hat{k}) - (2\hat{i} - \hat{j} + \hat{k}) = -\hat{i} + \hat{k}$$

$$\therefore |\bar{a}_2 - \bar{a}_1| = \sqrt{2}$$

$$\text{As } (\bar{a}_2 - \bar{a}_1) \cdot \bar{b} = (-\hat{i} + \hat{k}) \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = -2 + 0 - 2 = -4$$

$$\bar{b} = 2\hat{i} + \hat{j} - 2\hat{k}, \quad |\bar{b}| = 3$$

$$d = \sqrt{|\bar{a}_1 - \bar{a}_2|^2 - \left[\frac{(\bar{a}_1 - \bar{a}_2) \cdot \bar{b}}{|\bar{b}|} \right]^2} = \sqrt{2 - \left(-\frac{4}{3}\right)^2} = \sqrt{2 - \frac{16}{9}} = \frac{\sqrt{2}}{3} \text{ unit}$$

Ex.(18) Find the distance between parallel lines $\frac{x}{2} = \frac{y}{-1} = \frac{z}{2}$ and $\frac{x-1}{2} = \frac{y-1}{-1} = \frac{z-1}{2}$

Solution:

The vector equations of given lines $\bar{r} = \lambda(2\hat{i} - \hat{j} + 2\hat{k})$

$$\text{and } \bar{r} = (\hat{i} + \hat{j} + \hat{k}) + \mu(2\hat{i} - \hat{j} + 2\hat{k})$$

The distance between parallel lines $\bar{r} = \bar{a}_1 + \lambda\bar{b}$ and $\bar{r} = \bar{a}_2 + \lambda\bar{b}$ is given by $d = |(\bar{a}_2 - \bar{a}_1) \times \hat{b}|$

$$\text{Here } \bar{a}_1 = \bar{0}, \bar{a}_2 = \hat{i} + \hat{j} + \hat{k}, \bar{b} = 2\hat{i} - \hat{j} + 2\hat{k}$$

$$\therefore \bar{a}_2 - \bar{a}_1 = \hat{i} + \hat{j} + \hat{k}$$

$$\text{And } \hat{b} = \frac{2\hat{i} - \hat{j} + 2\hat{k}}{3}$$

$$\therefore (\bar{a}_2 - \bar{a}_1) \times \hat{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ \frac{2}{3} & \frac{-1}{3} & \frac{2}{3} \end{vmatrix} = \frac{1}{3} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 2 & -1 & 2 \end{vmatrix} = \frac{1}{3} \{3\hat{i} - 3\hat{k}\} = \hat{i} - \hat{k}$$

$$d = |(\bar{a}_2 - \bar{a}_1) \times \hat{b}| = \sqrt{2} \text{ unit}$$



Exercise 6.2

- (1) Find the length of the perpendicular from $(2, -3, 1)$ to the line $\frac{x+1}{2} = \frac{y-3}{3} = \frac{z+1}{-1}$
- (2) Find the co-ordinates of the foot of the perpendicular drawn from the point $2\hat{i} - \hat{j} + 5\hat{k}$ to the line $\bar{r} = (11\hat{i} - 2\hat{j} - 8\hat{k}) + \lambda(10\hat{i} - 4\hat{j} - 11\hat{k})$. Also find the length of the perpendicular.
- (3) Find the shortest distance between the lines $\bar{r} = (4\hat{i} - \hat{j}) + \lambda(\hat{i} + 2\hat{j} - 3\hat{k})$ and $\bar{r} = (\hat{i} - \hat{j} + 2\hat{k}) + \mu(\hat{i} + 4\hat{j} - 5\hat{k})$
- (4) Find the shortest distance between the lines $\frac{x+1}{7} = \frac{y+1}{-6} = \frac{z+1}{1}$ and $\frac{x-3}{1} = \frac{y-5}{-2} = \frac{z-7}{1}$
- (5) Find the perpendicular distance of the point $(1, 0, 0)$ from the line $\frac{x-1}{2} = \frac{y+1}{-3} = \frac{z+10}{8}$
Also find the co-ordinates of the foot of the perpendicular.
- (6) $A(1, 0, 4)$, $B(0, -11, 13)$, $C(2, -3, 1)$ are three points and D is the foot of the perpendicular from A to BC . Find the co-ordinates of D .
- (7) By computing the shortest distance, determine whether following lines intersect each other.
 - (i) $\bar{r} = (\hat{i} - \hat{j}) + \lambda(2\hat{i} + \hat{k})$ and $\bar{r} = (2\hat{i} - \hat{j}) + \mu(\hat{i} + \hat{j} - \hat{k})$
 - (ii) $\frac{x-5}{4} = \frac{y-7}{-5} = \frac{z+3}{-5}$ and $\frac{x-8}{7} = \frac{y-7}{1} = \frac{z-5}{3}$
- (8) If lines $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-1}{4}$ and $\frac{x-3}{1} = \frac{y-k}{2} = \frac{z}{1}$ intersect each other then find k .

Miscellaneous Exercise 6 A

- (1) Find the vector equation of the line passing through the point having position vector $3\hat{i} + 4\hat{j} - 7\hat{k}$ and parallel to $6\hat{i} - \hat{j} + \hat{k}$.
- (2) Find the vector equation of the line which passes through the point $(3, 2, 1)$ and is parallel to the vector $2\hat{i} + 2\hat{j} - 3\hat{k}$.

- (3) Find the Cartesian equations of the line which passes through the point $(-2, 4, -5)$ and parallel to the line $\frac{x+2}{3} = \frac{y-3}{5} = \frac{z+5}{6}$.
- (4) Obtain the vector equation of the line $\frac{x+5}{3} = \frac{y+4}{5} = \frac{z+5}{6}$.
- (5) Find the vector equation of the line which passes through the origin and the point $(5, -2, 3)$.
- (6) Find the Cartesian equations of the line which passes through points $(3, -2, -5)$ and $(3, -2, 6)$.
- (7) Find the Cartesian equations of the line passing through $A(3, 2, 1)$ and $B(1, 3, 1)$.
- (8) Find the Cartesian equations of the line passing through the point $A(1, 1, 2)$ and perpendicular to vectors $\vec{b} = \hat{i} + 2\hat{j} + \hat{k}$ and $\vec{c} = 3\hat{i} + 2\hat{j} - \hat{k}$.
- (9) Find the Cartesian equations of the line which passes through the point $(2, 1, 3)$ and perpendicular to lines $\frac{x-1}{1} = \frac{y-2}{2} = \frac{z-3}{3}$ and $\frac{x}{-3} = \frac{y}{2} = \frac{z}{5}$.
- (10) Find the vector equation of the line which passes through the origin and intersect the line $x-1 = y-2 = z-3$ at right angle.
- (11) Find the value of λ so that lines $\frac{1-x}{3} = \frac{7y-14}{2\lambda} = \frac{z-3}{2}$ and $\frac{7-7x}{3\lambda} = \frac{y-5}{1} = \frac{6-z}{5}$ are at right angle.
- (12) Find the acute angle between lines $\frac{x-1}{1} = \frac{y-2}{-1} = \frac{z-3}{2}$ and $\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-3}{1}$.
- (13) Find the acute angle between lines $x = y, z = 0$ and $x = 0, z = 0$.
- (14) Find the acute angle between lines $x = -y, z = 0$ and $x = 0, z = 0$.
- (15) Find the co-ordinates of the foot of the perpendicular drawn from the point $(0, 2, 3)$ to the line $\frac{x+3}{5} = \frac{y-1}{2} = \frac{z+4}{3}$.
- (16) By computing the shortest distance determine whether following lines intersect each other.
- (i) $\vec{r} = (\hat{i} + \hat{j} - \hat{k}) + \lambda(2\hat{i} - \hat{j} + \hat{k})$ and $\vec{r} = (2\hat{i} + 2\hat{j} - 3\hat{k}) + \mu(\hat{i} + \hat{j} - 2\hat{k})$
- (ii) $\frac{x-5}{4} = \frac{y-7}{5} = \frac{z+3}{5}$ and $x-6 = y-8 = z+2$.
- (17) If lines $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-1}{4}$ and $\frac{x-2}{1} = \frac{y+m}{2} = \frac{z-2}{1}$ intersect each other then find m .
- (18) Find the vector and Cartesian equations of the line passing through the point $(-1, -1, 2)$ and parallel to the line $2x-2 = 3y+1 = 6z-2$.
- (19) Find the direction cosines of the line $\vec{r} = \left(-2\hat{i} + \frac{5}{2}\hat{j} - \hat{k}\right) + \lambda(2\hat{i} + 3\hat{j})$.
- (20) Find the Cartesian equation of the line passing through the origin which is perpendicular to $x-1 = y-2 = z-1$ and intersects the $\frac{x-1}{2} = \frac{y+1}{3} = \frac{z-1}{4}$.
- (21) Write the vector equation of the line whose Cartesian equations are $y = 2$ and $4x - 3z + 5 = 0$.

(22) Find the co-ordinates of points on the line $\frac{x-1}{1} = \frac{y-2}{-2} = \frac{z-3}{2}$ which are at the distance 3 unit from the base point A(1, 2, 3).

6.4 Equations of Plane :

Introduction : A plane is a surface such that the line joining any two points on it lies entirely on it.

Plane can be determined by

- (i) two intersecting lines
- (ii) two parallel lines
- (iii) a line and a point outside it
- (iv) three non collinear points.

Definition: A line perpendicular to a plane is called a normal to the plane. A plane has several normals. They all have the proportional direction ratios. We require only direction ratios of normal therefore we refer normal as **the normal** to a plane.

Direction ratios of the normal to the XY plane are 0, 0, 1.

6.4.1 Equation of plane passing through a point and perpendicular to a vector.

Theorem 6.8: The equation of the plane passing through the point $A(\vec{a})$ and perpendicular to vector \vec{n} is $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$.

Proof: Let $P(\vec{r})$ be any point on the plane.

$\therefore \vec{AP}$ is perpendicular to \vec{n} .

$$\therefore \vec{AP} \cdot \vec{n} = 0$$

$$\therefore (\vec{r} - \vec{a}) \cdot \vec{n} = 0$$

$$\therefore \vec{r} \cdot \vec{n} - \vec{a} \cdot \vec{n} = 0$$

$$\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$$




Fig. 6.8

This is the equation of the plane passing through the point $A(\vec{a})$ and perpendicular to the vector \vec{n} .

Remark :

- Equation $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$ is called the **vector equation of plane in scalar product form**.
- If $\vec{a} \cdot \vec{n} = d$ then equation $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$ takes the form $\vec{r} \cdot \vec{n} = d$.

Cartesian form :

Theorem 6.9 : The equation of the plane passing through the point $A(x_1, y_1, z_1)$ and direction ratios of whose normal are a, b, c is $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$.

Proof: Let $P(x, y, z)$ be any point on the plane.

The direction ratios of AP are $x - x_1, y - y_1, z - z_1$.

The direction ratios of the normal are a, b, c . And AP is perpendicular to the normal.

$\therefore a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$. This is the required equation of plane.

Remark: Above equation may be written as $ax + by + cz + d = 0$

Ex.(1) Find the vector equation of the plane passing through the point having position vector $2\hat{i} + 3\hat{j} + 4\hat{k}$ and perpendicular to the vector $2\hat{i} + \hat{j} - 2\hat{k}$.

Solution: We know that the vector equation of the plane passing through $A(\bar{a})$ and normal to vector \bar{n} is given by $\bar{r} \cdot \bar{n} = \bar{a} \cdot \bar{n}$.

Here $\bar{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$, $\bar{n} = 2\hat{i} + \hat{j} - 2\hat{k}$

$$\bar{a} \cdot \bar{n} = (2\hat{i} + 3\hat{j} + 4\hat{k}) \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 4 + 3 - 8 = -1$$

The vector equation of the plane is $\bar{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = -1$.

Ex.(2) Find the Cartesian equation of the plane passing through $A(1, 2, 3)$ and the direction ratios of whose normal are 3, 2, 5.

Solution: The equation of the plane passing through $A(x_1, y_1, z_1)$ and normal to the line having direction ratios a, b, c is $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$.

Here $(x_1, y_1, z_1) \equiv (1, 2, 3)$ and direction ratios of the normal are 3, 2, 5.

The Cartesian equation of the plane is $3(x - 1) + 2(y - 2) + 5(z - 3) = 0$.

$\therefore 3x + 2y + 5z - 22 = 0$.

Ex.(3) The foot of the perpendicular drawn from the origin to a plane is $M(2, 1, -2)$. Find vector equation of the plane.

Solution: OM is normal to the plane.

\therefore The direction ratios of the normal are 2, 1, -2.

The plane passes through the point having position vector $2\hat{i} + \hat{j} - 2\hat{k}$ and vector $\overline{OM} = 2\hat{i} + \hat{j} - 2\hat{k}$ is normal to it.

Its vector equation is $\bar{r} \cdot \bar{n} = \bar{a} \cdot \bar{n}$.

$$\vec{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = (2\hat{i} + \hat{j} - 2\hat{k}) \cdot (2\hat{i} + \hat{j} - 2\hat{k})$$

$$\vec{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 9.$$

6.4.2 The vector equation of the plane passing through point $A(\vec{a})$ and parallel to \vec{b} and \vec{c} :

Theorem 6.10 : The vector equation of the plane passing through the point $A(\vec{a})$ and parallel to non-zero and non-parallel vectors \vec{b} and \vec{c} is $(\vec{r} - \vec{a}) \cdot (\vec{b} \times \vec{c}) = 0$.

Proof: As vectors \vec{b} and \vec{c} are parallel to the plane, vector $\vec{b} \times \vec{c}$ is normal to the plane. Plane passes through $A(\vec{a})$.

Let $P(\vec{r})$ be any point on the plane.

$\therefore \overline{AP}$ is perpendicular to $\vec{b} \times \vec{c}$

$$\therefore \overline{AP} \cdot (\vec{b} \times \vec{c}) = 0$$

$$\therefore (\vec{r} - \vec{a}) \cdot (\vec{b} \times \vec{c}) = 0$$

$$\therefore \vec{r} \cdot (\vec{b} \times \vec{c}) = \vec{a} \cdot (\vec{b} \times \vec{c}) \text{ is the required equation.}$$




Fig. 6.9

Remark: As \overline{AP} , \vec{b} and \vec{c} are parallel to the same plane, they are coplanar vectors. Therefore \overline{AP} can be expressed as the linear combination of \vec{b} and \vec{c} . Hence $\overline{AP} = \lambda\vec{b} + \mu\vec{c}$ for some scalars λ and μ .

$$\therefore \vec{r} - \vec{a} = \lambda\vec{b} + \mu\vec{c}$$

$\therefore \vec{r} = \vec{a} + \lambda\vec{b} + \mu\vec{c}$ This equation is called the **vector equation of plane in parametric form.**

Ex(4) Find the vector equation of the plane passing through the point $A(-1, 2, -5)$ and parallel to vectors $4\hat{i} - \hat{j} + 3\hat{k}$ and $\hat{i} + \hat{j} - \hat{k}$.

Solution: The vector equation of the plane passing through point $A(\vec{a})$ and parallel to \vec{b} and \vec{c} is $\vec{r} \cdot (\vec{b} \times \vec{c}) = \vec{a} \cdot (\vec{b} \times \vec{c})$.

Here $\vec{a} = -\hat{i} + 2\hat{j} - 5\hat{k}$, $\vec{b} = 4\hat{i} - \hat{j} + 3\hat{k}$, $\vec{c} = \hat{i} + \hat{j} - \hat{k}$

$$\vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & -1 & 3 \\ 1 & 1 & -1 \end{vmatrix} = -2\hat{i} + 7\hat{j} + 5\hat{k}$$

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = (-\hat{i} + 2\hat{j} - 5\hat{k}) \cdot (-2\hat{i} + 7\hat{j} + 5\hat{k}) = -9$$

The required equation is $\vec{r} \cdot (-2\hat{i} + 7\hat{j} + 5\hat{k}) = -9$

Ex.(5) Find the Cartesian equation of the plane $\vec{r} = (\hat{i} - \hat{j}) + \lambda(\hat{i} + \hat{j} + \hat{k}) + \mu(\hat{i} - 2\hat{j} + 3\hat{k})$

Solution: Given plane is perpendicular to vector \vec{n} , where

$$\vec{n} = \vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & -2 & 3 \end{vmatrix} = 5\hat{i} - 2\hat{j} - 3\hat{k}$$

∴ The direction ratios of the normal are 5, -2, -3.

And plane passes through A(1, -1, 0).

∴ Its Cartesian equation is $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$

$$\therefore 5(x - 1) - 2(y + 1) - 3(z - 0) = 0$$

$$\therefore 5x - 2y - 3z - 7 = 0$$

6.4.3 The vector equation of plane passing through three non-collinear points :

Theorem 6.11 : The vector equation of the plane passing through non-collinear points A(\vec{a}), B(\vec{b}) and C(\vec{c}) is $(\vec{r} - \vec{a}) \cdot (\vec{b} - \vec{a}) \times (\vec{c} - \vec{a}) = 0$

Proof: Let P(\vec{r}) be any point on the plane passing through non-collinear points A(\vec{a}), B(\vec{b}) and C(\vec{c}).

∴ \vec{AP} , \vec{AB} and \vec{AC} are coplanar .

$$\therefore \vec{AP} \cdot \vec{AB} \times \vec{AC} = 0$$

$$\therefore (\vec{r} - \vec{a}) \cdot \vec{AB} \times \vec{AC} = 0$$

$$\therefore (\vec{r} - \vec{a}) \cdot (\vec{b} - \vec{a}) \times (\vec{c} - \vec{a}) = 0$$

$$\therefore \left[\vec{r} - \vec{a} \quad \vec{b} - \vec{a} \quad \vec{c} - \vec{a} \right] = 0$$

This is the required equation of plane.

Cartesian form of the above equation is

$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0$$

Ex.(6) Find the vector equation of the plane passing through points A (1, 1, 2), B (0, 2, 3) and C (4, 5, 6).




Fig. 6.10

Solution; Let \bar{a} , \bar{b} and \bar{c} be position vectors of points A, B and C respectively .

$$\therefore \bar{a} = \hat{i} + \hat{j} + 2\hat{k}, \bar{b} = 2\hat{j} + 3\hat{k}, \bar{c} = 4\hat{i} + 5\hat{j} + 6\hat{k}$$

$$\therefore \bar{b} - \bar{a} = -\hat{i} + \hat{j} + \hat{k} \text{ and } \bar{c} - \bar{a} = 3\hat{i} + 4\hat{j} + 4\hat{k}$$

$$\therefore (\bar{b} - \bar{a}) \times (\bar{c} - \bar{a}) = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & 1 \\ 3 & 4 & 4 \end{vmatrix} = 7\hat{j} - 7\hat{k}$$

The plane passes through $\bar{a} = \hat{i} + \hat{j} + 2\hat{k}$ and $7\hat{j} - 7\hat{k}$ is normal to the plane.

$$\therefore \text{Its equation is } (\bar{r} - \bar{a}) \cdot (7\hat{j} - 7\hat{k}) = 0$$

$$\therefore (\bar{r} - (\hat{i} + \hat{j} + 2\hat{k})) \cdot (7\hat{j} - 7\hat{k}) = 0$$

$$\therefore \bar{r} \cdot (7\hat{j} - 7\hat{k}) = (\hat{i} + \hat{j} + 2\hat{k}) \cdot (7\hat{j} - 7\hat{k})$$

$$\therefore \bar{r} \cdot (7\hat{j} - 7\hat{k}) = -7 \text{ is the required equation.}$$

6.4.4 The normal form of equation of plane.

Theorem 6.12: The equation of the plane at distance p unit from the origin and to which unit vector \hat{n} is normal is $\bar{r} \cdot \hat{n} = p$

Proof: Let ON be the perpendicular to the plane $\therefore ON = p$

As \hat{n} is the unit vector along ON, $\overline{ON} = p\hat{n}$

Let P(\bar{r}) be any point on the plane.

$$\therefore \overline{NP} \perp \hat{n} \quad \therefore \overline{NP} \cdot \hat{n} = 0$$

$$\therefore (\bar{r} - p\hat{n}) \cdot \hat{n} = 0 \quad \therefore \bar{r} \cdot \hat{n} - p\hat{n} \cdot \hat{n} = 0$$

$$\therefore \bar{r} \cdot \hat{n} - p = 0 \quad \therefore \bar{r} \cdot \hat{n} = p$$




Fig. 6.11

This is called the **normal form of vector equation** of plane.

Remark:

- If l, m, n are direction cosines of the normal to a plane then $\hat{n} = l\hat{i} + m\hat{j} + n\hat{k}$.
- If N is the foot of the perpendicular drawn from the origin to the plane and $ON = p$ then the co-ordinates of N are (pl, pm, pn) .
- The equation of the plane is $lx + my + nz = p$. This is the **normal form of the Cartesian equation** of the plane.
- There are two planes at distance p units from origin and having \hat{n} as unit vector along normal, namely $\bar{r} \cdot \hat{n} = \pm p$

Ex.(7) Find the vector equation of the plane which is at a distance of 6 unit from the origin and to which the vector $2\hat{i} - \hat{j} + 2\hat{k}$ is normal.

Solution: Here $p = 6$ and $\vec{n} = 2\hat{i} - \hat{j} + 2\hat{k}$ $\therefore |\vec{n}| = 3$

$$\therefore \hat{n} = \frac{\vec{n}}{|\vec{n}|} = \frac{2\hat{i} - \hat{j} + 2\hat{k}}{3}$$

The required equation is $\therefore \vec{r} \cdot \hat{n} = p$

$$\therefore \vec{r} \cdot \left(\frac{2\hat{i} - \hat{j} + 2\hat{k}}{3} \right) = 6$$

$$\therefore \vec{r} \cdot (2\hat{i} - \hat{j} + 2\hat{k}) = 18$$

Ex.(8) Find the perpendicular distance of the origin from the plane $x - 3y + 4z - 6 = 0$

Solution : First we write the given Cartesian equation in normal form.

i.e. in the form $lx + my + nz = p$

Direction ratios of the normal are 1, -3, 4.

\therefore Direction cosines are $\frac{1}{\sqrt{26}}, \frac{-3}{\sqrt{26}}, \frac{4}{\sqrt{26}}$

Given equation can be written as $\frac{1}{\sqrt{26}}x - \frac{3}{\sqrt{26}}y + \frac{4}{\sqrt{26}}z = \frac{6}{\sqrt{26}}$

\therefore The distance of the origin from the plane is $\frac{6}{\sqrt{26}}$

Ex.(9) Find the coordinates of the foot of the perpendicular drawn from the origin to the plane $2x + y - 2z = 18$

Solution:

The Direction ratios of the normal to the plane $2x + y - 2z = 18$ are 2, 1, -2.

\therefore Direction cosines are $\frac{2}{3}, \frac{1}{3}, -\frac{2}{3}$

The normal form of the given Cartesian equation is $\frac{2}{3}x + \frac{1}{3}y - \frac{2}{3}z = 6$

$\therefore p = 6$

The coordinates of the foot of the perpendicular are $(lp, mp, np) = \left(6\left(\frac{2}{3}\right), 6\left(\frac{1}{3}\right), 6\left(-\frac{2}{3}\right) \right) \equiv (4, 2, -4)$

Ex.(10) Reduce the equation $\vec{r} \cdot (3\hat{i} - 4\hat{j} + 12\hat{k}) = 8$ to the normal form and hence find

- (i) the length of the perpendicular from the origin to the plane
- (ii) direction cosines of the normal.

Solution: Here $\vec{n} = 3\hat{i} - 4\hat{j} + 12\hat{k} \quad \therefore |\vec{n}| = 13$

The required normal form is $\vec{r} \cdot \frac{(3\hat{i} - 4\hat{j} + 12\hat{k})}{13} = \frac{8}{13}$

- (i) the length of the perpendicular from the origin to the plane is $\frac{8}{13}$
- (ii) direction cosines of the normal are $\frac{3}{13}, \frac{-4}{13}, \frac{12}{13}$.

6.4.5 Equation of plane passing through the intersection of two planes :

If planes $(\vec{r} \cdot \vec{n}_1 - d_1) = 0$ and $\vec{r} \cdot \vec{n}_2 - d_2 = 0$ intersect each other, then for every real value of λ equation $\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = (d_1 + \lambda d_2)$ represents a plane passing through the line of their intersection

If planes $a_1x + b_1y + c_1z + d_1 = 0$ and $a_2x + b_2y + c_2z + d_2 = 0$ intersect each other, then for every real value of λ , equation $(a_1x + b_1y + c_1z + d_1) + \lambda(a_2x + b_2y + c_2z + d_2) = 0$ represents a plane passing through the line of their intersection.




Fig. 6.12

Ex.(11) Find the vector equation of the plane passing through the point (1, 0, 2) and the line of intersection of planes $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 8$ and $\vec{r} \cdot (2\hat{i} + 3\hat{j} + 4\hat{k}) = 3$

Solution: The equation of the required plane is of the form equation $\therefore \vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) - (d_1 + \lambda d_2) = 0$

$$\therefore \vec{r} \cdot \left[(\hat{i} + \hat{j} + \hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 4\hat{k}) \right] = 8 + 3\lambda \quad \dots (1)$$

$$\therefore \vec{r} \cdot \left((1+2\lambda)\hat{i} + (1+3\lambda)\hat{j} + (1+4\lambda)\hat{k} \right) = 8 + 3\lambda$$

The plane passes through the point (1, 0, 2).

$$\therefore (\hat{i} + 2\hat{k}) \cdot \left((1+2\lambda)\hat{i} + (1+3\lambda)\hat{j} + (1+4\lambda)\hat{k} \right) = 8 + 3\lambda$$

$$\therefore (1+2\lambda) + 2(1+4\lambda) = 8 + 3\lambda$$

$$\therefore 1 + 2\lambda + 2 + 8\lambda = 8 + 3\lambda$$

$$\therefore 7\lambda = 5$$

$$\therefore \lambda = \frac{5}{7} \quad \dots (2)$$

From (1) and (2) we get

$$\therefore \vec{r} \cdot \left((\hat{i} + \hat{j} + \hat{k}) + \frac{5}{7}(2\hat{i} + 3\hat{j} + 4\hat{k}) \right) = 8 + 3\left(\frac{5}{7}\right)$$

$$\therefore \vec{r} \cdot (17\hat{i} + 22\hat{j} + 27\hat{k}) = 71$$



Exercise 6.3

- (1) Find the vector equation of a plane which is at 42 unit distance from the origin and which is normal to the vector $2\hat{i} + \hat{j} - 2\hat{k}$.
- (2) Find the perpendicular distance of the origin from the plane $6x - 2y + 3z - 7 = 0$.
- (3) Find the coordinates of the foot of the perpendicular drawn from the origin to the plane $2x + 6y - 3z = 63$.
- (4) Reduce the equation $\vec{r} \cdot (3\hat{i} + 4\hat{j} + 12\hat{k}) = 78$ to normal form and hence find
 - (i) the length of the perpendicular from the origin to the plane
 - (ii) direction cosines of the normal.
- (5) Find the vector equation of the plane passing through the point having position vector $\hat{i} + \hat{j} + \hat{k}$ and perpendicular to the vector $4\hat{i} + 5\hat{j} + 6\hat{k}$.
- (6) Find the Cartesian equation of the plane passing through A(-1, 2, 3), the direction ratios of whose normal are 0, 2, 5.
- (7) Find the Cartesian equation of the plane passing through A(7, 8, 6) and parallel to the XY plane.
- (8) The foot of the perpendicular drawn from the origin to a plane is M(1,0,0). Find the vector equation of the plane.
- (9) Find the vector equation of the plane passing through the point A(-2, 7, 5) and parallel to vectors $4\hat{i} - \hat{j} + 3\hat{k}$ and $\hat{i} + \hat{j} + \hat{k}$.
- (10) Find the Cartesian equation of the plane $\vec{r} = (5\hat{i} - 2\hat{j} - 3\hat{k}) + \lambda(\hat{i} + \hat{j} + \hat{k}) + \mu(\hat{i} - 2\hat{j} + 3\hat{k})$.
- (11) Find the vector equation of the plane which makes intercepts 1, 1, 1 on the co-ordinates axes.

6.5 Angle between planes: In this article we will discuss angles between two planes, angle between a line and a plane.

6.5.1 Angle between planes: Angle between planes can be determined from the angle between their normals. Planes $\vec{r} \cdot \vec{n}_1 = d_1$ and $\vec{r} \cdot \vec{n}_2 = d_2$ are perpendicular to each other if and only if $\vec{n}_1 \cdot \vec{n}_2 = 0$
 Planes $a_1x + b_1y + c_1z + d_1 = 0$ and $a_2x + b_2y + c_2z + d_2 = 0$ are perpendicular to each other if and only if $a_1a_2 + b_1b_2 + c_1c_2 = 0$

Definition : If two planes are not perpendicular to each other then **the angle between them** is defined as the **acute angle** between their normals.

The angle between the planes $\vec{r} \cdot \vec{n}_1 = d_1$ and $\vec{r} \cdot \vec{n}_2 = d_2$ is given by $\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| \cdot |\vec{n}_2|}$

Ex.(12) Find the angle between planes $\vec{r} \cdot (\hat{i} + \hat{j} - 2\hat{k}) = 8$ and $\vec{r} \cdot (-2\hat{i} + \hat{j} + \hat{k}) = 3$

Solution: Normal to the given planes are $\vec{n}_1 = \hat{i} + \hat{j} - 2\hat{k}$ and $\vec{n}_2 = -2\hat{i} + \hat{j} + \hat{k}$

The acute angle θ between normal is given by

$$\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| \cdot |\vec{n}_2|}$$

$$\therefore \cos \theta = \frac{|(\hat{i} + \hat{j} - 2\hat{k}) \cdot (-2\hat{i} + \hat{j} + \hat{k})|}{\sqrt{6} \cdot \sqrt{6}} = \frac{|-3|}{6} = \frac{1}{2}$$

$$\therefore \cos \theta = \frac{1}{2} \qquad \therefore \theta = 60^\circ = \frac{\pi}{3}$$

The acute angle between normals \vec{n}_1 and \vec{n}_2 is 60° .

\therefore The angle between given planes is 60° .

6.5.2 Angle between a line and a plane: Line $\vec{r} = \vec{a} + \lambda \vec{b}$ is perpendicular to the plane $\vec{r} \cdot \vec{n} = d$ if and only if \vec{b} and \vec{n} are collinear. i.e. if $\vec{b} = t\vec{n}$ for some $t \in \mathbb{R}$ Line $\vec{r} = \vec{a} + \lambda \vec{b}$ is parallel to the plane of $\vec{r} \cdot \vec{n} = d$ and only if \vec{b} and \vec{n} are perpendicular to each other. i.e. if $\vec{b} \cdot \vec{n} = 0$.

Definition : The angle between a line and a plane is defined as the complementary angle of the acute angle between the normal to the plane and the line.

Because of the definition, the angle between a line and a plane can't be obtuse.

If θ is the angle between the line $\vec{r} = \vec{a} + \lambda \vec{b}$ and the plane $\vec{r} \cdot \vec{n} = d$ then the acute angle between the line and the normal to the plane is $\frac{\pi}{2} - \theta$.

$$\therefore \cos\left(\frac{\pi}{2} - \theta\right) = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

$$\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$$

Ex.(13) Find the angle between the line $\vec{r} = (\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} + \hat{j} + \hat{k})$ and the plane $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 8$

Solution: The angle between the line $\vec{r} = \vec{a} + \lambda\vec{b}$ and the plane $\vec{r} \cdot \vec{n} = d$ is given by $\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$

Here $\vec{b} = \hat{i} + \hat{j} + \hat{k}$ and $\vec{n} = 2\hat{i} - \hat{j} + \hat{k}$

$$\therefore \vec{b} \cdot \vec{n} = (\hat{i} + \hat{j} + \hat{k}) \cdot (2\hat{i} - \hat{j} + \hat{k}) = 2 - 1 + 1 = 2$$

$$|\vec{b}| = \sqrt{1+1+1} = \sqrt{3} \quad \text{and} \quad |\vec{n}| = \sqrt{4+1+1} = \sqrt{6}$$

$$\therefore \sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|} = \frac{2}{\sqrt{3} \cdot \sqrt{6}} = \frac{\sqrt{2}}{3}$$

$$\therefore \theta = \sin^{-1} \left(\frac{\sqrt{2}}{3} \right)$$

6.6 Coplanarity of two lines:

We know that two parallel lines are always coplanar. If two non-parallel lines are coplanar then the shortest distance between them is zero. Conversely if the distance between two non-parallel lines is zero then they are coplanar.

Thus lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ are coplanar if and only if $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$

The plane determined by them passes through $A(\vec{a}_1)$ and $\vec{b}_1 \times \vec{b}_2$. is normal to the plane .

\therefore Its equation is $(\vec{r} - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$.

Lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$ and $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ are coplanar if and only if

$$\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

The equation of the plane determined by them is $\begin{vmatrix} x-x_1 & y-y_1 & z-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$.

Ex.(14) Show that lines $\vec{r} = (\hat{i} + \hat{j} - \hat{k}) + \lambda(2\hat{i} - 2\hat{j} + \hat{k})$ and $\vec{r} = (4\hat{i} - 3\hat{j} + 2\hat{k}) + \mu(\hat{i} - 2\hat{j} + 2\hat{k})$ are coplanar. Find the equation of the plane determined by them.

Solution: Lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ are coplanar if and only if $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$

Here $\vec{a}_1 = \hat{i} + \hat{j} - \hat{k}$, $\vec{a}_2 = 4\hat{i} - 3\hat{j} + 2\hat{k}$

$\vec{b}_1 = 2\hat{i} - 2\hat{j} + \hat{k}$, $\vec{b}_2 = \hat{i} - 2\hat{j} + 2\hat{k}$

$\therefore \vec{a}_2 - \vec{a}_1 = 3\hat{i} - 4\hat{j} + 3\hat{k}$

$$(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = \begin{vmatrix} 3 & -4 & 3 \\ 2 & -2 & 1 \\ 1 & -2 & 2 \end{vmatrix} = 3(-2) + 4(3) + 3(-2) = -6 + 12 - 6 = 0$$

\therefore Given lines are coplanar.

Now $\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -2 & 1 \\ 1 & -2 & 2 \end{vmatrix} = -2\hat{i} - 3\hat{j} - 2\hat{k}$

The equation of the plane determined by them is $(\vec{r} - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$

$\therefore \vec{r} \cdot (\vec{b}_1 \times \vec{b}_2) = \vec{a}_1 \cdot (\vec{b}_1 \times \vec{b}_2)$

$\therefore \vec{r} \cdot (-2\hat{i} - 3\hat{j} - 2\hat{k}) = (\hat{i} + \hat{j} - \hat{k}) \cdot (-2\hat{i} - 3\hat{j} - 2\hat{k})$

$\therefore \vec{r} \cdot (-2\hat{i} - 3\hat{j} - 2\hat{k}) = -3$

$\therefore \vec{r} \cdot (2\hat{i} + 3\hat{j} + 2\hat{k}) = 3$

6.7 Distance of a point from a plane.

To find the distance of the point $A(\vec{a})$ from the plane $\vec{r} \cdot \hat{n} = p$

The distance of the origin from the plane $\vec{r} \cdot \hat{n} = p$ is p .

The equation of the plane passing through $A(\vec{a})$ and parallel to the plane

$\vec{r} \cdot \hat{n} = p$ is $\vec{r} \cdot \hat{n} = \vec{a} \cdot \hat{n}$.

The distance of the origin from the plane $\vec{r} \cdot \hat{n} = \vec{a} \cdot \hat{n}$ is $|\vec{a} \cdot \hat{n}|$




Fig. 6.13




Fig. 6.14

The distance of the point $A(\bar{a})$ from the plane $\bar{r} \cdot \hat{n} = p$ is given by $\left| p - \left| \bar{a} \cdot \hat{n} \right| \right|$.

Remark : For finding distance of a point from a plane, the equation of the plane must be in the normal form.

Ex.(15) Find the distance of the point $4\hat{i} - 3\hat{j} + 2\hat{k}$ from the plane $\bar{r} \cdot (-2\hat{i} + \hat{j} - 2\hat{k}) = 6$

Solution: Here $\bar{a} = 4\hat{i} - 3\hat{j} + 2\hat{k}, \bar{n} = -2\hat{i} + \hat{j} - 2\hat{k}$

$$\therefore |\bar{n}| = \sqrt{(-2)^2 + (1)^2 + (-2)^2} = 3$$

$$\therefore \hat{n} = \frac{(-2\hat{i} + \hat{j} - 2\hat{k})}{3}$$

The normal form of the equation of the given plane is

$$\bar{r} \cdot \frac{(-2\hat{i} + \hat{j} - 2\hat{k})}{3} = 2 \quad \therefore p = 2$$

$$\begin{aligned} \text{Now, } \bar{a} \cdot \hat{n} &= (4\hat{i} - 3\hat{j} + 2\hat{k}) \cdot \frac{(-2\hat{i} + \hat{j} - 2\hat{k})}{3} \\ &= \frac{(4\hat{i} - 3\hat{j} + 2\hat{k}) \cdot (-2\hat{i} + \hat{j} - 2\hat{k})}{3} = \frac{15}{3} = -5 \end{aligned}$$

$$\therefore \left| \bar{a} \cdot \hat{n} \right| = 5$$

The required distance is given by $\left| p - \left| \bar{a} \cdot \hat{n} \right| \right| = |2 - 5| = 3$

Therefore the distance of the point $4\hat{i} - 3\hat{j} + 2\hat{k}$ from the plane $(-2\hat{i} + \hat{j} - 2\hat{k}) = 6$ is 3 unit.



Exercise 6.4

- (1) Find the angle between planes $\bar{r} \cdot (\hat{i} + \hat{j} + 2\hat{k}) = 13$ and $\bar{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 31$.
- (2) Find the acute angle between the line $\bar{r} \cdot (\hat{i} + 2\hat{j} + 2\hat{k}) + \lambda(2\hat{i} + 3\hat{j} - 6\hat{k})$ and the plane $\bar{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 0$.
- (3) Show that lines $\bar{r} = (2\hat{j} - 3\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$ and $\bar{r} = (2\hat{i} + 6\hat{j} + 3\hat{k}) + \mu(2\hat{i} + 3\hat{j} + 4\hat{k})$ are coplanar. Find the equation of the plane determined by them.
- (4) Find the distance of the point $4\hat{i} - 3\hat{j} + \hat{k}$ from the plane $\bar{r} \cdot (2\hat{i} + 3\hat{j} - 6\hat{k}) = 21$.
- (5) Find the distance of the point $(1, 1, -1)$ from the plane $3x + 4y - 12z + 20 = 0$.

Remember This: Line

- The vector equation of the line passing through $A(\vec{a})$ and parallel to vector \vec{b} is $\vec{r} = \vec{a} + \lambda\vec{b}$
- The vector equation of the line passing through $A(\vec{a})$ and $B(\vec{b})$ is $\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$.
- The Cartesian equations of the line passing through $A(x_1, y_1, z_1)$ and having direction ratios a, b, c are $\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}$
- The distance of point $P(\vec{\alpha})$ from the line $\vec{r} = \vec{a} + \lambda\vec{b}$ is given by $\sqrt{|\vec{\alpha} - \vec{a}|^2 - \left[\frac{(\vec{\alpha} - \vec{a}) \cdot \vec{b}}{|\vec{b}|} \right]^2}$
- The shortest distance between lines $\vec{r} = \vec{a}_1 + \lambda_1\vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2\vec{b}_2$ is given by $d = \left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right|$
- Lines $\vec{r} = \vec{a}_1 + \lambda_1\vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2\vec{b}_2$ intersect each other if and only if $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$
- Lines $\frac{x - x_1}{a_1} = \frac{y - y_1}{b_1} = \frac{z - z_1}{c_1}$ and $\frac{x - x_2}{a_2} = \frac{y - y_2}{b_2} = \frac{z - z_2}{c_2}$ intersect each other if and only if $\begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$
- The distance between parallel lines $\vec{r} = \vec{a}_1 + \lambda\vec{b}$ and $\vec{r} = \vec{a}_2 + \lambda\vec{b}$ is given by $d = \left| (\vec{a}_2 - \vec{a}_1) \times \hat{b} \right|$

Plane

- The vector equation of the plane passing through $A(\vec{a})$ and normal to vector \vec{n} is $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$
- Equation $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$ is called the **vector equation of plane in scalar product form**.
- If $\vec{a} \cdot \vec{n} = d$ then equation $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$ takes the form $\vec{r} \cdot \vec{n} = d$
- The equation of the plane passing through $A(x_p, y_p, z_p)$ and normal to the line having direction ratios a, b, c is $a(x - x_p) + b(y - y_p) + c(z - z_p) = 0$.

- The vector equation of the plane passing through point $A(\vec{a})$ and parallel to \vec{b} and \vec{c} is $\vec{r} \cdot (\vec{b} \times \vec{c}) = a \cdot (\vec{b} \times \vec{c})$
- Equation $\vec{r} = \vec{a} + \lambda \vec{b} + \mu \vec{c}$ is called the vector equation of plane in parametric form.
- The vector equation of the plane passing through non-collinear points $A(\vec{a})$, $B(\vec{b})$ and $C(\vec{c})$ is $(\vec{r} - \vec{a}) \cdot (\vec{b} - \vec{a}) \times (\vec{c} - \vec{a}) = 0$

- Cartesian form of the above equation is
$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0$$

- The equation of the plane at distance p unit from the origin and to which unit vector \hat{n} is normal is $\vec{r} \cdot \hat{n} = p$
- If l, m, n are direction cosines of the normal to a plane which is at distance p unit from the origin then its equation is $lx + my + nz = p$.
- If N is the foot of the perpendicular drawn from the origin to a plane and $ON = p$ then the coordinates of N are (pl, pm, pn) .
- If planes $(\vec{r} \cdot \vec{n}_1 - d_1) = 0$ and $\vec{r} \cdot \vec{n}_2 - d_2 = 0$ intersect each other, then for every real value of λ , equation $\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) - (d_1 + \lambda d_2) = 0$ represents a plane passing through the line of their intersection.
- If planes $a_1x + b_1y + c_1z + d_1 = 0$ and $a_2x + b_2y + c_2z + d_2 = 0$ intersect each other, then for every real value of λ , equation $(a_1x + b_1y + c_1z + d_1) + \lambda (a_2x + b_2y + c_2z + d_2) = 0$ represents a plane passing through the line of their intersection.

- The angle between the planes $\vec{r} \cdot \vec{n}_1 = d_1$ and $\vec{r} \cdot \vec{n}_2 = d_2$ is given by $\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| |\vec{n}_2|}$

- The acute angle between the line $\vec{r} = \vec{a} + \lambda \vec{b}$ and the plane $\vec{r} \cdot \vec{n} = d$ is given by $\sin \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| |\vec{n}|}$

- Lines $\vec{r} = \vec{a}_1 + \lambda_1 \vec{b}_1$ and $\vec{r} = \vec{a}_2 + \lambda_2 \vec{b}_2$ are coplanar if and only if $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$ and the equation of the plane determined by them is $(\vec{r} - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$

- Lines $\frac{x - x_1}{a_1} = \frac{y - y_1}{b_1} = \frac{z - z_1}{c_1}$ and $\frac{x - x_2}{a_2} = \frac{y - y_2}{b_2} = \frac{z - z_2}{c_2}$.

are coplanar if and only if
$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$
, and the equation of the plane determined

by them is
$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

- The distance of the point $A(\bar{a})$ from the plane $\bar{r} \cdot \hat{n} = p$ is given by $\left| p - |\bar{a} \cdot \hat{n}| \right|$

Miscellaneous Exercise 6 (B)

I Choose correct alternatives.

- If the line $\frac{x}{3} = \frac{y}{4} = z$ is perpendicular to the line $\frac{x-1}{k} = \frac{y+2}{3} = \frac{z-3}{k-1}$ then the value of k is:
 A) $\frac{11}{4}$ B) $-\frac{11}{4}$ C) $\frac{11}{2}$ D) $\frac{4}{11}$
- The vector equation of line $2x - 1 = 3y + 2 = z - 2$ is
 A) $\bar{r} = \left(\frac{1}{2}\bar{i} - \frac{2}{3}\bar{j} + 2\bar{k} \right) + \lambda(3\bar{i} + 2\bar{j} + 6\bar{k})$
 B) $\bar{r} = \bar{i} - \bar{j} + (2\bar{i} + \bar{j} + \bar{k})$
 C) $\bar{r} = \left(\frac{1}{2}\bar{i} - \bar{j} \right) + \lambda(\bar{i} - 2\bar{j} + 6\bar{k})$
 D) $\bar{r} = (\bar{i} + \bar{j}) + \lambda(\bar{i} - 2\bar{j} + 6\bar{k})$
- The direction ratios of the line which is perpendicular to the two lines $\frac{x-7}{2} = \frac{y+17}{-3} = \frac{z-6}{1}$ and $\frac{x+5}{1} = \frac{y+3}{2} = \frac{z-6}{-2}$ are
 A) 4,5,7 B) 4, -5, 7 C) 4, -5,-7 D) -4, 5, 8
- The length of the perpendicular from (1, 6,3) to the line $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$
 A) 3 B) $\sqrt{11}$ C) $\sqrt{13}$ D) 5

- 5) The shortest distance between the lines $\vec{r} = (\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} - \hat{j} - \hat{k})$ and $\vec{r} = (2\hat{i} - \hat{j} - \hat{k}) + \mu(2\hat{i} + \hat{j} + 2\hat{k})$ is
- A) $\frac{1}{\sqrt{3}}$ B) $\frac{1}{\sqrt{2}}$ C) $\frac{3}{\sqrt{2}}$ D) $\frac{\sqrt{3}}{2}$
- 6) The lines $\frac{x-2}{1} = \frac{y-3}{1} = \frac{z-4}{-k}$ and $\frac{x-1}{k} = \frac{y-4}{2} = \frac{z-5}{1}$ are coplanar if
- A) $k = 1$ or -1 B) $k = 0$ or -3 C) $k = \pm 3$ D) $k = 0$ or -1
- 7) The lines $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$ and $\frac{x-1}{-2} = \frac{y-2}{-4} = \frac{z-3}{6}$ are
- A) perpendicular B) intersecting C) skew D) coincident
- 8) Equation of X-axis is
- A) $x = y = z$ B) $y = z$
 C) $y = 0, z = 0$ D) $x = 0, y = 0$
- 9) The angle between the lines $2x = 3y = -z$ and $6x = -y = -4z$ is
- A) 45° B) 30° C) 0° D) 90°
- 10) The direction ratios of the line $3x + 1 = 6y - 2 = 1 - z$ are
- A) 2, 1, 6 B) 2, 1, -6 C) 2, -1, 6 D) -2, 1, 6
- 11) The perpendicular distance of the plane $2x + 3y - z = k$ from the origin is $\sqrt{14}$ units, the value of k is
- A) 14 B) 196 C) $2\sqrt{14}$ D) $\frac{\sqrt{14}}{2}$
- 12) The angle between the planes $\vec{r} \cdot (\hat{i} - 2\hat{j} + 3\hat{k}) + 4 = 0$ and $\vec{r} \cdot (2\hat{i} + \hat{j} - 3\hat{k}) + 7 = 0$ is
- A) $\frac{\pi}{2}$ B) $\frac{\pi}{3}$ C) $\cos^{-1}\left(\frac{3}{4}\right)$ D) $\cos^{-1}\left(\frac{9}{14}\right)$
- 13) If the planes $\vec{r} \cdot (2\hat{i} - \lambda\hat{j} + \hat{k}) = 3$ and $\vec{r} \cdot (4\hat{i} - \hat{j} + \mu\hat{k}) = 5$ are parallel, then the values of λ and μ are respectively.
- A) $\frac{1}{2}, -2$ B) $-\frac{1}{2}, 2$ C) $-\frac{1}{2}, -2$ D) $\frac{1}{2}, 2$

- 14) The equation of the plane passing through (2, -1, 3) and making equal intercepts on the coordinate axes is
 A) $x + y + z = 1$ B) $x + y + z = 2$ C) $x + y + z = 3$ D) $x + y + z = 4$
- 15) Measure of angle between the planes $5x - 2y + 3z - 7 = 0$ and $15x - 6y + 9z + 5 = 0$ is
 A) 0° B) 30°
 C) 45° D) 90°
- 16) The direction cosines of the normal to the plane $2x - y + 2z = 3$ are
 A) $\frac{2}{3}, \frac{-1}{3}, \frac{2}{3}$ B) $\frac{-2}{3}, \frac{1}{3}, \frac{-2}{3}$
 C) $\frac{2}{3}, \frac{1}{3}, \frac{2}{3}$ D) $\frac{2}{3}, \frac{-1}{3}, \frac{-2}{3}$
- 17) The equation of the plane passing through the points (1, -1, 1), (3, 2, 4) and parallel to Y-axis is :
 A) $3x + 2z - 1 = 0$ B) $3x - 2z = 1$ C) $3x + 2z + 1 = 0$ D) $3x + 2z = 2$
- 18) The equation of the plane in which the line $\frac{x-5}{4} = \frac{y-7}{4} = \frac{z+3}{-5}$ and $\frac{x-8}{7} = \frac{y-4}{1} = \frac{z+5}{3}$ lie, is
 A) $17x - 47y - 24z + 172 = 0$
 B) $17x + 47y - 24z + 172 = 0$
 C) $17x + 47y + 24z + 172 = 0$
 D) $17x - 47y + 24z + 172 = 0$
- 19) If the line $\frac{x+1}{2} = \frac{y-m}{3} = \frac{z-4}{6}$ lies in the plane $3x - 14y + 6z + 49 = 0$, then the value of m is:
 A) 5 B) 3 C) 2 D) -5
- 20) The foot of perpendicular drawn from the point (0,0,0) to the plane is (4, -2, -5) then the equation of the plane is
 A) $4x + y + 5z = 14$ B) $4x - 2y - 5z = 45$
 C) $x - 2y - 5z = 10$ D) $4x + y + 6z = 11$

II. Solve the following :

- Find the vector equation of the plane which is at a distance of 5 unit from the origin and which is normal to the vector $2\hat{i} + \hat{j} + 2\hat{k}$
- Find the perpendicular distance of the origin from the plane $6x + 2y + 3z - 7 = 0$
- Find the coordinates of the foot of the perpendicular drawn from the origin to the plane $2x + 3y + 6z = 49$.
- Reduce the equation $\vec{r} \cdot (6\hat{i} + 8\hat{j} + 24\hat{k}) = 13$ to normal form and hence find
 (i) the length of the perpendicular from the origin to the plane
 (ii) direction cosines of the normal.

- (5) Find the vector equation of the plane passing through the points $A(1, -2, 1)$, $B(2, -1, -3)$ and $C(0, 1, 5)$.
- (6) Find the Cartesian equation of the plane passing through $A(1, -2, 3)$ and the direction ratios of whose normal are $0, 2, 0$.
- (7) Find the Cartesian equation of the plane passing through $A(7, 8, 6)$ and parallel to the plane $\vec{r} \cdot (6\hat{i} + 8\hat{j} + 7\hat{k}) = 0$
- (8) The foot of the perpendicular drawn from the origin to a plane is $M(1, 2, 0)$. Find the vector equation of the plane.
- (9) A plane makes non zero intercepts a, b, c on the co-ordinates axes. Show that the vector equation of the plane is $\vec{r} \cdot (bc\hat{i} + ca\hat{j} + ab\hat{k}) = abc$
- (10) Find the vector equation of the plane passing through the point $A(-2, 3, 5)$ and parallel to vectors $4\hat{i} + 3\hat{k}$ and $\hat{i} + \hat{j}$
- (11) Find the Cartesian equation of the plane $\vec{r} = \lambda(\hat{i} + \hat{j} - \hat{k}) + \mu(\hat{i} + 2\hat{j} + 3\hat{k})$
- (12) Find the vector equations of planes which pass through $A(1, 2, 3)$, $B(3, 2, 1)$ and make equal intercepts on the co-ordinates axes.
- (13) Find the vector equation of the plane which makes equal non-zero intercepts on the co-ordinates axes and passes through $(1, 1, 1)$.
- (14) Find the angle between planes $\vec{r} \cdot (-2\hat{i} + \hat{j} + 2\hat{k}) = 17$ and $\vec{r} \cdot (2\hat{i} + 2\hat{j} + \hat{k}) = 71$.
- (15) Find the acute angle between the line $\vec{r} = \lambda(\hat{i} - \hat{j} + \hat{k})$ and the plane $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) = 23$
- (16) Show that lines $\vec{r} = (\hat{i} + 4\hat{j}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k})$ and $\vec{r} = (3\hat{j} - \hat{k}) + \mu(2\hat{i} + 3\hat{j} + 4\hat{k})$ are coplanar. Find the equation of the plane determined by them.
- (17) Find the distance of the point $3\hat{i} + 3\hat{j} + \hat{k}$ from the plane $\vec{r} \cdot (2\hat{i} + 3\hat{j} + 6\hat{k}) = 21$
- (18) Find the distance of the point $(13, 13, -13)$ from the plane $3x + 4y - 12z = 0$.
- (19) Find the vector equation of the plane passing through the origin and containing the line $\vec{r} = (\hat{i} + 4\hat{j} + \hat{k}) + \lambda(\hat{i} + 2\hat{j} + \hat{k})$
- (20) Find the vector equation of the plane which bisects the segment joining $A(2, 3, 6)$ and $B(4, 3, -2)$ at right angle.
- (21) Show that lines $x = y, z = 0$ and $x + y = 0, z = 0$ intersect each other. Find the vector equation of the plane determined by them.



7

Linear Programming



Let's Study

7.1 Linear Inequations in two variables.

7.1.1 Convex Sets.

7.1.2 Graphical representation of linear inequations in two variables.

7.1.3 Graphical solution of linear inequation.

7.2 Linear Programming Problem (L.P.P.).

7.2.1 Meaning of Linear Programming Problem.

7.2.2 Mathematical formulation of L.P. P.

7.2.3 Solution of L. P. P. by graphical methods.



Let's Recall

A linear equation in two variables $ax + by + c = 0$, where $a, b, c \in \mathbb{R}$ and a and b are not zero simultaneously represents a straight line. A straight line makes disjoint parts of the plane. The points lying on the straight line and two half planes on either side, which are represented by $ax + by + c < 0$ and $ax + by + c > 0$. We will now study the two half planes made by a line.

The sets of points $\{(x, y) \mid ax + by + c < 0\}$ and $\{(x, y) \mid ax + by + c > 0\}$ are two open half planes. The sets of points $\{(x, y) \mid ax + by + c \leq 0\}$ and $\{(x, y) \mid ax + by + c \geq 0\}$ are two half, planes with common points. The sets $\{(x, y) \mid ax + by + c \leq 0\}$ and $\{(x, y) \mid ax + by + c \geq 0\}$ have the common boundary $\{(x, y) \mid ax + by + c = 0\}$.

Linear inequation in two variables.

Definition : A linear inequation in two variables x, y is a mathematical expression of the form $ax + by < c$ or $ax + by > c$ where $a \neq 0, b \neq 0$ simultaneously and $a, b \in \mathbb{R}$.

Activity : Check which of the following ordered pairs is a solution of $2x + 3y - 6 \leq 0$.

1. (1, -1) 2. (2, 1) 3. (-2, 1) 4. (-1, -2) 5. (-3, 4)

Solution :

Sr. No.	(x, y)	Inequation	Conclusion
1.	(1, -1)	$2(1) + 3(-1) - 6 = -1 < 0$	is a solution
2.	(2, 1)	$2(2) + 3(1) - 6 = 1 > 0$	
3.	(-2, 1)		
4.	(1, -2)		
5.	(-3, 4)		

Graphical representation of linear inequation in two variables $ax + by (\leq \text{ or } \geq) c$ is a region on any one side of the straight line $ax + by = c$ in the coordinate system, depending on the sign of inequality.

7.1.1 : Convex set :

Definition : A set of points in a plane is said to be a convex set if line segment joining any two points of the set entirely lies within the set.

The following sets are convex sets :




fig 7.1(a)




fig 7.1(b)

The following sets are not convex sets :




fig 7.1(c)




fig 7.1(d)

Note :

(i) The convex sets may be bounded. Following are bounded convex sets.




fig 7.2(a)




fig 7.2(b)

(ii) Convex sets may be unbounded.




fig 7.3(a)




fig 7.3(b)

Note :

1) Graphical representations of $x \leq h$ and $x \geq h$ on the Cartesian coordinate system. Draw the line $x = h$ in XOY plane.

The solution set is the set of points lying on the Left side or Right side of the line $x = h$.




fig 7.4




fig 7.5

- 2) Graphical representation of $y \leq k$ and $y \geq k$ on the Cartesian coordinate system. Draw the line $y = k$ in XOY plane.

The solution set is the set of points lying below or above the line $y = k$.




fig 7.6




fig 7.7

- 3) Graphical representations of $ax + by \leq 0$ and $ax + by \geq 0$ on the Cartesian coordinate system. The line $ax + by = 0$ passes through the origin, see the following graphs.




fig 7.8




fig 7.9

- 4) To find the solutions of $ax + by \leq c$ and $ax + by \geq c$ graphically, Draw the line $ax + by = c$ in XOY system. It divides the plane into two parts, each part is called half plane :




fig 7.10




fig 7.11

The half plane H_1 , containing origin is called origin side of the line and the other half plane H_2 is non-origin side of the line $ax + by = c$.

For convenience, consider $O(0, 0)$ as test point. If $O(0, 0)$ satisfies the given inequation

$ax + by \leq c$, then the required region is on origin side. Hence shade the region the H_1 otherwise shade the other half plane H_2 .

The shaded portion represents the solution set of the given inequation.

Note that the points $\{(x, y) \mid ax + by = c\}$ form the common boundary of the two half planes.



Solved Examples

Ex.1 Show the solution sets for the following inequations graphically.

- a) $x \leq 3$ b) $y \geq -2$ c) $x + 2y \leq 0$
- d) $2x + 3y \geq 6$ e) $2x - 3y \geq -6$ f) $4x - 5y \leq 20$.

Solution :

- a) To draw : $x \leq 3$
Draw line: $x = 3$




fig 7.12

- b) To draw : $y \geq -2$
Draw line: $y = -2$




fig 7.13

- c) To draw : $x + 2y \leq 0$
Draw line: $x = -2y$

x	0	-2	-4	2
$y = -\frac{x}{2}$	0	1	2	-1




fig 7.14

As the origin lies on the line $x = -2y$, we have to choose another point as a test point. Let us choose $(-2, -2) = -2 - 4 = -6 < 0$
 $x + 2y = -2 - 4 = -6 < 0$

- d) To draw : $2x + 3y \geq 6$
Draw line : $2x + 3y = 6$

x	y	(x, y)
3	0	(3, 0)
0	2	(0, 2)

$2x + 3y|_{(0,0)} = 0 < 6$. Therefore, the required region is the non-origin side of the line.




fig 7.15

- e) To draw : $2x - 3y \geq -6$
Draw line : $2x - 3y = -6$

x	y	(x, y)
-3	0	(-3, 0)
0	2	(0, 2)

$2x + 3y|_{(0,0)} = 0 > -6$. Therefore, the required region is the origin side of the line.




fig 7.16

f) To draw : $4x - 5y \leq 20$
 Draw line : $4x - 5y = 20$

x	y	(x, y)
5	0	(5, 0)
0	-4	(0, -4)




fig 7.17

$4x - 5y|_{(0,0)} = 0 < 20$. Therefore, the required region is the origin side of the line.

Ex. 2 : Represent the solution set of inequation $3x + 2y \leq 6$ graphically.

Solution :

To draw : $3x + 2y \leq 6$
 Draw the line : $3x + 2y = 6$

x	y	(x, y)
2	0	(2, 0)
0	3	(0, 3)




fig 7.18

$3x + 2y|_{(0,0)} = 0 < 6$. Therefore, the required region is the origin side of the line.

Ex. 3 : Find the common region of the solutions of the inequations $x + 2y \geq 4$, $2x - y \leq 6$.

Solution :

To find the common region of : $x + 2y \geq 4$ and $2x - y \leq 6$
 Draw the lines : $x + 2y = 4$ and $2x - y = 6$

Equation of line	x	y	Line passes through (x, y)	Sign	Region
$x + 2y = 4$	4	0	(4, 0)	\geq	Non-origin side
	0	2	(0, 2)		
$2x - y = 6$	3	0	(3, 0)	\leq	Origin side
	0	-6	(0, -6)		




fig 7.19

Notes :

There are two methods of plotting the lines :

- 1) Find the intersection points of the line with X axis and Y axis.
- 2) Write the equation of the straight line in the double intercept form.
For example : consider the line $3x + 2y = 6$.
- 1) The intersection with X axis is given when $y = 0$, So A (2, 0) is the point of intersection with the x axis, The intersection with Y axis is given when $x = 0$.
So B (0, 3) is the point of intersection with the y axis. We draw the line through A and B.
- 2) The equation of line is $3x + 2y = 6$ Divide both sides by 6.
We get the double intercept form $\frac{x}{2} + \frac{y}{3} = 1$
∴ Intercepts on X axis and Y axis are 2 and 3 respectively.
The points (2, 0), (0, 3) lie on the line.

Ex. 4 : Find the graphical solution of $3x + 4y \leq 12$, and $x - 4y \leq 4$

Solution : To find the graphical solution of $3x + 4y \leq 12$ and $x - 4y \leq 4$
Draw the lines $L_1 : 3x + 4y = 12$ and $L_2 : x - 4y = 4$.

Equation of line	x	y	Line passes through (x, y)	Sign	Region
$3x + 4y = 12$	4	0	(4, 0)	\leq	Origin side
	0	3	(0, 3)		
$x - 4y = 4$	4	0	(4, 0)	\leq	Origin side
	0	-1	(0, -1)		

The common shaded region is graphical solution.




fig 7.20

Exercise 7.1

- 1) Solve graphically :
 - i) $x \geq 0$
 - ii) $x \leq 0$
 - iii) $y \geq 0$
 - iv) $y \leq 0$
- 2) Solve graphically :
 - i) $x \geq 0$ and $y \geq 0$
 - ii) $x \leq 0$ and $y \geq 0$
 - iii) $x \leq 0$ and $y \leq 0$
 - iv) $x \geq 0$ and $y \leq 0$
- 3) Solve graphically :
 - i) $2x - 3 \geq 0$
 - ii) $2y - 5 \geq 0$
 - iii) $3x + 4 \leq 0$
 - iv) $5y + 3 \leq 0$
- 4) Solve graphically :
 - i) $x + 2y \leq 6$
 - ii) $2x - 5y \geq 10$
 - iii) $3x + 2y \geq 0$
 - iv) $5x - 3y \leq 0$

5) Solve graphically :

i) $2x + y \geq 2$ and $x - y \leq 1$

ii) $x - y \leq 2$ and $x + 2y \leq 8$

iii) $x + y \geq 6$ and $x + 2y \leq 10$

iv) $2x + 3y \leq 6$ and $x + 4y \geq 4$

v) $2x + y \geq 5$ and $x - y \leq 1$



Solved Examples

Ex. 1 : Find the graphical solution of the system of inequations.

$2x + y = 10, 2x - y = 2, x \geq 0, y \geq 0$

Solution : To find the solution of the system of given inequations -

Draw lines	x	y	Line passes through (x, y)	Sign	Region
L_1 $2x + y = 10$	0	10	(0, 10)	\leq	Origin side of L_1
	5	0	(5, 0)		
L_2 $2x - y = 2$	0	-2	(0, -2)	\geq	Origin side of L_2
	1	0	(1, 0)		

The common shaded region OABCO is the graphical solution. This graphical solution is known as feasible solution.




fig 7.21

Remark : The restrictions $x \geq 0, y \geq 0$, are called non-negativity constraints.

Definition : A solution which satisfies all the constraints is called a feasible solution.

Ex. 2 : Find the feasible solution of the system of inequations $3x + 4y \geq 12, 2x + 5y \geq 10, x \geq 0, y \geq 0$.

To draw (Inequations)	Draw line	x	y	Line passes through (x, y)	Sign	Region
$3x + 4y \geq 12$	L_1 $3x + 4y = 12$	0	3	(0, 3)	\geq	Non-origin side of Line L_1
		4	0	(4, 0)		
$2x + 5y \geq 10$	L_2 $2x + 5y = 10$	0	2	(0, 2)	\geq	Non-origin side of Line L_2
		5	0	(5, 0)		

Solution :

Common shaded region is the feasible solution.

Ex. 3 :

A manufacturer produces two items A and B. Both are processed on two machines I and II. A needs 2 hours on machine I and 2 hours on machine II. B needs 3 hours on machine I and 1 hour on machine II. If machine I can run maximum 12 hours per day and II for 8 hours per day, construct a problem in the form of inequations and find its feasible solution graphically.

Solution :

Let x units of product A and y units of product B be produced.
 $x \geq 0, y \geq 0$.

Tabular form is:

Machine	Product A (x)	Product B (y)	Availability
I	2	3	12
II	2	1	08

Inequations are $2x + 3y \leq 12, 2x + y \leq 8, x \geq 0, y \geq 0$.

To draw graphs of the above inequations :




fig 7.22




fig 7.23

To draw	Draw line	x	y	Line passes through (x, y)	Sign	Region lie on
$2x + 3y \leq 12$	L_1 $2x + 3y = 12$	0	4	(0, 4)	\leq	Origin side of Line L_1
		6	0	(6, 0)		
$2x + y \leq 8$	L_2 $2x + y = 8$	0	8	(0, 8)	\leq	Origin side of Line L_2
		4	0	(4, 0)		

The common shaded region OABCO the feasible region.

Exercise 7.2

1) Find the feasible solution of the following inequations graphically.

- $3x + 2y \leq 18, 2x + y \leq 10, x \geq 0, y \geq 0$
- $2x + 3y \leq 6, x + y \geq 2, x \geq 0, y \geq 0$
- $3x + 4y \geq 12, 4x + 7y \leq 28, y \geq 1, x \geq 0$
- $x + 4y \leq 24, 3x + y \leq 21, x + y \leq 9, x \geq 0, y \geq 0$
- $0 \leq x \leq 3, 0 \leq y \leq 3, x + y \leq 5, 2x + y \geq 4$
- $x - 2y \leq 2, x + y \geq 3, -2x + y \leq 4, x \geq 0, y \geq 0$
- A company produces two types of articles A and B which requires silver and gold. Each unit of A requires 3 gm of silver and 1 gm of gold, while each unit of B requires 2 gm of silver and 2 gm of gold. The company has 6 gm of silver and 4 gm of gold. Construct the inequations and find the feasible solution graphically.
- A furniture dealer deals in tables and chairs. He has Rs.1,50,000 to invest and a space to store at most 60 pieces. A table costs him Rs.1500 and a chair Rs.750. Construct the inequations and find the feasible solution.

7.2 Linear Programming Problems (L.P.P.) :

L.P.P. is an optimization technique used in different fields such as management, planning, production, transportation etc. It is developed during the second world war to optimize the utilization of limited resources to get maximum returns. Linear Programming is used to minimize the cost of production and maximizing the profit. These problems are related to efficient use of limited resources like raw materials, man-power, availability of machine time and cost of the material and so on.

Linear Programming is mathematical technique designed to help managers in the planning and decision making. Programming problems are also known as optimization problems. The mathematical programming involves optimization of a certain function, called objective function, subject to given conditions or restrictions known as constraints.

7.2.1 Meaning of L.P.P. :

Linear implies all the mathematical functions contain variables of index at most one. A L.P.P. may be defined as the problem of maximizing or minimizing a linear function subject to linear constraints. These constraints may be equations or inequations.

Now, we formally define the terms related to L.P.P. as follows :

- 1) **Decision variables** : The variables involved in L.P.P. are called decision variables.
- 2) **Objective function** : A linear function of variables which is to be optimized, i.e. either maximized or minimized, is called an objective function.
- 3) **Constraints** : Conditions under which the objective function is to be optimized are called constraints. These constraints are in the form of equations or inequations.
- 4) **Non-negativity constraints** : In some situations, the values of the variables under considerations may be positive or zero due to the imposed constraints, Such constraints are referred as non-negativity constraints.

7.2.2 Mathematical formulations of L.P.P. :

- Step 1) : Identify the decision variables (x, y) or (x_1, x_2) .
- Step 2) : Identify the objective function and write it as mathematical expression in terms of decision variables.
- Step 3) : Identify the different constraints and express them as mathematical equations / inequations.

- Note** : i) We shall study L.P.P. with at most two variables.
ii) We shall restrict ourselves to L.P.P. involving non-negativity constraints.



Solved examples

Ex. 1 : A Toy manufacturer produces bicycles and tricycles, each of which must be processed through two machine A and B. Machine A has maximum of 120 hours available and machine B has a maximum of 180 hours available. Manufacturing a bicycle requires 4 hours on machine A and 10 hours on machine B. Manufacturing a tricycle required 6 hours on machine A and 3 hours on machine B. If profits are Rs.65 for a bicycle and Rs.45 for a tricycle, formulate L.P.P. to have maximum profit.

Solution :

Let z be the profit, which can be made by manufacturing and selling x tricycles and y bicycles.

$$x \geq 0, y \geq 0$$

$$\therefore \text{Total Profit } z = 45x + 65y$$

$$\text{Maximize } z = 45x + 65y$$

It is given that

Machine	Tricycles (x)	Bicycles (y)	Availability
A	6	4	120
B	3	10	180

From the above table, remaining conditions are $6x + 4y \leq 120$, $3x + 10y \leq 180$.

\therefore The required formulated L.P.P. is as follows :

$$\begin{aligned} &\text{Maximize} && z = 45x + 65y \\ &\text{Subject to the constraints} && x \geq 0, y \geq 0 \\ &&& 6x + 4y \leq 120 \\ &&& 3x + 10y \leq 180 \end{aligned}$$

Ex. 2 : A company manufactures two types of toys A and B. Each toy of type A requires 2 minutes for cutting and 1 minute for assembling. Each toy of type B requires 3 minutes for cutting and 4 minutes for assembling. There are 3 hours available for cutting and 2 hours are available for assembling. On selling a toy of type A the company gets a profit of Rs.10 and that on toy of type B is Rs. 20. Formulate the L.P.P. to maximize profit.

Solution :

Suppose, the company manufactures x toys of type A and y toys of type B.

$$x \geq 0, y \geq 0$$

Let P be the total profit

On selling a toy of type A, company gets Rs.10 and that on a toy of type B is Rs.20.

\therefore total profit on selling x toys of type A and y toys of type B is $p = 10x + 20y$.

\therefore maximize $p = 10x + 20y$.

The conditions are

$$2x + 3y \leq 180, \quad x + 4y \leq 120, \quad x \geq 0, \quad y \geq 0.$$

Ex. 3 : A horticulturist wishes to mix two brands of fertilizers that will provide a minimum of 15 units of potash, 20 units of nitrate and 24 units of phosphate. One unit of brand I provides 3 units of potash, 1 unit of nitrate, 3 units of phosphate. One unit of brand II provides 1 unit of potash, 5 units of nitrate and 2 units of phosphates. One unit of brand I costs Rs. 120 and one unit of brand II costs Rs.60 per unit. Formulate this problems as L.P.P. to minimize the cost.

Solution :

Let z be the cost of mixture prepared by mixing x units of brand I and y units of brand II.

Then $x \geq 0, y \geq 0$.

Since, 1 unit of brand I costs Rs.120.

1 unit of brand II costs Rs.60.

\therefore total cost $z = 120x + 60y$.

\therefore Minimize $z = 120x + 60y$.

Content \ Brand	I per unit	II per unit	Minimum requirement
Potash	3	1	15
Nitrate	1	5	20
Phosphate	3	2	24

The conditions are

$$\begin{aligned} 3x + y &\geq 15, \\ x + 5y &\geq 20, \\ 3x + 2y &\geq 24. \end{aligned}$$

The L.P.P. is

Maximize $z = 120x + 60y$ subject to the above constraints.



Exercise 7.3

- 1) A manufacturing firm produces two types of gadgets A and B, which are first processed in the foundry and then sent to machine shop for finishing. The number of man hours of labour required in each shop for production of A and B and the number of man hours available for the firm are as follows :

Gadgets	Foundry	Machine Shop
A	10	5
B	6	4
Time available (hour)	60	35

Profit on the sale of A is Rs. 30 and B is Rs. 20 per units. Formulate the L.P.P. to have maximum profit.

- 2) In a cattle breeding firm, it is prescribed that the food ration for one animal must contain 14, 22 and 1 units of nutrients A, B and C respectively. Two different kinds of fodder are available. Each unit of these two contains the following amounts of these three nutrients :

Fodder	Fodder 1	Fodder 2
Nutrients A	2	1
Nutrients B	2	3
Nutrients C	1	1

The cost of fodder 1 is Rs.3 per unit and that of fodder Rs. 2, Formulate the L.P.P. to minimize the cost.

- 3) A company manufactures two types of chemicals A and B. Each chemical requires two types of raw material P and Q. The table below shows number of units of P and Q required to manufacture one unit of A and one unit of B and the total availability of P and Q.

Chemical	A	B	Availability
P	3	2	120
Q	2	5	160

The company gets profits of Rs.350 and Rs.400 by selling one unit of A and one unit of B respectively. (Assume that the entire production of A and B can be sold). How many units of the chemicals A and B should be manufactured so that the company get maximum profit? Formulate the problem as L.P.P. to maximize the profit.

- 4) A printing company prints two types of magazines A and B. The company earns Rs. 10 and Rs. 15 on magazines A and B per copy. These are processed on three machines I, II, III. Magazine A requires 2 hours on Machine I, 5 hours on Machine II and 2 hours on Machine III. Magazine B requires 3 hours on Machine I, 2 hours on Machine II and 6 hours on Machine III. Machines I, II, III are available for 36, 50, 60 hours per week respectively. Formulate the L.P.P. to determine weekly production of A and B, so that the total profit is maximum.
- 5) A manufacture produces bulbs and tubes. Each of these must be processed through two machines M_1 and M_2 . A package of bulbs require 1 hour of work on Machine M_1 and 3 hours of work on M_2 . A package of tubes require 2 hours on Machine M_1 and 4 hours on Machine M_2 . He earns a profit of Rs. 13.5 per package of bulbs and Rs. 55 per package of tubes. Formulate the LLP to maximize the profit, if he operates the machine M_1 , for atmost 10 hours a day and machine M_2 for atmost 12 hours a day.

- 6) A company manufactures two types of fertilizers F_1 and F_2 . Each type of fertilizer requires two raw materials A and B. The number of units of A and B required to manufacture one unit of fertilizer F_1 and F_2 and availability of the raw materials A and B per day are given in the table below :

Fertilizers Raw Material	F1	F2	Availability
A	2	3	40
B	1	4	70

By selling one unit of F_1 and one unit of F_2 , company gets a profit of Rs. 500 and Rs. 750 respectively. Formulate the problem as L.P.P. to maximize the profit.

- 7) A doctor has prescribed two different units of foods A and B to form a weekly diet for a sick person. The minimum requirements of fats, carbohydrates and proteins are 18, 28, 14 units respectively. One unit of food A has 4 units of fats, 14 units of carbohydrates and 8 units of protein. One unit of food B has 6 units of fat, 12 units of carbohydrates and 8 units of protein. The price of food A is 4.5 per unit and that of food B is 3.5 per unit. Form the L.P.P. so that the sick person's diet meets the requirements at a minimum cost.
- 8) If John drives a car at a speed of 60 kms/hour he has to spend Rs. 5 per km on petrol. If he drives at a faster speed of 90 kms/hour, the cost of petrol increases to 8 per km. He has Rs. 600 to spend on petrol and wishes to travel the maximum distance within an hour. Formulate the above problem as L.P.P.
- 9) The company makes concrete bricks made up of cement and sand. The weight of a concrete brick has to be least 5 kg. Cement costs Rs.20 per kg. and sand costs of Rs.6 per kg. strength consideration dictate that a concrete brick should contain minimum 4 kg. of cement and not more than 2 kg. of sand. Form the L.P.P. for the cost to be minimum.

7.2.3 Formal definitions related to L.P.P. :

- 1) **Solution of L.P.P. :** A set of values of the decision variables x_1, x_2, \dots, x_n which satisfy the conditions of given linear programming problem is called a solution to that problem.
- 2) **Feasible solution :** A solution which satisfies the given constraints is called a feasible solution.
- 3) **Optimal feasible solution :** A feasible solution which maximizes or minimizes the objective function as per the requirements is called an optimal feasible solution.
- 4) **Feasible region :** The common region determined by all the constraints of the L.P.P. is called the feasible region.

Solution of L.P.P. :

There are two methods to find the solution of L.P.P. :

- 1) Graphical method,
- 2) Simplex method.

Note : We shall restrict ourselves to graphical method.

Some definitions :

Solution :

A set of values of the variables which satisfies all the constraints of the L.P.P. is called the solution of the L.P.P.

Optimum feasible solution :

A feasible solution which optimizes (either maximizes or minimizes) the objective function of L.P.P. is called optimum feasible solution.

Theorems (without proof) :

Theorem 1 : The set of all feasible solutions of L.P.P. is a convex set.

Theorem 2 : The objective function of L.P.P. attains its optimum value (either maximum or minimum) at least at one of the vertices of convex polygon. This is known as convex polygon theorem.

Corner - Point Method :

- 1) Convert all inequations of the constraints into equations..
- 2) Draw the lines in X - Y plane.
- 3) Locate common region indicated by the constraints. This common region is feasible region.
- 4) Find the vertices of feasible region.
- 5) Find the value of the objective function z at all vertices of feasible region.

Suppose, we are expected to maximize or minimize a given objective function $z = ax + by$ in the feasible region. The feasible region is a convex region bounded by straight lines. If any linear function $z = ax + by$, is maximized in the feasible region at some point, then the point is vertex of the polygon. This can be verified by drawing a line $ax + by = c$ which passes through the feasible region and moves with different values of c.

Solve graphically the following Linear Programming Problems :


Example 1 : Maximize : $z = 9x + 13y$ subject to $2x + 3y \leq 18, 2x + y \leq 10, x \geq 0, y \geq 0$.

Solution : To draw $2x + 3y \leq 18$ and $2x + y \leq 10$
 Draw line $2x + 3y = 18$ and $2x + y = 10$

To draw	x	y	Line passes through (x, y)	Sign	Region lies on
L_1 $2x + 3y = 18$	0	6	(0, 6)	\leq	Origin side of Line L_1
	9	0	(9, 0)		
L_2 $2x + y = 10$	0	10	(0, 10)	\leq	Origin side of Line L_2
	5	0	(5, 0)		

The common shaded region is O A B C O is a feasible region with vertices O (0, 0), A (5, 0), B (3, 4), C (0, 6).

(x, y) Vertex of S	Value of $z = 9x + 13y$ at (x, y)
O (0, 0)	0
A (5, 0)	45
B (3, 4)	79
C (0, 6)	78



From the table, maximum value of $z = 79$, occurs at B (3, 4) i.e. when $x = 3, y = 4$.

fig 7.24

Solve graphically the following Linear Programming Problems :

Example 2 : Maximize : $z = 5x + 2y$ subject to $5x + y \geq 10$, $x + y \geq 6$, $x \geq 0$, $y \geq 0$.

Solution : To draw $5x + y \geq 10$ and $x + y \geq 6$
 Draw line $5x + y = 10$ and $x + y = 6$.

To draw	x	y	Line passes through (x, y)	Sign	Region lies on
L_1 $5x + y = 10$	0	10	(0, 10)	\geq	Non-origin side of Line L_1
	2	0	(2, 0)		
L_2 $x + y = 6$	0	6	(0, 6)	\geq	Non-origin side of Line L_2
	6	0	(6, 0)		

The common shaded region is feasible region with vertices A (6, 0), B (1, 5), C (0, 10).

(x, y) Vertex of S	Value of $z = 5x + 2y$ at (x, y)
A (6, 0)	30
B (1, 5)	15
C (0, 10)	20




fig 7.25

From the table, maximum value of $z = 15$, occurs at B (1, 5) i.e. when $x = 1, y = 5$.

Example 3: Maximize: $z = 3x + 4y$ subject to $x - y \geq 0$, $-x + 3y \leq 3$, $x \geq 0$, $y \geq 0$.

Solution : To draw $x - y \geq 0$ and $-x + 3y \leq 3$
 Draw line $x - y \geq 0$ and $-x + 3y = 3$.

To draw	Draw line	x	y	Line passes through (x, y)	Sign	Region lies on
$x - y \geq 0$	L_1 $x = y$	0	0	(0, 0)	\geq	A side
		1	1	(1, 1)		
$-x + 3y \leq 3$	L_2 $-x + 3y = 3$	0	1	(0, 1)	\leq	Origin side of Line L_2
		-3	0	(-3, 0)		




fig 7.26

From graph, we can see that the common shaded area is the feasible region which is unbounded (not a polygon). In such cases, the iso-profit lines can be moved away from the origin indefinitely. \therefore There is no finite maximum value of z within the feasible region.

Example 4 : Maximize : $z = 5x + 2y$ subject to $3x + 5y \leq 15$, $5x + 2y \leq 10$, $x \geq 0$, $y \geq 0$.

Solution : To draw $3x + 5y \leq 15$ and $5x + 2y \leq 10$
 Draw line $3x + 5y = 15$ and $5x + 2y = 10$.

To draw	x	y	Line passes through (x, y)	Sign	Region lies on
L_1 $3x + 5y = 15$	5	0	(5, 0)	\leq	Origin side of Line L_1
	0	3	(0, 3)		
L_2 $5x + 2y = 10$	2	0	(2, 0)	\leq	Origin side of Line L_2
	0	5	(0, 5)		

The shaded region O A B C is the feasible region with the vertices O (0, 0), A (2, 0), B($\frac{20}{19}, \frac{45}{19}$), C (0, 3)

$Z_0 = 0, Z_A = 10, Z_B = 10, Z_C = 6.$

Maximum value of z occurs at A and B and is $z = 10.$

Maximum value of z occurs at every point lying on the segment AB.

Hence there are infinite number of optimal solutions.




fig 7.27

Note : If the two distinct points produce the same minimum value then the minimum value of objective function occurs at every point on the segment joining them.

Exercise 7.4

Solve the following L.P.P. by graphical method :

- 1) Maximize : $z = 11x + 8y$ subject to $x \leq 4, y \leq 6,$
 $x + y \leq 6, x \geq 0, y \geq 0.$
- 2) Maximize : $z = 4x + 6y$ subject to $3x + 2y \leq 12,$
 $x + y \geq 4, x, y \geq 0.$
- 3) Maximize : $z = 7x + 11y$ subject to $3x + 5y \leq 26$
 $5x + 3y \leq 30, x \geq 0, y \geq 0.$
- 4) Maximize : $z = 10x + 25y$ subject to $0 \leq x \leq 3,$
 $0 \leq y \leq 3, x + y \leq 5$ also find maximum value of $z.$
- 5) Maximize : $z = 3x + 5y$ subject to $x + 4y \leq 24, 3x + y \leq 21,$
 $x + y \leq 9, x \geq 0, y \geq 0$ also find maximum value of $z.$
- 6) Minimize : $z = 7x + y$ subject to $5x + y \geq 5, x + y \geq 3,$
 $x \geq 0, y \geq 0.$
- 7) Minimize : $z = 8x + 10y$ subject to $2x + y \geq 7, 2x + 3y \geq 15,$
 $y \geq 2, x \geq 0, y \geq 0.$
- 8) Minimize : $z = 6x + 21y$ subject to $x + 2y \geq 3, x + 4y \geq 4,$
 $3x + y \geq 3, x \geq 0, y \geq 0.$



Let's remember!

* Working rule to formulate the L.P.P. :

Step 1 : Identify the decision variables and assign the symbols x, y or x_1, x_2 to them. Introduce non-negativity constraints.

Step 2 : Identify the set of constraints and express them as linear inequation in terms of the decision variables.

Step 3 : Identify the objective function to be optimized (i.e. maximized or minimized) and express it as a linear function of decision variables.

* Let R be the feasible region (convex polygon) for a L.P.P. and Let $z = ax + by$ be the objective functions then the optimal value (maximum or minimum) of z occurs at least one of the corner points (vertex) of the feasible region.

* Corner point method for solving L.P.P. graphically :

Step 1 : Find the feasible region of the L.P.P.

Step 2 : Determine the vertices of the feasible region either by inspection or by solving the two equations of the lines intersecting at that points.

Step 3 : Find the value of the objective function z , at all vertices of feasible region.

Step 4 : Determine the feasible solution which optimizes the value of the objective function.

Miscellaneous Exercise

I) Select the appropriate alternatives for each of the following :

- 1) The value of objective function is maximum under linear constraints _____.
 A) at the centre of feasible region
 B) at (0, 0)
 C) at a vertex of feasible region
 D) the vertex which is of maximum distance from (0, 0)
- 2) Which of the following is correct _____.
 A) every L.P.P. has an optimal solution
 B) a L.P.P. has unique optimal solution
 C) if L.P.P. has two optimal solutions then it has infinite number of optimal solutions
 D) the set of all feasible solution of L.P.P. may not be convex set
- 3) Objective function of L.P.P. is _____.
 A) a constraint
 B) a function to be maximized or minimized
 C) a relation between the decision variables
 D) equation of a straight line
- 4) The maximum value of $z = 5x + 3y$ subjected to the constraints $3x + 5y \leq 15$, $5x + 2y \leq 10$, $x, y \geq 0$ is _____.
 A) 235
 B) $\frac{235}{9}$
 C) $\frac{235}{19}$
 D) $\frac{235}{3}$
- 5) The maximum value of $z = 10x + 6y$ subjected to the constraints $3x + y \leq 12$, $2x + 5y \leq 34$, $x \geq 0$, $y \geq 0$. _____.
 A) 56
 B) 65
 C) 55
 D) 66
- 6) The point at which the maximum value of $x + y$ subject to the constraints $x + 2y \leq 70$, $2x + y \leq 95$, $x \geq 0$, $y \geq 0$ is obtained at _____.
 A) (30, 25)
 B) (20, 35)
 C) (35, 20)
 D) (40, 15)
- 7) Of all the points of the feasible region, the optimal value of z obtained at the point lies _____.
 A) inside the feasible region
 B) at the boundary of the feasible region
 C) at vertex of feasible region
 D) outside the feasible region

- 8) Feasible region is the set of points which satisfy _____.
- A) the objective function B) all of the given constraints
C) some of the given constraints D) only one constraint
- 9) Solution of L.P.P. to minimize $z = 2x + 3y$ s.t. $x \geq 0, y \geq 0, 1 \leq x + 2y \leq 10$ is _____.
- A) $x = 0, y = \frac{1}{2}$ B) $x = \frac{1}{2}, y = 0$
C) $x = 1, y = 2$ D) $x = \frac{1}{2}, y = \frac{1}{2}$
- 10) The corner points of the feasible solution given by the inequation $x + y \leq 4, 2x + y \leq 7, x \geq 0, y \geq 0$ are _____.
- A) (0, 0), (4, 0), (7, 1), (0, 4) B) (0, 0), ($\frac{7}{2}$, 0), (3, 1), (0, 4)
C) (0, 0), ($\frac{7}{2}$, 0), (3, 1), (0, 7) D) (0, 0), (4, 0), (3, 1), (0, 7)
- 11) The corner points of the feasible solution are (0, 0), (2, 0), ($\frac{12}{7}, \frac{3}{7}$), (0, 1). Then $Z = 7x + y$ is maximum at _____.
- A) (0, 0) B) (2, 0)
C) ($\frac{12}{7}, \frac{3}{7}$) D) (0, 1)
- 12) If the corner points of the feasible solution are (0, 0), (3, 0), (2, 1) and ($0, \frac{7}{3}$) the maximum value of $z = 4x + 5y$ is _____.
- A) 12 B) 13
C) $\frac{35}{3}$ D) 0
- 13) If the corner points of the feasible solution are (0, 10), (2, 2) and (4, 0) then the point of minimum $z = 3x + 2y$ is _____.
- A) (2, 2) B) (0, 10)
C) (4, 0) D) (3, 4)
- 14) The half plane represented by $3x + 2y < 8$ constraints the point _____.
- A) ($1, \frac{5}{2}$) B) (2, 1)
C) (0, 0) D) (5, 1)
- 15) The half plane represented by $4x + 3y > 14$ contains the point _____.
- A) (0, 0) B) (2, 2)
C) (3, 4) D) (1, 1)

II) Solve the following :

- 1) Solve each of the following inequations graphically using X Y plane.
- i) $4x - 18 \geq 0$ ii) $-11x - 55 \leq 0$
iii) $5y - 12 \geq 0$ iv) $y \leq -3.5$
- 2) Sketch the graph of each of following inequations in XOY co-ordinate system.
- i) $x \geq 5y$ ii) $x + y \leq 0$
iii) $2y - 5x \geq 0$ iv) $1x + 51 \leq y$
- 3) Find graphical solution for each of the following system of linear inequation.
- i) $2x + y \geq 2, x - y \leq 1$ ii) $x + 2y \geq 4, 2x - y \leq 6$
iii) $3x + 4y \leq 12, x - 2y \geq 2, y \geq -1$
- 4) Find feasible solution for each of the following system of linear inequations graphically.
- i) $2x + 3y \leq 12, 2x + y \leq 8, x \geq 0, y \geq 0$
ii) $3x + 4y \geq 12, 4x + 7y \leq 28, x \geq 0, y \geq 0$

- 5) Solve each of the following L.P.P.
- i) Maximize $z = 5x_1 + 6x_2$ subject to $2x_1 + 3x_2 \leq 18$, $2x_1 + x_2 \leq 12$, $x_1 \geq 0$, $x_2 \geq 0$
 - ii) Maximize $z = 4x + 2y$ subject to $3x + y \geq 27$, $x + y \geq 21$
 - iii) Maximize $z = 6x + 10y$ subject to $3x + 5y \leq 10$, $5x + 3y \leq 15$, $x \geq 0$, $y \geq 0$
 - iv) Maximize $z = 2x + 3y$ subject to $x - y \geq 3$, $x \geq 0$, $y \geq 0$
- 6) Solve each of the following L.P.P.
- i) Maximize $z = 4x_1 + 3x_2$ subject to $3x_1 + x_2 \leq 15$, $3x_1 + 4x_2 \leq 24$, $x_1 \geq 0$, $x_2 \geq 0$
 - ii) Maximize $z = 60x + 50y$ subject to $x + 2y \leq 40$, $3x + 2y \leq 60$, $x \geq 0$, $y \geq 0$
 - iii) Maximize $z = 4x + 2y$ subject to $3x + y \geq 27$, $x + y \geq 21$, $x + 2y \geq 30$; $x \geq 0$, $y \geq 0$
- 7) A carpenter makes chairs and tables. Profits are Rs.140/- per chair and < 210/- per table. Both products are processed on three machines : Assembling, Finishing and Polishing. The time required for each product in hours and availability of each machine is given by following table:

Product Machine	Chair (x)	Table (y)	Available time (hours)
Assembling	3	3	36
Finishing	5	2	50
Polishing	2	6	60

Formulate the above problem as L.P.P. Solve it graphically.

Formulate and solve the following Linear Programming Problems using graphical method :

- 8) A company manufactures bicycles and tricycles, each of which must be processed through two machines A and B. Maximum availability of Machine A and B is respectively 120 and 180 hours. Manufacturing a bicycle requires 6 hours on Machine A and 3 hours on Machine B. Manufacturing a tricycles requires 4 hours on Machine A and 10 hours on Machine B. If profits are Rs.180/- for a bicycle and Rs.220/- for a tricycle. Determine the number of bicycles and tricycles that should be manufactured in order to maximize the profit.
- 9) A factory produced two types of chemicals A and B. The following table gives the units of ingredients P and Q (per kg) of chemicals A and B as well as minimum requirements of P and Q and also cost per kg. chemicals A and B :

Chemicals in units Ingredients per kg.	A (x)	B (y)	Minimum requirements in units
P	1	2	80
Q	3	1	75
Cost (in Rs.)	4	6	--

Find the number of units of chemicals A and B should be produced sp as to minimize the cost.

- 10) A company produces mixers and food processors. Profit on selling one mixer and one food processor is Rs. 2,000/- and Rs. 3,000/- respectively. Both the products are processed through three Machines A, B, C. The time required in hours by each product and total time available in hours per week on each machine are as follows :

Product Machine	Mixer (per unit)	Food Processor (per unit)	Available time
A → A	3	3	36
B → B	5	2	50
C → C	2	6	60

How many mixers and food processors should be produced to maximize the profit?

- 11) A chemical company produces a chemical containing three basic elements A, B, C so that it has at least 16 liters of A, 24 liters of B and 18 liters of C. This chemical is made by mixing two compounds I and II. Each unit of compound I has 4 liters of A, 12 liters of B, 2 liters of C. Each unit of compound II has 2 liters of A, 2 liters of B and 6 liters of C. The cost per unit of compound I is Rs.800/- and that of compound II is Rs.640/-. Formulate the problem as L.P.P. and solve it to minimize the cost.
- 12) A person makes two types of gift items A and B requires the services of a cutter and a finisher. Gift item A requires 4 hours of cutter's time and 2 hours of finisher's time. B requires 2 hours of cutter's time and 4 hours of finisher's time. The cutter and finisher have 208 hours and 152 hours available times respectively every month. The profit of one gift item of type A is Rs.75/- and on gift item B is Rs.125/-. Assuming that the person can sell all the gift items produced, determine how many gift items of each type should he make every month to obtain the best returns?
- 13) A firm manufactures two products A and B on which profit earned per unit Rs.3/- and Rs.4/- respectively. Each product is processed on two machines M_1 and M_2 . The product A requires one minute of processing time on M_1 and two minute of processing time on M_2 , B requires one minute of processing time on M_1 and one minute of processing time on M_2 . Machine M_1 is available for use for 450 minutes while M_2 is available for 600 minutes during any working day. Find the number of units of product A and B to be manufactured to get the maximum profit.
- 14) A firm manufacturing two types of electrical items A and B, can make a profit of Rs.20/- per unit of A and Rs.30/- per unit of B. Both A and B make use of two essential components a motor and a transformer. Each unit of A requires 3 motors and 2 transformers and each units of B requires 2 motors and 4 transformers. The total supply of components per month is restricted to 210 motors and 300 transformers. How many units of A and B should the manufacture per month to maximize profit? How much is the maximum profit?



ANSWERS

1. Mathematical Logic



Exercise 1.1

- 1) (i) Statement, F (ii) Not statement (iii) Not statement
 (iv) Statement, T (v) Not statement (vi) Statement, T
 (vii) Not statement (viii) Statement, T (ix) Not statement
 (x) Statement, T (xi) Statement, F (xii) Not statement
 (xiii) Statement, T (xiv) Statement, T (xv) Not statement
- 2) (i) $p \wedge q$ (ii) $p \vee q$ (iii) $p \leftrightarrow q$
 (iv) $\sim p \wedge \sim q$ (v) $p \rightarrow q$ (vi) $p \leftrightarrow q$
 (vii) $\sim p \wedge q$
- 3) (i) F (ii) F (iii) F
 (iv) T (v) T (vi) T
 (vii) T
- 4) (i) T (ii) T (iii) F
 (iv) T (v) F (vi) F
 (vii) T (viii) T
- 5) (i) Tirupati is not in Andhra Pradesh.
 (ii) 3 is a root of the equation $x^2 + 3x - 18 = 0$.
 (iii) $\sqrt{2}$ is not a rational number.
 (iv) Polygon ABCDE is not a pentagon.
 (v) $7 + 3 \neq 5$



Exercise 1.2

- 1) (i) TTFT (ii) FFFF (iii) TTFT FFFT
 (iv) FTTTTTTT (v) FFFF (vi) TFFT
 (vii) TTTT (viii) TTTTTTTT (ix) FTTTTTTT
 (x) TFTFTTFF
- 3) (i) Tautology (ii) Tautology (iii) Contingency
 (iv) Contingency (v) Tautology (vi) Contingency
 (vii) Contingency (viii) Contingency (ix) Contingency
 (x) Contradiction



Exercise 1.3

- 1) (i) T (ii) T (iii) F
 (iv) F (v) T (vi) T

- 2) (i) $p \wedge (q \vee r)$ (ii) $p \vee (q \vee r)$ (iii) $(p \wedge q) \vee (r \wedge s)$
 (iv) $p \vee \sim q$ (v) $(\sim p \wedge q) \vee (\sim r \vee s)$
 (vi) $\sim p \vee (\sim q \vee (p \wedge q) \vee \sim r)$
 (vii) $[\sim(p \wedge q) \vee [p \wedge \sim(q \vee \sim s)]]$
 (viii) $t \vee \{p \vee (q \wedge r)\}$
 (ix) $\sim p \wedge (q \vee r) \vee c$
 (x) $(p \wedge q) \wedge t$

- 3) (i) $x + 8 \leq 11$ and $y - 3 \neq 6$
 (ii) $11 \geq 15$ and $25 \leq 20$
 (iii) Quadrilateral is a square but not rhombus or quadrilateral is a rhombus but not a square.
 (iv) It is not cold or not raining.
 (v) It is raining and we will not go or not play football.
 (vi) $\sqrt{2}$ is not a rational number.
 (vii) Some natural numbers are not whole numbers.
 (viii) $\exists n \in \mathbb{N}, n^2 + n + 2$ is not divisible by 4.
 (ix) $\forall x \in \mathbb{N}, x - 17 \geq 20$.

- 4) (i) Converse :If $x^2 < y^2$ then $x < y$
 Inverse :If $x \geq y$ then $x^2 \geq y^2$.
 Contrapositive :If $x^2 \geq y^2$ then $x \geq y$.
 (ii) Converse :If a family becomes literate then the woman in it is literate.
 Inverse :If the woman in the family is not literate then the family does not become literate.
 Contrapositive :If a family does not become literate then the woman in the family is not literate.
 (iii) Converse :If pressure increases then surface area decreases.
 Inverse :If surface area does not decrease then pressure does not increase.
 Contrapositive :If pressure does not increase then surface area does not decrease.
 (iv) Converse :If current decreases then voltage increases.
 Inverse :If voltage does not increase then current does not decrease.
 Contrapositive :If current does not decrease then voltage does not increase.




Exercise 1.4

- 1) (i) $\sim q \wedge \sim p$ (ii) $\sim p \vee q$ (iii) $\sim p \wedge q$
 (iv) $(\sim p \wedge q) \vee \sim r$ (v) $p \wedge (\sim p \wedge q)$ (vi) $(p \wedge q) \wedge (\sim p \wedge q)$
 (vii) $(p \vee \sim q) \wedge (\sim p \vee q)$ (viii) $(p \wedge q) \wedge (\sim p \vee q)$
- 2) (i) A man is not a judge or he is honest.
 (ii) 2 is not rational number or is $\sqrt{2}$ irrational number.
 (iii) $f(2) \neq 0$ or $f(x)$ is divisible by $(x - 2)$.



Exercise 1.5

- 1) i) $p \vee (q \wedge r)$ 11111000
 ii) $(\sim p \wedge q) \vee (p \wedge \sim q)$ 0110
 iii) $[(p \wedge (\sim q \vee r)) \vee [\sim q \wedge \sim r]]$ 10110001
 iv) $(p \vee q) \wedge \sim r \wedge (\sim p \vee r)$ 01000100
 v) $[p \vee (\sim p \wedge \sim q)] \vee (p \wedge q)$ 1101
 vi) $(p \vee q) \wedge (q \vee r) \wedge (r \vee p)$ 11101000



- 4) (i) $(p \vee \sim q) \vee (\sim p \wedge q)$ 1111
 The lamp will glow irrespective of the status of the switches.
 (ii) $[p \vee (\sim p \wedge \sim q)] \vee (p \wedge q)$ 1101
 The lamp will not glow when switch S_1 is OFF and S_2 is ON otherwise it will glow.

(iii) $[p \vee \sim q \vee \sim r] \wedge [(p \vee (q \wedge \sim r))]$ 11110000
 The lamp will glow if S_1 is ON and any status of S_2 .

- 5) (i) P
 (ii) $\sim p \vee \sim q$
 (iii) P
 (iv) $(q \wedge r) \vee$

Miscellaneous Exercise - 1

1)

i	ii	iii	iv	v	vi	vii
B	A	C	B	A	D	C

- 2) (i) Statement, T (ii) Statement, T (iii) Statement, F (iv) Not a statement
 (v) Statement, T (vi) Statement, T
- 3) (i) T(ii) F (iii) T (iv) T(v) T (vi) F
- 4) (i) T(ii) F (iii) T (iv) F
- 5) (i) $\exists n \in \mathbb{N}$ such that $n+7 \not> 6$. $\exists n \in \mathbb{N}$ such that $n + t \leq 6$
 (ii) $\forall x \in A, x + 9 \not> 15$ on $x \in A, \forall x + 9 > 15$.
 (iii) All triangles are not equilateral triangles.

6) (i)

p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$
T	T	T	T
T	F	T	T
F	T	F	T
F	F	T	T

(ii)

(I)	(II)	(III)	(IV)	(V)	(VI)	(VII)	(VIII)
p	q	$\sim p$	$\sim q$	$p \wedge q$	$\sim (p \wedge q)$	$\sim p \vee \sim q$	$(vi) \leftrightarrow (vii)$
T	T	F	F	T	F	F	T
T	T	F	T	F	T	T	T
F	T	T	F	F	T	T	T
F	T	T	T	F	T	T	T

(iii)

(I)	(II)	(III)	(IV)	(V)	(VI)	(VII)
p	q	$\sim p$	$\sim q$	$\sim p \wedge \sim q$	$\sim (\sim p \wedge \sim q)$	$\sim (\sim p \wedge \sim q) \wedge q$
T	T	F	F	F	T	T
T	F	F	T	F	T	T
F	T	T	F	F	T	T
F	F	T	T	T	T	T

(iv)


(I)	(II)	(III)	(IV)	(V)	(VI)	(VII)	(IX)
p	q	r	$p \wedge q$	$(p \wedge q) \vee q$	$\sim r$	$\sim r \vee (\text{iv})$	$(\text{v}) \wedge (\text{vii})$
T	T	T	T	T	F	T	T
T	T	F	T	T	T	T	T
T	F	T	F	T	F	F	F
T	F	F	F	F	T	T	F
F	T	T	F	T	F	F	F
F	T	F	F	F	T	T	F
F	F	T	F	T	F	F	F
F	F	F	F	F	T	T	F

(v)


(I)	(II)	(III)	(IV)	(V)	(VI)	(VII)	(VIII)	(IX)
p	q	r	$\sim p$	$\sim p \wedge q$	$q \rightarrow r$	$p \rightarrow r$	$(\text{i}) \wedge (\text{vi})$	$(\text{viii}) \rightarrow (\text{vii})$
T	T	T	F	T	T	T	T	T
T	T	F	F	T	F	F	F	T
T	F	T	F	F	T	T	F	T
T	F	F	F	F	T	F	F	T
F	T	T	T	T	T	T	T	T
F	T	F	T	T	F	T	F	T
F	F	T	T	T	T	T	T	T
F	F	F	T	T	T	T	T	T

- 7) (i) Tautology (ii) Contradiction (iii) Contradiction (iv) Tautology
 (v) Tautology (vi) Tautology (vii) Contingency (viii) Tautology
- 8) (i) T, T (ii) T, F (iii) T, F or F, T or F, F
- 11) (i) $\sim q \wedge (\sim p \vee r)$ (ii) $\sim p \vee (\sim q \wedge \sim r)$ (iii) $(p \wedge \sim q) \vee r$ (iv) $(p \vee \sim q) \wedge (\sim p \vee q)$
- 12) (i) $(p \wedge q) \vee \sim p \vee (p \wedge \sim q)$ 11111 (ii) $(p \vee q) \wedge (p \vee r)$ 11111000

13) (i)




(ii)



14) (i) Logically equivalent
15)

(ii) Logically equivalent




16) Current always flows

17) $(p \vee \sim q \vee \sim r) \wedge [p \vee (q \wedge r)]$

$1 \ 1 \ 1 \ 1 \ 0 \ 0 \ 0 \ 0$ which is same as p .

Hence we can conclude that the given switching circuit is equivalent to a simple circuit with only one switch S_1 .



2. Matrics



Exercise 2.1

1) $\begin{bmatrix} -1 & 3 \\ 1 & 0 \end{bmatrix}$

2) $\begin{bmatrix} -1 & -6 & -1 \\ 2 & 5 & 4 \end{bmatrix}$

3) $A \sim \begin{bmatrix} 4 & 5 \\ 3 & 1 \end{bmatrix}$ $B \sim \begin{bmatrix} 4 & 5 \\ 3 & 1 \end{bmatrix}$ The new matrices are equal.

4) $\begin{bmatrix} -2 & 4 & -7 \\ 2 & 6 & 8 \end{bmatrix}$

5) $\begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 9 & 9 & 21 \end{bmatrix}$

6) $\begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & 2 \\ 9 & 9 & 21 \end{bmatrix}$

\therefore The transformations are commutative.

7) $\begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix}$

9) $\begin{bmatrix} 1 & -1 & 2 \\ 0 & 3 & -1 \\ 0 & 0 & \frac{17}{3} \end{bmatrix}$



Exercise 2.2

1) (i) 4, 3, -2, -1.

(ii) -3, -12, 6, -1, 3, 2, -11, -9, 1.

2) (i) $\begin{bmatrix} -1 & -4 \\ -3 & 1 \end{bmatrix}$

(ii) $\begin{bmatrix} -11 & -10 & -6 \\ 6 & -5 & 3 \\ -2 & -7 & 1 \end{bmatrix}$

3) (i) $\begin{bmatrix} 5 & 3 \\ -3 & 2 \end{bmatrix}$

(ii) $\begin{bmatrix} -3 & -1 & -11 \\ -12 & 3 & -9 \\ 6 & 2 & 1 \end{bmatrix}$

5) (i) $\frac{1}{13} \begin{bmatrix} 2 & -5 \\ 3 & -1 \end{bmatrix}$

(ii) $\frac{1}{14} \begin{bmatrix} 3 & 2 \\ -4 & 2 \end{bmatrix}$

(iii) $-\frac{1}{3} \begin{bmatrix} -3 & 0 & 0 \\ 3 & -1 & 0 \\ -9 & -2 & 3 \end{bmatrix}$

(iv) $-\frac{1}{10} \begin{bmatrix} 10 & -10 & 2 \\ 0 & 2 & 4 \\ 0 & 0 & 2 \end{bmatrix}$

6) (i) $-\frac{1}{5} \begin{bmatrix} -1 & -2 \\ -2 & 1 \end{bmatrix}$

(ii) $\begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$

(iii) $\begin{bmatrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \\ 4 & 3 & 1 \\ \frac{5}{3} & \frac{3}{2} & \frac{1}{2} \end{bmatrix}$

(iv) $\begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$

Miscellaneous Exercise - 2(A)

1) Using $C_1 - 2C_2$, $C_1 + 3C_3$ and $C_2 - 3C_3$, We get the required result.

2) Using $R_1 - R_2$, $R_3 - R_2$, $-R_2$, $R_1 - R_2$, $R_3 - R_2$, $-R_3$, $R_1 - R_3$, $R_2 - R_3$, we get the required result.
(There can be another sequence of the transformations.)

3) The invertible matrices are (i), (iii), (v), (vi), (vii) and not invertible matrices are (ii), (iv), (viii) and (ix).

4) $AB = \begin{bmatrix} 6 & -3 \\ -4 & 1 \end{bmatrix}$ and it is invertible.

$$5) \quad A^{-1} = \begin{bmatrix} \frac{1}{x} & 0 & 0 \\ 0 & \frac{1}{y} & 0 \\ 0 & 0 & \frac{1}{z} \end{bmatrix} \quad A^{-1} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

$$6) \quad (i) \quad X = \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & \frac{1}{2} \end{bmatrix}$$

$$7) \quad (i) \quad \frac{1}{5} \begin{bmatrix} 3 & 1 \\ -2 & 1 \end{bmatrix} \quad (ii) \quad -\frac{1}{3} \begin{bmatrix} -1 & -1 \\ -1 & 2 \end{bmatrix}$$

$$(iii) \quad \begin{bmatrix} 7 & -3 \\ -2 & 1 \end{bmatrix} \quad (iv) \quad \frac{1}{29} \begin{bmatrix} 7 & 3 \\ -5 & 2 \end{bmatrix}$$

$$(v) \quad \begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix} \quad (vi) \quad \begin{bmatrix} 7 & -10 \\ 2 & -3 \end{bmatrix}$$

$$(vii) \quad -\frac{1}{25} \begin{bmatrix} 10 & 0 & -15 \\ -5 & -5 & 0 \\ -10 & 5 & 10 \end{bmatrix} \quad (viii) \quad \frac{1}{25} \begin{bmatrix} 25 & -10 & -15 \\ -10 & 4 & 11 \\ -15 & 1 & 9 \end{bmatrix}$$

$$(ix) \quad \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix} \quad (x) \quad \begin{bmatrix} 3 & 6 & 2 \\ 1 & 2 & 1 \\ 2 & 5 & 2 \end{bmatrix}$$

$$8) \quad A^{-1} = \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$9) \quad AB = \begin{bmatrix} 11 & 3 \\ 7 & 2 \end{bmatrix}, (AB)^{-1} = \begin{bmatrix} 2 & -3 \\ -7 & 11 \end{bmatrix}$$

$$A^{-1} = \begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix} \quad B^{-1} = \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix}$$

$$11) \quad X = \begin{bmatrix} \frac{4}{5} & 1 \\ \frac{2}{5} & 1 \end{bmatrix} \quad 12) \quad X = -\frac{1}{3} \begin{bmatrix} 1 \\ 7 \\ -6 \end{bmatrix}$$

$$13) X = \begin{bmatrix} 2 & 3 \\ 1 & 1 \end{bmatrix} \qquad 14) \begin{bmatrix} 13 & 2 & -7 \\ -3 & -1 & 2 \\ -2 & 0 & 1 \end{bmatrix}$$

$$15) -\frac{1}{6} \begin{bmatrix} -4 & 2 & -2 \\ 3 & 0 & -3 \\ -2 & -2 & 2 \end{bmatrix} \qquad 16) \frac{1}{3} \begin{bmatrix} 0 & -2 & 1 \\ 6 & 1 & -5 \\ -3 & 0 & 3 \end{bmatrix}$$

$$17) \frac{1}{6} \begin{bmatrix} 4 & -2 & 2 \\ -3 & 0 & 3 \\ 2 & 2 & -2 \end{bmatrix} \qquad 18) \begin{bmatrix} 13 & 2 & -7 \\ -3 & -1 & 2 \\ -2 & 0 & 1 \end{bmatrix}$$

19) Hint : Use the definition of the co-factors and the value of the determinant by considering.

$$A = [a_{ij}]_{3 \times 3}$$

$$20) X = \frac{1}{6} \begin{bmatrix} 4 & 4 & 2 \\ 11 & 8 & -5 \\ 10 & 10 & 2 \end{bmatrix}$$



Exercise 2.3

- 1) (i) 0, 1 (ii) 3, 1 (iii) Not solvable
- 2) (i) 4, -3 (ii) $\frac{1}{2}, \frac{1}{2}$ (iii) 1, 2 (iv) 2, -3
- 3) Rs. 5 for a pencil Rs. 8 for a pen and Rs.8 for an eraser.
- 4) The numbers are 1, -2, 3.
- 5) The cost price of one T.V. set is Rs.3000 and of one V.C.R. is Rs. 13,000.
The selling price of one T.V.Set is Rs.4000 and that of V.C.R. is Rs. 13,500.

Miscellaneous exercise - 2 (B)

I)

1	2	3	4	5	6	7	8	9	10	11	12
A	B	D	B	B	B	B	A	B	B	B	D

- II) 1) (i) $-\frac{5}{11}, \frac{12}{11}$ (ii) $2 - \frac{4}{a}, 0, -1 + \frac{4}{a}$
- (iii) $x = 3, y = 2, z = -2$ (iv) $x = 2, y = -3$
- (v) $x = \frac{5}{2}, y = \frac{3}{2}, z = -2,$
- 2) (i) 1, 1, 1 (ii) $\frac{1}{3}, \frac{2}{3}, 1$

- (iii) 1, 2, 1 (iv) 1, 2, 3
 (v) 3, 2, 1 (vi) -1, 1, 2

- 3) The numbers are 1, 2, 3
 4) Cost of a pencil, a pen and a book is respectively Rs.10, Rs.15 and Rs.25.
 5) The costs are $3, \frac{5}{3}, \frac{4}{3}$
 6) The numbers are 1, -1, 2
 7) 1750, 1500, 1750
 8) Maths Rs.150, Phy. Rs.30, Chem. Rs. 30



3. Trigonometric Functions



Exercise 3.1

- 1) (i) $\frac{\pi}{3}, \frac{5\pi}{3}$ (ii) $\frac{\pi}{6}, \frac{11\pi}{6}$ (iii) $\frac{\pi}{6}, \frac{7\pi}{6}$ (iv) $0, \pi$
- 2) (i) $\frac{7\pi}{6}, \frac{11\pi}{6}$ (ii) $\frac{3\pi}{4}, \frac{7\pi}{4}$ (iii) $\frac{4\pi}{3}, \frac{5\pi}{3}$
- 3) (i) $n\pi + (-1)^n \frac{\pi}{6}, n \in \mathbb{Z}$ (ii) $2n\pi \pm \frac{\pi}{6}, n \in \mathbb{Z}$
 (iii) $n\pi + \frac{\pi}{6}, n \in \mathbb{Z}$ (iv) $n\pi + \frac{\pi}{2}, n \in \mathbb{Z}$
- 4) (i) $2n\pi \pm \frac{\pi}{4}, n \in \mathbb{Z}$ (ii) $n\pi + (-1)^n \frac{5\pi}{4}, n \in \mathbb{Z}$ (iii) $n\pi + \frac{3\pi}{4}, n \in \mathbb{Z}$
- 5) (i) $\frac{n\pi}{2} + (-1)^n \frac{\pi}{12}, n \in \mathbb{Z}$ (ii) $\frac{3n\pi}{2} + \frac{\pi}{2}, n \in \mathbb{Z}$ (iii) $\frac{n\pi}{4} + \frac{3\pi}{16}, n \in \mathbb{Z}$
- 6) (i) $n\pi \pm \frac{\pi}{6}, n \in \mathbb{Z}$ (ii) $n\pi \pm \frac{\pi}{6}, n \in \mathbb{Z}$ (iii) $\frac{n\pi}{3}, n \in \mathbb{Z}$
- 7) (i) $n\pi, n \in \mathbb{Z}$ (ii) $n\pi$ or $n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z}$
 (iii) $2n\pi$ or $2n\pi \pm \frac{\pi}{2}, n \in \mathbb{Z}$
- 8) (i) and (iv) have solutions (ii) and (iii) do not have solutions



Exercise 3.2

- 1) (i) (1,1) (ii) (0,4) (iii) $\left(-\frac{3}{4\sqrt{2}}, \frac{3}{4\sqrt{2}}\right)$ (iv) $\left(\frac{1}{4}, \frac{\sqrt{3}}{4}\right)$

2) (i) $\left(2, \frac{\pi}{4}\right)$ (ii) $\left(\frac{1}{2}, \frac{\pi}{2}\right)$ (iii) $\left(2, \frac{5\pi}{3}\right)$ (iv) $\left(3, \frac{\pi}{3}\right)$

3) (i) $2 : \sqrt{6} : 1 + \sqrt{3}$

10) (i) $\frac{4}{5}$ (ii) $\frac{1}{\sqrt{10}}$ (iii) $\frac{3}{\sqrt{10}}$ (iv) $\frac{1}{3}$ (v) 216 (vi) $\frac{3}{5}$



Exercise 3.3

1) (i) $\frac{\pi}{6}$ (ii) $\frac{\pi}{6}$ (iii) $-\frac{\pi}{4}$ (iv) $-\frac{\pi}{3}$ (v) $\frac{\pi}{4}$ (vi) $\frac{2\pi}{3}$

2) (i) $\frac{3\pi}{4}$ (ii) $\frac{2\pi}{3}$ (iii) $-\frac{\pi}{3}$ (iv) $-\frac{\pi}{12}$

Miscellaneous exercise - 3

I)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
B	A	A	A	D	C	A	B	A	C	B	D	A	B	D	A	B	A	B	B

II) i) $\left\{\frac{7\pi}{12}, \frac{11\pi}{12}, \frac{19\pi}{12}, \frac{23\pi}{12}\right\}$ ii) $\left\{\frac{3\pi}{12}, \frac{7\pi}{12}, \frac{11\pi}{12}, \frac{15\pi}{12}, \frac{19\pi}{12}, \frac{23\pi}{12}\right\}$

2) (i) $\left\{\frac{5\pi}{8}, \frac{7\pi}{8}, \frac{13\pi}{8}, \frac{15\pi}{8}\right\}$

(ii) $\left\{\frac{3\pi}{20}, \frac{7\pi}{20}, \frac{11\pi}{20}, \frac{15\pi}{20}, \frac{19\pi}{20}, \frac{23\pi}{20}, \frac{27\pi}{20}, \frac{31\pi}{20}, \frac{35\pi}{20}, \frac{39\pi}{20}\right\}$

(iii) $\left\{\frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}\right\}$

3) (i) and (ii) have solution, (iii) and iv) do not have solutions

4) (i) $n\pi + \frac{2\pi}{3}, n \in \mathbb{Z}$ (ii) $n\pi \pm \frac{\pi}{3}, n \in \mathbb{Z}$ (iii) $(2n+1)\pi$ or $2n\pi + \frac{\pi}{2}, n \in \mathbb{Z}$

iv) $2n\pi \pm \frac{\pi}{2}, n \in \mathbb{Z}$

10) $c = \sqrt{6}, A = 105^\circ, B = 15^\circ$

19) (i) $\frac{3\pi}{5}$ (ii) $\frac{\pi}{6}$

26) $\frac{\pi}{4}$

27) $\frac{1}{\sqrt{3}}$

28) 0

29) $\frac{1}{6}$



4. Pair of Straight Lines



Exercise 4.1

- 1) (i) $6x^2 + xy - y^2 = 0$
 (ii) $x^2 - xy - 6y^2 + x + 7y - 2 = 0$
 (iii) $xy - 3x - 2y + 6 = 0$
 (iv) $6x^2 - 7xy - 3y^2 - 3x + 32y - 45 = 0$
 (v) $3x^2 + 11xy + 6y^2 - 16x - 13y + 5 = 0$
- 2) (i) $y = 0, 7x + 3y = 0$
 (ii) $\sqrt{5}x - 3y = 0, \sqrt{5}x + 3y = 0,$
 (iii) $x = 0, x - 4y = 0$
 (iv) $3x + 2y = 0, x - 4y = 0$
 (v) $3x + \sqrt{3}y = 0, x = \sqrt{3}y = 0$
 (vi) $(\operatorname{cosec} \alpha - \cot \alpha)x + y = 0, (\operatorname{cosec} \alpha + \cot \alpha)x - y = 0$
 (vii) $(\sec \alpha - \tan \alpha)x + y = 0, (\sec \alpha + \tan \alpha)x - y = 0$
- 3) (i) $3x^2 + 8xy + 5y^2 = 0$
 (ii) $x^2 + 2xy - 5y^2 = 0$
 (iii) $3x^2 - xy = 0$
 (iv) $4xy + 3y^2 = 0$
- 4) (i) -2 (ii) ± 2 (iii) 12
- 5) (i) $25a + 16b = 40h$ (ii) $9a + 6h + b = 0$
- 6) $ap^2 + 2hpq + bq^2 = 0$
- 7) $3x^2 - y^2 = 0$



Exercise 4.2

- 3) $k = 4$ 4) i) 30° ii) $\tan^{-1} \left(\frac{3}{5} \right)$ iii) 45° iv) 60°
- 5) $23x^2 + 48xy + 3y^2 = 0$ 7) $x^2 - 3y^2 = 0$



Exercise 4.3

- 1) (i) $2x^2 + 3xy - 9y^2 - 5x - 24y - 7 = 0$ (ii) $x^2 + xy - y^2 - x - 8y - 11 = 0$
- 2) $h^2 - ab = -1 < 0$
- 3) $2x - 3y + 4 = 0$ and $x + y - 5 = 0$ are separate equations of lines.
- 4) $2x - y + 3 = 0$ and $x + y - 1 = 0$ are separate equations. $\theta = \tan^{-1}(3)$.
- 5) (i) $x - y - 3 = 0, x - 2y - 4 = 0$ (ii) $2x - y + 4 = 0, 5x + 3y - 1 = 0$
- 6) (i) -12 (ii) 15 (iii) -6
- 7) $p = -3, q = -8$
- 8) $p = 8, q = 1$
- 9) $36x^2 - 25xy - 252x + 350y - 784 = 0$
- 10) $7x - 8y = 0$
- 11) $(1, 0)$

Miscellaneous exercise - 4

I.

1	2	3	4	5	6	7	8	9	10	11	12	13	14
B	B	B	A	D	D	A	B	B	B	C	C	D	D

- II. 1) (i) $x^2 - y^2 = 0$ (ii) $2x^2 + 3xy + y^2 - 7x - 4y + 3 = 0$ (iii) $6x^2 - 5xy + y^2 = 0$
 (iv) $3x^2 - y^2 = 0$ (v) $xy - 2x - y + 2 = 0$ (vi) $xy - 2x - 3y + 6 = 0$
 (vii) $8x^2 + 2xy - 3y^2 + 12x + 14y - 8 = 0$ (viii) $2x^2 + 2xy - y^2 = 0$
 (ix) $x^2 - 81 = 0$ (x) $x^2 - 2xy - 2x + 6y - 3 = 0$ (xi) $2x^2 - 7xy + 3y^2 = 0$

- 3) (i) $2x - 3y = 0, 3x + 2y = 0$ (ii) $x - 2y = 0, x + 2y = 0$
 (iii) $\sqrt{3}x + y = 0, \sqrt{3}x - y = 0$ (iv) $(\sqrt{3} - 1)x + y = 0, (\sqrt{3} + 1)x - y = 0$
- 4) (i) $5x^2 + 4xy - y^2 = 0$ (ii) $9x^2 - 3xy - 2y^2 = 0$ (iii) $x^2 + xy - y^2 = 0$
- 5) (i) 0 (ii) -1 (iii) 1 (iv) 8 (v) 1 (vi) 6 (vii) 5
- 6) $3x^2 + 2xy - 3y^2 = 0$
- 7) $x^2 - 3y^2 = 0$
- 8) $\frac{50}{\sqrt{3}}$
- 10) $x^2 - 2xy - y^2 = 0$
- 11) -4
- 13) (i) 0° (ii) $\tan^{-1}(3)$ (iii) $\tan^{-1}(3)$
- 14) $x^2 - 3y^2 = 0$
- 18) Area = $\sqrt{3}$ sq. unit, Perimeter = 6 unit
- 22) $e = 0$ or $bd = ae$ 26) $a = 1, c = 0$.



5. Vectors

Exercise 5.1

- 1) 25
- 2) (i) $2\vec{a} - 2\vec{b}$ (ii) $\vec{a} + \vec{b}$ (iii) $\vec{b} - \vec{a}$
- 3) $\vec{OC} = 2\vec{a} + 2\vec{b}$, $\vec{OD} = -3\vec{a} + 2\vec{b}$, $\vec{OE} = -2\vec{a} + \vec{b}$
- 5) Vectors do not form a triangle.
- 6) $\vec{c} = \frac{1}{2}\vec{a} + \frac{1}{2}\vec{b}$. $\vec{d} = \frac{1}{2}\vec{b} - \frac{1}{2}\vec{a}$.
- 7) $\frac{7}{\sqrt{5}}\hat{i} - \frac{14}{\sqrt{5}}\hat{j}$
- 8) (a) 6 (b) 4 (c) 2 (d) $2\sqrt{10}$ (e) $2\sqrt{13}$ (f) $2\sqrt{5}$
- 9) (a) $x = -3, y = 4, z = 5$ (b) $(0, 1, 6)$
- 10) $\frac{\sqrt{3}}{2}$ sq. units
- 11) Terminal Point is $(3, 1, 7)$
- 12) $q = \frac{5}{2}$
- 13) Non coplanar
- 14) $\vec{r} = 2\vec{a} + 2\vec{b} - 3\vec{c}$

Exercise 5.2

- 1) (i) $\frac{1}{5}(-11, 4, -9)$ (ii) $(-19, 8, -21)$
- 2) $M(6, -1, 5)$
- 3) (i) C divides externally in the ratio 3:1. (ii) $p = 9, q = 2$.
- 6) 15 : 4 and 10 : 9 respectively
- 9) $C \equiv (-2, 0, 2)$
- 10) $OP : PD = 3 : 2$
- 11) $\sqrt{107}$
- 12) $G \equiv (4, -3, 2)$

Exercise 5.3

- 1) $\pm \left(\frac{2}{\sqrt{17}}\hat{i} + \frac{2}{\sqrt{17}}\hat{j} + \frac{3}{\sqrt{17}}\hat{k} \right)$
- 6) (i) Parallel (ii) Orthogonal (iii) Orthogonal (iv) Neither parallel nor orthogonal
- 7) $\angle P = 45^\circ$
- 8) (i) $\frac{1}{2}$ (ii) $\frac{1}{2}$

10) $\frac{\pi}{3}$

11) $0, \frac{-1}{\sqrt{2}}, \frac{1}{\sqrt{2}}$

12) $\frac{2}{11}, \frac{-6}{11}, \frac{9}{11}$

13) (0, 5, 7) or (8, -3, 3)

14) -1, 1, 2 or 1, 2, 3.



Exercise 5.4

1) $-4\hat{i} + 10\hat{j} + 22\hat{k}$

2) $\pm \left(\frac{2}{3}\hat{i} - \frac{2}{3}\hat{j} + \frac{1}{3}\hat{k} \right)$

3) 60°

4) $\pm \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$

5) (i) ± 6

ii) 1.6

7) $6i + 12j + 6k$

8) $\sqrt{146}$ sq. units

10) $\sqrt{42}$ sq. units

12) $\bar{b} = \frac{1}{3}(5\hat{i} + 2\hat{j} + 2\hat{k})$

13) $2\hat{j} + \hat{k}$

14) $\frac{3\pi}{4}$

16) i) -3, 5, 11

ii) 4, -4, 4

17) $\left(\frac{-8}{5}, \frac{16}{5}, \frac{24}{5} \right)$



Exercise 5.5

1) 110

2) 23 cubic units

3) $p = 2$

6) (i) -12

ii) 16

iii) $|\bar{u} + \bar{v}|^2$

7) $\frac{16}{3}$ cubic units

9) (i) $6\hat{i} + 3\hat{j} - 6\hat{k}$ (ii) $-2\hat{i} + 4\hat{j}$

Not same ; as $\vec{a} \times (\vec{b} \times \vec{c})$ lies in the plane of \vec{b} and \vec{c} whereas $(\vec{a} \times \vec{b}) \times \vec{c}$ lies in the plane of \vec{a} and \vec{b} .

Miscellaneous exercise - 5

I.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
C	B	B	B	A	D	C	A	B	B	A	B	A	A	A	B	C	B	A	A

II. 1) (i) $\vec{b} - \frac{1}{2} \vec{a}$ (ii) $\vec{b} - 3\vec{a}$ (iii) $\frac{3}{2} \vec{a} - \vec{b}$ (iv) $2\vec{a} - \vec{b}$

2) $-\frac{1}{2} \vec{a} - \frac{1}{2} \vec{b} + \vec{c}$

4) $\vec{AB} = -2\hat{i} + 5\hat{j} + \hat{k}$ and $\vec{AD} = 4\hat{i} - 2\hat{j} + 3\hat{k}$

5) 3

6) $\sqrt{2}$

7) (i) Right angled triangle ii) Isosceles triangle

8) (i) $2j \pm 2\sqrt{3} \hat{k}$ ii) $\pm 5\sqrt{2} \hat{i} + 5\sqrt{2} \hat{k}$

9) $\frac{1}{\sqrt{17}}(3i + 2j + 2k)$ and $\frac{1}{\sqrt{21}}(-i - 2j + 4k)$

11) $\pm \frac{1}{\sqrt{17}}(i + 4j)$

12) $\hat{i} + 4\hat{j} - 4\hat{k} = 1(2\hat{i} - \hat{j} + 3\hat{k}) + 2(\hat{i} - 2\hat{j} + 4\hat{k}) + 3(-\hat{i} + 3\hat{j} - 5\hat{k})$

14) $7(\hat{i} + \hat{j} + \hat{k})$

15) (-4, 9, 6)

20) OP : PD = 3 : 2

21) $3\hat{i} + 2\hat{k}$

22) $-\frac{3}{2}$

24) $\vec{a}_1 = 6\hat{i} + 2\hat{k}$ and $\vec{a}_2 = -\hat{i} - 2\hat{j} + 3\hat{k}$

25) $\pm \left(\frac{1}{\sqrt{3}} \hat{i} + \frac{1}{\sqrt{3}} \hat{j} + \frac{1}{\sqrt{3}} \hat{k} \right)$

26) $\cos \theta = \frac{7}{5\sqrt{2}}$

27) $\cos \alpha = \frac{2}{3}, \cos \beta = \frac{1}{3}$ and $\cos \gamma = \frac{2}{3}$ $\cos \alpha = \frac{1}{4}, \cos \beta = \cos \gamma = \frac{2}{3}$

28) $2\hat{i} - \hat{j}$

- 30) $\cos^{-1}\left(\frac{1}{6}\right)$
- 31) $\left(\frac{19}{9}, \frac{28}{9}, \frac{41}{9}\right)$
- 33) $\frac{bc\hat{i} + ac\hat{j} + ab\hat{k}}{\sqrt{b^2c^2 + a^2c^2 + a^2b^2}}$ and $\text{area} = \frac{1}{2}\sqrt{b^2c^2 + a^2c^2 + a^2b^2}$
- 34) a) meaningful, scalar b) meaningless c) meaningful, vector
 d) meaningless e) meaningless f) meaningful, scalar
 g) meaningless h) meaningful, vector i) meaningful, scalar
 j) meaningful scalar k) meaningless l) meaningless
- 36) (i) No ii) No iii) Yes
- 37) $\sqrt{286}$ sq. units.
- 40) $a = \pm \frac{1}{\sqrt{3}}$
- 41) $2a^3$ cu. units.
- 44) 2 cubic units, $\frac{1}{3}$ cubic units



6. Line and Plane



Exercise 6.1

- 1) $\vec{r} = (-2\hat{i} + \hat{j} + \hat{k}) + \lambda(4\hat{i} - \hat{j} + 2\hat{k})$
- 2) $\vec{r} = (3\hat{i} + 4\hat{j} - 7\hat{k}) + \lambda(3\hat{i} - 5\hat{j} + 8\hat{k})$
- 3) $\vec{r} = (5\hat{i} + 4\hat{j} + 3\hat{k}) + \lambda(-3\hat{i} + 4\hat{j} + 2\hat{k})$
- 4) $\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda(2\hat{i} + \hat{j} - 3\hat{k})$
- 5) $\vec{r} = (-\hat{i} - \hat{j} + 2\hat{k}) + \lambda(3\hat{i} + 2\hat{j} + \hat{k})$
- 6) $\frac{x+1}{2} = \frac{y-2}{3} = \frac{z-1}{1}$
- 7) $\frac{x-2}{-1} = \frac{y-2}{1} = \frac{z-1}{-1}$
- 8) $\frac{x+2}{3} = \frac{y-3}{-2} = \frac{z-4}{-2}$
- 9) $(-11, -4, 5)$
- 10) $\vec{r} = (3\hat{i} - \hat{j} + 2\hat{k}) + \lambda(-2\hat{i} - 3\hat{j} - 2\hat{k})$



Exercise 6.2

- 1) $\sqrt{35}$
- 2) $(1, 2, 3), \sqrt{14}$
- 3) $\frac{1}{\sqrt{3}}$
- 4) $2\sqrt{29}$
- 5) $2\sqrt{6}, (3, -4, -2)$
- 6) $\left(\frac{99}{53}, \frac{-187}{53}, \frac{95}{53}\right)$
- 7) a) do not intersect b) do not intersect
- 8) $\frac{9}{2}$

Miscellaneous exercise - 6A

- 1) $\vec{r} = (3\hat{i} + 4\hat{j} - 7\hat{k}) + \lambda(6\hat{i} - \hat{j} + \hat{k})$
- 2) $\vec{r} = (3\hat{i} + 2\hat{j} + \hat{k}) + \lambda(2\hat{i} + 2\hat{j} - 3\hat{k})$
- 3) $\vec{r} = (-2\hat{i} + 4\hat{j} - 5\hat{k}) + \lambda(3\hat{i} + 5\hat{j} + 6\hat{k})$
- 4) $\vec{r} = (-5\hat{i} - 4\hat{j} - 5\hat{k}) + \lambda(3\hat{i} + 5\hat{j} + 6\hat{k})$
- 5) $\vec{r} = \lambda(5\hat{i} - 2\hat{j} + 3\hat{k})$
- 6) $x = 3, y = -2$
- 7) $\frac{x-3}{-2} = \frac{y-2}{1}; z = 1$
- 8) $x - 1 = y - 1 = z - 2$
- 9) $\frac{x-2}{2} = \frac{y-1}{-7} = \frac{z-3}{4}$
- 10) $\vec{r} = \lambda(-\hat{i} + \hat{k})$
- 11) $-\frac{10}{11}$
- 12) 60°
- 13) 45°
- 14) 45°
- 15) $(2, 3, -1)$
- 16) i) intersect ii) intersect

- 17) -1
- 18) $\frac{x+1}{3} = \frac{y+1}{2} = \frac{z-2}{1}$, $\vec{r} = (-\hat{i} - \hat{j} + 2\hat{k}) + \lambda(3\hat{i} + 2\hat{j} + \hat{k})$
- 19) $\frac{2}{\sqrt{13}}, \frac{3}{\sqrt{13}}, 0$
- 20) $\frac{x}{7} = \frac{y}{-12} = \frac{z}{5}$
- 21) $\vec{r} = \left(2\hat{j} + \frac{5}{3}\hat{k}\right) + \lambda(3\hat{i} + 4\hat{k})$
- 22) $(2, 0, 5), (0, 4, 1)$



Exercise 6.3

- 1) $\vec{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) = 126$
- 2) 1
- 3) $\left(\frac{18}{7}, \frac{54}{7}, \frac{-27}{7}\right)$
- 4) $\vec{r} \cdot \left(\frac{3}{13}\hat{i} + \frac{4}{13}\hat{j} + \frac{12}{13}\hat{k}\right) = 6$, (i) 6 (ii) $\left(\frac{3}{13}, \frac{4}{13}, \frac{12}{13}\right)$
- 5) $\vec{r} \cdot (4\hat{i} + 5\hat{j} + 6\hat{k}) = 15$
- 6) $2y + 5z = 19$
- 7) $z = 6$
- 8) $\vec{r} \cdot (\hat{i}) = 1$
- 9) $\vec{r} \cdot (-4\hat{i} - \hat{j} + 5\hat{k}) = 26$
- 10) $5x - 2y - 3z = 38$
- 11) $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 1$



Exercise 6.4

- 1) 60°
- 2) $\sin^{-1}\left(\frac{5}{7\sqrt{6}}\right)$
- 3) $\vec{r} \cdot (-\hat{i} + 2\hat{j} - \hat{k}) = 7$
- 4) 2
- 5) $\frac{1}{13}$

Miscellaneous exercise - 6B

I.

1	2	3	4	5	6	7	8	9	10
B	A	A	C	D	B	B	C	D	B

11	12	13	14	15	16	17	18	19	20
A	D	D	D	A	A	B	B	A	B

- II.
- 1) $\vec{r} \cdot (2\hat{i} + \hat{j} + 2\hat{k}) = 15$
 - 2) 1
 - 3) (2, 3, 6)
 - 4) i) $\frac{1}{2}$ ii) $\frac{3}{13}, \frac{4}{13}, \frac{12}{13}$
 - 5) $\vec{r} \cdot (16\hat{i} + 4\hat{k}) = 20$
 - 6) $y + 2 = 0$
 - 7) $6x + 8y + 7z = 148$
 - 8) $\vec{r} \cdot (\hat{i} + 2\hat{j}) = 5$
 - 9) $\vec{r} \cdot (bc\hat{i} + ca\hat{j} + ab\hat{k}) = abc$
 - 10) $\vec{r} \cdot (-3\hat{i} + 3\hat{j} + 4\hat{k}) = 35$
 - 11) $\vec{r} \cdot (5\hat{i} - 4\hat{j} + \hat{k}) = 0$
 - 12) $x + y + z = 6, x - 2y + z = 0$
 - 13) $x + y + z = 3$
 - 14) 90°
 - 15) $\sin^{-1}\left(\frac{2\sqrt{2}}{3}\right)$
 - 16) $\vec{r} \cdot (-\hat{i} + 2\hat{j} - \hat{k}) = 7$
 - 17) 0 unit
 - 18) 19 units
 - 19) $\vec{r} \cdot (\hat{i} - \hat{k}) = 0$
 - 20) $\vec{r} \cdot (\hat{i} - 4\hat{k}) = -5$
 - 21) $\vec{r} \cdot (\hat{k}) = 0$



7. Linear Programming



Exercise 7.3

- 1) maximize $z = 30x + 20y$ subject to $10x + 6y \leq 60, 5x + 4y \leq 35, x \geq 0, y \geq 0$
- 2) maximize $z = 3x + 2y$ subject to $2x + y \geq 14, 2x + 3y \geq 22, x + y \geq 1, x \geq 0, y \geq 0$
- 3) maximize $p = 350x + 400y$ subject to $3x + 2y \leq 120, 2x + 5y \leq 160, x \geq 0, y \geq 0$
- 4) maximize $z = 10x + 15y$ subject to $2x + 3y \leq 36, 5x + 2y \leq 50, 2x + 6y \leq 60, x \geq 0, y \geq 0$
- 5) maximize $p = 13.5x + 55y$ subject to $x + 2y \leq 10, 3x + 4y \leq 12, x \geq 0, y \geq 0$
- 6) maximize $z = 500x + 750y$ subject to $2x + 3y \leq 40, x + 4y \leq 70, x \geq 0, y \geq 0$
- 7) minimize $z = 4.5x + 3.5y$ subject to $4x + 6y \geq 18, 14x + 12y \geq 28, 7x + 8y \geq 14, x \geq 0, y \geq 0$
- 8) maximize $z = x_1 + x_2$ subject to $\frac{x_1}{60} + \frac{x_2}{90} \leq 1, 5x_1 + 8x_2 \leq 600, x \geq 0, x_2 \geq 0$
- 9) minimize $C = 20x_1 + 6x_2$ s. t $x_1 > 4, x_2 < 2, x_1 + x_2 \geq 5, x \geq 0, x_2 \geq 0$.



Exercise 7.4

- 1) Maximum at (4, 2), 60
- 2) Maximum at (0, 6), maximum value = 36
- 3) Maximum at (4.5, 2.5), 59
- 4) Maximum at (2, 3), maximum value = 95
- 5) Maximum at (4, 5), maximum $z = 37$
- 6) Maximum at (0, 5), 5
- 7) Maximum at (1.5, 4), 52
- 8) Maximum at (2, 0.5), 22.5

Miscellaneous exercise - 7


I.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
A	C	B	C	A	D	C	B	A	B	B	B	A	C	C

- 5)
 - (i) $x_1 = 4.5, x_2 = 3$ $\max z = 40.5$.
 - (ii) $x = 3, y = 18$ $\min z = 48$.
 - (iii) infinite number of optimum solutions on the line $3x + 5y = 10$ between $A\left(\frac{45}{16}, \frac{5}{16}\right)$ and $B(0, 2)$.

- 6) (i) $x = 4, y = 3$ maximize $z = 25$.
 (ii) $x = 10, y = 15$ maximize $z = 1350$.
 (iii) $x = 3, y = 18$ maximize $z = 48$.
- 7) maximize $z = 140x + 210y$ s.t. $3x + 3y \leq 36, 5x + 2y \leq 50, 2x + 6y \leq 60$
 $x, y \geq 0$ where $x =$ no. of tables = 3
 $y =$ no. of chairs = 9
 maximize $z =$ maximum profit = 2310/-
- 8) Maximize $z = 180x + 220y$ s.t. $6x + 4y \leq 120, 3x + 10y \leq 180, x \geq 0, y \geq 0$.
 Ans. $x = 10, y = 15$.
- 9) Minimize $z = 4x + 6y$ s.t. $x + 2y \geq 80, 3x + y \geq 75, x \geq 0, y \geq 0$.
 Ans. $x = 14, y = 33$.
- 10) Maximize $z = 2000x + 3000y$ s.t. $3x + 3y \leq 36, 5x + 2y \leq 50, 2x + 6y \leq 60, x \geq 0, y \geq 0$.
 Ans. $x = 3, y = 9$.
- 11) Minimize $z = 800x + 640y$ s.t. $4x + 2y \geq 16, 12x + 2y \geq 24, 2x + 6y \geq 18, x \geq 0, y \geq 0$.
 Ans. Minimum cost ₹3680/- when $x = 3, y = 2$.
- 12) Maximize $z = 75x + 125y$ s.t. $4x + 2y \leq 208, 2x + 4y \leq 152, x \geq 0, y \geq 0$.
 Ans. $x = 44, y = 16$.
- 13) Maximize $z = -3x + 4y$ s.t. $x + y \leq 450, 2x + y \leq 600, x \geq 0, y \geq 0$
 maximum profit = Rs. 1800/- at $(0, 450)$
- 14) Maximize $z = 20x + 30y$ s.t. $2x + 2y \leq 210, 3x + 4y \leq 300, x \geq 0, y \geq 0$
 maximum profit = Rs. 2400/- at $(30, 60)$





**Maharashtra State Bureau of Textbook
Production and Curriculum Research,
Pune - 411 004.**

गणित आणि संख्याशास्त्र (कला व विज्ञान) भाग-१, इ. १२ वी (इंग्रजी माध्यम)



₹ 146.00



Mathematics & Statistics

Arts & Science Part 2

STANDARD XII



The Coordination Committee formed by GR No. Abhyas - 2116/(Pra.Kra.43/16) SD - 4
Dated 25.4.2016 has given approval to prescribe this textbook in its meeting held on
30.01.2020 and it has been decided to implement it from the educational year 2020-21.

Mathematics and Statistics

(Arts and Science)

Part - II

STANDARD - XII



**Maharashtra State Bureau of Textbook Production and Curriculum Research,
Pune - 411 004**



B6K9J5

Download DIKSHA App on your smartphone. If you scan the Q.R.Code on this page of your textbook, you will be able to access full text and the audio-visual study material relevant to each lesson provided as teaching and learning aids.

First Edition : 2020 © Maharashtra State Bureau of Textbook Production and Curriculum Research, Pune- 411 004.

Maharashtra State Bureau of Textbook Production and Curriculum Research reserves all rights relating to the book. No part of this book should be reproduced without the written permission of the Director, Maharashtra State Bureau of Textbook Production and curriculum Research, Pune.

**Mathematics and Statistics
(Arts and Science)
Committee Members**

Dr. Mangla Narlikar	(Chairman)
Dr. Sharad Gore	(Member)
Shri. Prahallad Chippalagatti	(Member)
Shri. Prasad Kunte	(Member)
Shri. Sujit Shinde	(Member)
Smt. Prajkti Gokhale	(Member)
Shri. Ramakant Sarode	(Member)
Smt. Ujjwala Godbole	(Member-Secretary)

Co-ordinator

Ujjwala Shrikant Godbole
I/C Special Officer for Mathematics

**Cover &
Computer Drawings**

Shri. Sandip Koli, Artist, Mumbai

Production

Sachchitanand Aphale
Chief Production Officer
Sanjay Kamble
Production Officer
Prashant Harne
Asst. Production Officer

**Mathematics and Statistics
(Arts and Science)
Study Group Members**

Dr. Ishwar Patil	Dr. Pradeep Mugale
Dr. Pradnyankumar Bhojekar	Shri. Uday Mahindrakar
Shri. Prafullachandra Pawar	Shri. Balkrishna Mapari
Shri. Sachin Batwal	Shri. Devanand Bagul
Shri. Swapnil Shinde	Smt. Swati Powar
Shri. Pradipkumar Bhavsar	Smt. Mahadevi Mane
Shri. Amit Totade	Shri. Dilip Panchamia
Smt. Gauri Prachand	Shri. Pramod Deshpande
Shri. Sharadchandra Walagade	Shri. Milind Patil
Shri. Dhananjay Panhalkar	Smt. Supriya Abhyankar
Smt. Deepti Kharapas	Smt. Nileema Khaladkar

Paper

70 GSM Cream wove

Print Order No.

N/PB/2020-21/75,000

Printer


SOHAIL ENTERPRISES, THANE

Layout & Typesetters

Samarth Graphics,
522, Narayan Peth, Pune 411 030.

Publisher

Vivek Uttam Gosavi, Controller
Maharashtra State Textbook Bureau,
Prabhadevi Mumbai- 400 025



NATIONAL ANTHEM

Jana-gana-mana-adhināyaka jaya hē
Bhārata-bhāgya-vidhātā,

Panjāba-Sindhu-Gujarāta-Marāthā
Drāvida-Utkala-Banga

Vindhya-Himāchala-Yamunā-Gangā
uchchala-jaladhi-taranga

Tava subha nāmē jāgē, tava subha āsisa māgē,
gāhē tava jaya-gāthā,

Jana-gana-mangala-dāyaka jaya hē
Bhārata-bhāgya-vidhātā,

Jaya hē, Jaya hē, Jaya hē,
Jaya jaya jaya, jaya hē.

PLEDGE

India is my country. All Indians
are my brothers and sisters.

I love my country, and I am proud
of its rich and varied heritage. I shall
always strive to be worthy of it.

I shall give my parents, teachers
and all elders respect, and treat
everyone with courtesy.

To my country and my people,
I pledge my devotion. In their
well-being and prosperity alone lies
my happiness.

PREFACE

Dear Students,

Welcome to Standard XII, an important milestone in your life.

Standard XII or Higher Secondary School Certificate opens the doors of higher education. Alternatively, you can pursue other career paths like joining the workforce. Either way, you will find that mathematics education helps you considerably. Learning mathematics enables you to think logically, consistently, and rationally. The curriculum for Standard XII Mathematics and Statistics for Science and Arts students has been designed and developed keeping both of these possibilities in mind.

The curriculum of Mathematics and Statistics for Standard XII for Science and Arts students is divided in two parts. Part I deals with topics like Mathematical Logic, Matrices, Vectors and Introduction to three dimensional geometry. Part II deals with Differentiation, Integration and their applications, Introduction to random variables and statistical methods.

The new text books have three types of exercises for focused and comprehensive practice. First, there are exercises on every important topic. Second, there are comprehensive exercises at the end of all chapters. Third, every chapter includes activities that students must attempt after discussion with classmates and teachers. Additional information has been provided on the E-balbharati website (www.ebalbharati.in).

We are living in the age of Internet. You can make use of modern technology with the help of the Q.R. code given on the title page. The Q.R. code will take you to links that provide additional useful information. Your learning will be fruitful if you balance between reading the text books and solving exercises. Solving more problems will make you more confident and efficient.

The text books are prepared by a subject committee and a study group. The books (Paper I and Paper II) are reviewed by experienced teachers and eminent scholars. The Bureau would like to thank all of them for their valuable contribution in the form of creative writing, constructive and useful suggestions for making the text books valuable. The Bureau hopes and wishes that the text books are very useful and well received by students, teachers and parents.

Students, you are now ready to study. All the best wishes for a happy learning experience and a well deserved success. Enjoy learning and be successful.



(Vivek Gosavi)
Director

Pune

Date : 21 February 2020

Bharatiya Saur : 2 Phalguna 1941

Maharashtra State Bureau of Textbook
Production and Curriculum Research, Pune.

Mathematics and Statistics XII (Part II)
Arts and Science

Sr. No	Area / Topic	Sub Unit	Competency Statement
1.	Differentiation	Differentiation	<p>The students will be able to</p> <ul style="list-style-type: none"> • state and use standard formulas of derivative of standard functions • use chain rule of derivatives • find derivatives of the logarithm, implicit, inverse and parametric functions • find second and higher order derivatives.
2.	Applications of Derivatives	Applications of Derivatives	<ul style="list-style-type: none"> • find equations of tangents and normal to a curve • determine nature of the function-increasing or decreasing • find approximate values of the function • examine function for maximum and minimum values • verify mean value theorems
3.	Indefinite Integration	Indefinite Integration	<ul style="list-style-type: none"> • understand the relation between derivative and integral • use the method of substitution • solve integrals with the help of integration by parts • solve the integrals by the method of partial fractions
4.	Definite Integration	Definite Integration	<ul style="list-style-type: none"> • understand integral as a limit of sum • the properties of definite integral • state the properties of definite integral and use them to solve problems

5.	Application of Definite Integration	Application of Definite Integration	<ul style="list-style-type: none"> • find the area under the curve, bounded by the curves using definite integrals.
6.	Differential Equation	Differential Equation	<ul style="list-style-type: none"> • form a differential equation and find its order and degree • solve the first order and first degree differential equation by various methods • apply the differential equations to study the population, growth and decay in amount of substance and physics.
7.	Probability Distribution	Probability Distribution	<ul style="list-style-type: none"> • understand the random variable and its types. • find probability mass function and its probability distribution. • find the expected value, variance and the standard deviation • find the probability density function of continuous random variable • find distribution function of c.r.v.
8	Binomial Distribution	Binomial Distribution	<ul style="list-style-type: none"> • understand random experiment with two or more outcomes. • determine probability distribution of random experiment with parameters n and p. • find mean, variance, expected value and standard deviation for the binomial distribution.

INDEX

Sr. No.	Chapter Name	Page No.
1	Differentiation	1-64
2	Applications of Derivatives	65-94
3	Indefinite Integration	95-150
4	Definite Integration	151-177
5	Application of Definite Integration	178-190
6	Differential Equations	191-218
7	Probability Distributions	219-244
8	Binomial Distribution	245-255
	Answers	256-276

1. DIFFERENTIATION

Let us Study

- Derivatives of Composite functions.
- Derivatives of Inverse functions
- Derivatives of Implicit functions.
- Higher order Derivatives.
- Geometrical meaning of Derivative.
- Logarithmic Differentiation
- Derivatives of Parametric functions.

Let us Recall

- The derivative of $f(x)$ with respect to x , at $x = a$ is given by $f'(a) = \lim_{h \rightarrow 0} \left[\frac{f(a+h) - f(a)}{h} \right]$
- The derivative can also be defined for $f(x)$ at any point x on the open interval as $f'(x) = \lim_{h \rightarrow 0} \left[\frac{f(x+h) - f(x)}{h} \right]$. If the function is given as $y = f(x)$ then its derivative is written as $\frac{dy}{dx} = f'(x)$.
- For a differentiable function $y = f(x)$ if δx is a small increment in x and the corresponding increment in y is δy then $\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \frac{dy}{dx}$.
- Derivatives of some standard functions.

$y = f(x)$	$\frac{dy}{dx} = f'(x)$
c (Constant)	0
x^n	nx^{n-1}
$\frac{1}{x}$	$-\frac{1}{x^2}$
$\frac{1}{x^n}$	$-\frac{n}{x^{n+1}}$
\sqrt{x}	$\frac{1}{2\sqrt{x}}$
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
$\tan x$	$\sec^2 x$

$y = f(x)$	$\frac{dy}{dx} = f'(x)$
$\sec x$	$\sec x \tan x$
$\operatorname{cosec} x$	$-\operatorname{cosec} x \cot x$
$\cot x$	$-\operatorname{cosec}^2 x$
e^x	e^x
a^x	$a^x \log a$
$\log x$	$\frac{1}{x}$
$\log_a x$	$\frac{1}{x \log a}$

Table 1.1.1

Rules of Differentiation :

If u and v are differentiable functions of x such that

$$(i) \quad y = u \pm v \quad \text{then} \quad \frac{dy}{dx} = \frac{du}{dx} \pm \frac{dv}{dx} \qquad (ii) \quad y = uv \quad \text{then} \quad \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$(iii) \quad y = \frac{u}{v} \quad \text{where } v \neq 0 \quad \text{then} \quad \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Introduction :

The history of mathematics presents the development of calculus as being accredited to Sir Isaac Newton (1642-1727) an English physicist and mathematician and Gottfried Wilhelm Leibnitz (1646-1716) a German physicist and mathematician. The Derivative is one of the fundamental ideas of calculus. It's all about rate of change in a function. We try to find interpretations of these changes in a mathematical way. The symbol δ will be used to represent the change, for example δx represents a small change in the variable x and it is read as "change in x " or "increment in x ". δy is the corresponding change in y if y is a function of x .

We have already studied the basic concept, derivatives of standard functions and rules of differentiation in previous standard. This year, in this chapter we are going to study the geometrical meaning of derivative, derivatives of Composite, Inverse, Logarithmic, Implicit and Parametric functions and also higher order derivatives. We also add some more rules of differentiation.

**Let us Learn****1.1.1 Derivatives of Composite Functions (Function of another function) :**

So far we have studied the derivatives of simple functions like $\sin x$, $\log x$, e^x etc. But how about the derivatives of $\sin \sqrt{x}$, $\log(\sin(x^2 + 5))$ or $e^{\tan x}$ etc? These are known as composite functions. In this section let us study how to differentiate composite functions.


1.1.2 Theorem : If $y = f(u)$ is a differentiable function of u and $u = g(x)$ is a differentiable function of x such that the composite function $y = f[g(x)]$ is a differentiable function of x then $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$.

Proof : Given that $y = f(u)$ and $u = g(x)$. We assume that u is not a constant function. Let there be a small increment in the value of x say δx then δu and δy are the corresponding increments in u and y respectively.

As δx , δu , δy are small increments in x , u and y respectively such that $\delta x \neq 0$, $\delta u \neq 0$ and $\delta y \neq 0$.

$$\text{We have } \frac{\delta y}{\delta x} = \frac{\delta y}{\delta u} \times \frac{\delta u}{\delta x}.$$

Taking the limit as $\delta x \rightarrow 0$ on both sides we get,



$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta u} \right) \times \lim_{\delta x \rightarrow 0} \left(\frac{\delta u}{\delta x} \right)$$

As $\delta x \rightarrow 0$, we get, $\delta u \rightarrow 0$ ($\because u$ is a continuous function of x)

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \lim_{\delta u \rightarrow 0} \left(\frac{\delta y}{\delta u} \right) \times \lim_{\delta x \rightarrow 0} \left(\frac{\delta u}{\delta x} \right) \quad \dots \dots \text{(I)}$$

Since y is a differentiable function of u and u is a differentiable function of x . we have,

$$\lim_{\delta u \rightarrow 0} \left(\frac{\delta y}{\delta u} \right) = \frac{dy}{du} \quad \text{and} \quad \lim_{\delta x \rightarrow 0} \left(\frac{\delta u}{\delta x} \right) = \frac{du}{dx} \quad \dots \dots \text{(II)}$$

From (I) and (II), we get

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \frac{dy}{du} \times \frac{du}{dx} \quad \dots \dots \text{(III)}$$

The R.H.S. of (III) exists and is finite, implies L.H.S. of (III) also exists and is finite

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \frac{dy}{dx}. \quad \text{Then equation (III) becomes,}$$

$$\boxed{\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}}$$

Note:

1. The derivative of a composite function can also be expressed as follows. $y = f(u)$ is a differentiable function of u and $u = g(x)$ is a differentiable function of x such that the composite function $y = f[g(x)]$ is defined then

$$\frac{dy}{dx} = f'[g(x)] \cdot g'(x).$$

2. If $y = f(v)$ is a differentiable function of v and $v = g(u)$ is a differentiable function of u and $u = h(x)$ is a differentiable function of x then

$$\frac{dy}{dx} = \frac{dy}{dv} \times \frac{dv}{du} \times \frac{du}{dx}.$$

3. If y is a differentiable function of u_1 , u_i is a differentiable function of u_{i+1} for $i = 1, 2, \dots, n-1$ and u_n is a differentiable function of x , then

$$\frac{dy}{dx} = \frac{dy}{du_1} \times \frac{du_1}{du_2} \times \frac{du_2}{du_3} \times \dots \times \frac{du_{n-1}}{du_n} \times \frac{du_n}{dx}$$

This rule is also known as **Chain rule**.



1.1.3 Derivatives of some standard Composite Functions :

y	$\frac{dy}{dx}$
$[f(x)]^n$	$n [f(x)]^{n-1} \cdot f'(x)$
$\sqrt{f(x)}$	$\frac{f'(x)}{2\sqrt{f(x)}}$
$\frac{1}{[f(x)]^n}$	$-\frac{n \cdot f'(x)}{[f(x)]^{n+1}}$
$\sin [f(x)]$	$\cos [f(x)] \cdot f'(x)$
$\cos [f(x)]$	$-\sin [f(x)] \cdot f'(x)$
$\tan [f(x)]$	$\sec^2 [f(x)] \cdot f'(x)$
$\sec [f(x)]$	$\sec [f(x)] \cdot \tan [f(x)] \cdot f'(x)$

y	$\frac{dy}{dx}$
$\cot [f(x)]$	$-\operatorname{cosec}^2 [f(x)] \cdot f'(x)$
$\operatorname{cosec} [f(x)]$	$-\operatorname{cosec} [f(x)] \cdot \cot [f(x)] \cdot f'(x)$
$a^{f(x)}$	$a^{f(x)} \cdot \log a \cdot f'(x)$
$e^{f(x)}$	$e^{f(x)} \cdot f'(x)$
$\log [f(x)]$	$\frac{f'(x)}{f(x)}$
$\log_a [f(x)]$	$\frac{f'(x)}{f(x) \log a}$

Table 1.1.2



SOLVED EXAMPLES

Ex. 1 : Differentiate the following w. r. t. x .

(i) $y = \sqrt{x^2 + 5}$

(ii) $y = \sin (\log x)$

(iii) $y = e^{\tan x}$

(iv) $\log (x^5 + 4)$

(v) $5^{3 \cos x - 2}$

(vi) $y = \frac{3}{(2x^2 - 7)^5}$

Solution : (i) $y = \sqrt{x^2 + 5}$

Method 1 :

Let $u = x^2 + 5$ then $y = \sqrt{u}$, where y is a differentiable function of u and u is a differentiable function of x then

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} \quad \dots \dots (I)$$

Now, $y = \sqrt{u}$

Differentiate w. r. t. u

$$\frac{dy}{du} = \frac{d}{du}(\sqrt{u}) = \frac{1}{2\sqrt{u}} \text{ and } u = x^2 + 5$$

Differentiate w. r. t. x

$$\frac{du}{dx} = \frac{d}{dx}(x^2 + 5) = 2x$$

Now, equation (I) becomes,

$$\frac{dy}{dx} = \frac{1}{2\sqrt{u}} \times 2x = \frac{x}{\sqrt{x^2 + 5}}$$

Method 2 :

We have $y = \sqrt{x^2 + 5}$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx}(\sqrt{x^2 + 5})$$

[Treat $x^2 + 5$ as u in mind and use the formula of derivative of \sqrt{u}]

$$\frac{dy}{dx} = \frac{1}{2\sqrt{x^2 + 5}} \cdot \frac{d}{dx}(x^2 + 5)$$

$$\frac{dy}{dx} = \frac{1}{2\sqrt{x^2 + 5}}(2x)$$

$$\frac{dy}{dx} = \frac{x}{\sqrt{x^2 + 5}}$$

(ii) $y = \sin(\log x)$

Method 1 :

Let $u = \log x$ then $y = \sin u$, where y is a differentiable function of u and u is a differentiable function of x then

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} \quad \dots\dots (I)$$

Now, $y = \sin u$

Differentiate w. r. t. u

$$\frac{dy}{du} = \frac{d}{du}(\sin u) = \cos u \text{ and } u = \log x$$

Differentiate w. r. t. x

$$\frac{du}{dx} = \frac{d}{dx}(\log x) = \frac{1}{x}$$

Now, equation (I) becomes,

$$\frac{dy}{dx} = \cos u \times \frac{1}{x} = \frac{\cos(\log x)}{x}$$

Note : Hence onwards let's use Method 2.

(iii) $y = e^{\tan x}$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx}[e^{\tan x}]$$

$$\frac{dy}{dx} = e^{\tan x} \times \frac{d}{dx}(\tan x)$$

$$\frac{dy}{dx} = e^{\tan x} \cdot \sec^2 x = \sec^2 x \cdot e^{\tan x}$$

(v) Let $y = 5^{3 \cos x - 2}$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx}[5^{3 \cos x - 2}]$$

$$\frac{dy}{dx} = 5^{3 \cos x - 2} \cdot \log 5 \times \frac{d}{dx}(3 \cos x - 2)$$

$$\frac{dy}{dx} = -3 \sin x \cdot 5^{3 \cos x - 2} \cdot \log 5$$

Method 2 :

We have $y = \sin(\log x)$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx}[\sin(\log x)]$$

[Treat $\log x$ as u in mind and use the formula of derivative of $\sin u$]

$$\frac{dy}{dx} = \cos(\log x) \times \frac{d}{dx}(\log x)$$

$$\frac{dy}{dx} = \cos(\log x) \cdot \frac{1}{x}$$

$$\frac{dy}{dx} = \frac{\cos(\log x)}{x}$$

(iv) Let $y = \log(x^5 + 4)$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx}[\log(x^5 + 4)]$$

$$\frac{dy}{dx} = \frac{1}{x^5 + 4} \times \frac{d}{dx}(x^5 + 4)$$

$$\frac{dy}{dx} = \frac{1}{x^5 + 4} (5x^4) = \frac{5x^4}{x^5 + 4}$$

(vi) Let $y = \frac{3}{(2x^2 - 7)^5}$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} \left(\frac{3}{(2x^2 - 7)^5} \right) = 3 \frac{d}{dx} \left(\frac{1}{(2x^2 - 7)^5} \right)$$

$$= 3 \times \frac{-5}{(2x^2 - 7)^6} \times \frac{d}{dx}(2x^2 - 7)$$

$$= -\frac{15}{(2x^2 - 7)^6} (4x)$$

$$\frac{dy}{dx} = -\frac{60x}{(2x^2 - 7)^6}$$

Ex. 2 : Differentiate the following w. r. t. x.

- (i) $y = \sqrt{\sin x^3}$ (ii) $y = \cot^2(x^3)$ (iii) $y = \log [\cos(x^5)]$
 (iv) $y = (x^3 + 2x - 3)^4(x + \cos x)^3$ (v) $y = (1 + \cos^2 x)^4 \times \sqrt{x + \sqrt{\tan x}}$

Solution :

(i) $y = \sqrt{\sin x^3}$

Differentiate w. r. t. x

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\sqrt{\sin x^3}) \\ &= \frac{1}{2\sqrt{\sin x^3}} \times \frac{d}{dx}(\sin x^3) \\ &= \frac{1}{2\sqrt{\sin x^3}} \times \cos x^3 \times \frac{d}{dx}(x^3) \\ &= \frac{1}{2\sqrt{\sin x^3}} \times \cos x^3 \times (3x^2) \\ \therefore \frac{dy}{dx} &= \frac{3x^2 \cos x^3}{2\sqrt{\sin x^3}} \end{aligned}$$

(ii) $y = \cot^2(x^3)$

Differentiate w. r. t. x

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\cot^2(x^3)) \\ &= \frac{d}{dx}[\cot(x^3)]^2 \\ &= 2 \cot(x^3) \frac{d}{dx}[\cot(x^3)] \\ &= 2 \cot(x^3)[- \operatorname{cosec}^2(x^3)] \frac{d}{dx}(x^3) \\ &= -2 \cot(x^3) \operatorname{cosec}^2(x^3)(3x^2) \\ \therefore \frac{dy}{dx} &= -6x^2 \cot(x^3) \operatorname{cosec}^2(x^3) \end{aligned}$$

(iii) $y = \log [\cos(x^5)]$

Differentiate w. r. t. x

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\log [\cos(x^5)]) \\ &= \frac{1}{\cos(x^5)} \cdot \frac{d}{dx}[\cos(x^5)] \\ &= \frac{1}{\cos(x^5)} (-\sin(x^5)) \frac{d}{dx}(x^5) \end{aligned}$$

$\therefore \frac{dy}{dx} = -\tan(x^5)(5x^4) = -5x^4 \tan(x^5)$

(iv) $y = (x^3 + 2x - 3)^4(x + \cos x)^3$

Differentiate w. r. t. x

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}[(x^3 + 2x - 3)^4(x + \cos x)^3] \\ &= (x^3 + 2x - 3)^4 \cdot \frac{d}{dx}(x + \cos x)^3 + (x + \cos x)^3 \cdot \frac{d}{dx}(x^3 + 2x - 3)^4 \end{aligned}$$



$$= (x^3 + 2x - 3)^4 \cdot 3(x + \cos x)^2 \cdot \frac{d}{dx}(x + \cos x) + (x + \cos x)^3 \cdot 4(x^3 + 2x - 3)^3 \cdot \frac{d}{dx}(x^3 + 2x - 3)$$

$$= (x^3 + 2x - 3)^4 \cdot 3(x + \cos x)^2(1 - \sin x) + (x + \cos x)^3 \cdot 4(x^3 + 2x - 3)^3(3x^2 + 2)$$

$$\therefore \frac{dy}{dx} = 3(x^3 + 2x - 3)^4(x + \cos x)^2(1 - \sin x) + 4(3x^2 + 2)(x^3 + 2x - 3)^3(x + \cos x)^3$$

(v) $y = (1 + \cos^2 x)^4 \times \sqrt{x + \sqrt{\tan x}}$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} \left[(1 + \cos^2 x)^4 \times \sqrt{x + \sqrt{\tan x}} \right]$$

$$= (1 + \cos^2 x)^4 \frac{d}{dx}(\sqrt{x + \sqrt{\tan x}}) + (\sqrt{x + \sqrt{\tan x}}) \frac{d}{dx}(1 + \cos^2 x)^4$$

$$= (1 + \cos^2 x)^4 \cdot \frac{1}{2\sqrt{x + \sqrt{\tan x}}} \cdot \frac{d}{dx}(x + \sqrt{\tan x}) + (\sqrt{x + \sqrt{\tan x}}) \cdot 4(1 + \cos^2 x)^3 \frac{d}{dx}[1 + (\cos x)^2]$$

$$= (1 + \cos^2 x)^4 \cdot \frac{1}{2\sqrt{x + \sqrt{\tan x}}} \left(1 + \frac{1}{2\sqrt{\tan x}} \cdot \frac{d}{dx}(\tan x) \right) + (\sqrt{x + \sqrt{\tan x}}) \cdot 4(1 + \cos^2 x)^3 (2 \cos x) \frac{d}{dx}(\cos x)$$

$$= (1 + \cos^2 x)^4 \cdot \frac{1}{2\sqrt{x + \sqrt{\tan x}}} \left(1 + \frac{\sec^2 x}{2\sqrt{\tan x}} \right) + (\sqrt{x + \sqrt{\tan x}}) \cdot 4(1 + \cos^2 x)^3 (2 \cos x)(-\sin x)$$

$$= (1 + \cos^2 x)^4 \cdot \frac{1}{2\sqrt{x + \sqrt{\tan x}}} \left(\frac{2\sqrt{\tan x} + \sec^2 x}{2\sqrt{\tan x}} \right) - (\sqrt{x + \sqrt{\tan x}}) \cdot 4(1 + \cos^2 x)^3 (2 \sin x \cos x)$$

$$\frac{dy}{dx} = \frac{(1 + \cos^2 x)^4 (2\sqrt{\tan x} + \sec^2 x)}{4\sqrt{\tan x} \sqrt{x + \sqrt{\tan x}}} - 4 \sin 2x (1 + \cos^2 x)^3 \sqrt{x + \sqrt{\tan x}}$$

Ex. 3 : Differentiate the following w. r. t. x .

(i) $y = \log_3(\log_5 x)$ (ii) $y = \log \left[e^{3x} \cdot \frac{(3x - 4)^{\frac{2}{3}}}{\sqrt[3]{2x + 5}} \right]$

(iii) $y = \log \left[\frac{1 - \cos\left(\frac{3x}{2}\right)}{1 + \cos\left(\frac{3x}{2}\right)} \right]$ (iv) $y = \log \left[\frac{x + \sqrt{x^2 + a^2}}{\sqrt{x^2 + a^2} - x} \right]$

(v) $y = (4)^{\log_2(\sin x)} + (9)^{\log_3(\cos x)}$ (vi) $y = a^{a^{\log_a(\cot x)}}$

Solution :

(i) $y = \log_3 (\log_5 x)$

$$= \log_3 \left(\frac{\log x}{\log 5} \right) = \log_3 (\log x) - \log_3 (\log 5)$$

$$\therefore y = \frac{\log(\log x)}{\log 3} - \log_3 (\log 5)$$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} \left[\frac{\log(\log x)}{\log 3} - \log_3 (\log 5) \right]$$

$$= \frac{1}{\log 3} \frac{d}{dx} [\log(\log x)] - \frac{d}{dx} [\log_3 (\log 5)]$$

$$= \frac{1}{\log 3} \times \frac{1}{\log x} \frac{d}{dx} (\log x) - 0 \quad [\text{Note that } \log_3 (\log 5) \text{ is constant}]$$

$$= \frac{1}{\log 3} \times \frac{1}{\log x} \times \frac{1}{x}$$

$$\therefore \frac{dy}{dx} = \frac{1}{x \log x \log 3}$$

(ii) $y = \log \left[e^{3x} \cdot \frac{(3x-4)^{\frac{2}{3}}}{\sqrt[3]{2x+5}} \right] = \log \left[\frac{e^{3x} \cdot (3x-4)^{\frac{2}{3}}}{(2x+5)^{\frac{1}{3}}} \right]$

$$= \log \left[e^{3x} \cdot (3x-4)^{\frac{2}{3}} \right] - \log \left[(2x+5)^{\frac{1}{3}} \right]$$

$$= \log e^{3x} + \log(3x-4)^{\frac{2}{3}} - \log(2x+5)^{\frac{1}{3}}$$

$$\therefore y = 3x + \frac{2}{3} \log(3x-4) - \frac{1}{3} \log(2x+5) \quad [\because \log e = 1]$$

Differentiate w. r. t. x


$$\frac{dy}{dx} = \frac{d}{dx} \left[3x + \frac{2}{3} \log(3x-4) - \frac{1}{3} \log(2x+5) \right]$$

$$= 3 \frac{d}{dx} (x) + \frac{2}{3} \cdot \frac{d}{dx} [\log(3x-4)] - \frac{1}{3} \cdot \frac{d}{dx} [\log(2x+5)]$$

$$= 3(1) + \frac{2}{3} \cdot \frac{1}{3x-4} \cdot \frac{d}{dx} (3x-4) - \frac{1}{3} \cdot \frac{1}{2x+5} \cdot \frac{d}{dx} (2x+5)$$

$$= 3(1) + \frac{2}{3} \cdot \frac{1}{3x-4} \cdot (3) - \frac{1}{3} \cdot \frac{1}{2x+5} \cdot (2)$$

$$\therefore \frac{dy}{dx} = 3 + \frac{2}{3x-4} - \frac{2}{3(2x+5)}$$



$$(iii) \quad y = \log \left[\sqrt{\frac{1 - \cos\left(\frac{3x}{2}\right)}{1 + \cos\left(\frac{3x}{2}\right)}} \right] = \log \left[\sqrt{\frac{2 \sin^2\left(\frac{3x}{4}\right)}{2 \cos^2\left(\frac{3x}{4}\right)}} \right]$$

$$\therefore y = \log \left[\tan\left(\frac{3x}{4}\right) \right]$$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} \left\{ \log \left[\tan\left(\frac{3x}{4}\right) \right] \right\}$$

$$= \frac{1}{\tan\left(\frac{3x}{4}\right)} \cdot \frac{d}{dx} \left[\tan\left(\frac{3x}{4}\right) \right]$$

$$= \cot\left(\frac{3x}{4}\right) \cdot \sec^2\left(\frac{3x}{4}\right) \cdot \frac{d}{dx} \left(\frac{3x}{4}\right)$$

$$= \frac{\cos\left(\frac{3x}{4}\right)}{\sin\left(\frac{3x}{4}\right)} \times \frac{1}{\cos^2\left(\frac{3x}{4}\right)} \times \frac{3}{4}$$

$$= \frac{3}{2 \left[2 \sin\left(\frac{3x}{4}\right) \cdot \cos\left(\frac{3x}{4}\right) \right]} = \frac{3}{2 \sin\left(\frac{3x}{2}\right)}$$

$$\therefore \frac{dy}{dx} = \frac{3}{2} \operatorname{cosec}\left(\frac{3x}{2}\right)$$

$$(iv) \quad y = \log \left[\frac{x + \sqrt{x^2 + a^2}}{\sqrt{x^2 + a^2} - x} \right] = \log \left[\frac{\sqrt{x^2 + a^2} + x}{\sqrt{x^2 + a^2} - x} \times \frac{\sqrt{x^2 + a^2} + x}{\sqrt{x^2 + a^2} + x} \right]$$

$$= \log \left[\frac{(\sqrt{x^2 + a^2} + x)^2}{x^2 + a^2 - x^2} \right]$$

$$= \log \left[\frac{(\sqrt{x^2 + a^2} + x)^2}{a^2} \right]$$

$$= \log(\sqrt{x^2 + a^2} + x)^2 - \log(a^2)$$

$$\therefore y = 2 \log(\sqrt{x^2 + a^2} + x) - \log(a^2)$$

Differentiate w. r. t. x



$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} \left[2 \log(\sqrt{x^2 + a^2} + x) - \log(a^2) \right] \\ &= 2 \frac{d}{dx} \left[\log(\sqrt{x^2 + a^2} + x) \right] - \frac{d}{dx} [\log(a^2)] \\ &= 2 \times \frac{1}{\sqrt{x^2 + a^2} + x} \cdot \frac{d}{dx} [\sqrt{x^2 + a^2} + x] - 0 \\ &= \frac{2}{\sqrt{x^2 + a^2} + x} \cdot \left[\frac{1}{2\sqrt{x^2 + a^2}} \cdot \frac{d}{dx} (x^2 + a^2) + 1 \right] \\ &= \frac{2}{\sqrt{x^2 + a^2} + x} \cdot \left[\frac{1}{2\sqrt{x^2 + a^2}} (2x) + 1 \right] \\ &= \frac{2}{\sqrt{x^2 + a^2} + x} \cdot \left[\frac{x + \sqrt{x^2 + a^2}}{\sqrt{x^2 + a^2}} \right] \\ \frac{dy}{dx} &= \frac{2}{\sqrt{x^2 + a^2}} \end{aligned}$$

(v) $y = (4)^{\log_2(\sin x)} + (9)^{\log_3(\cos x)}$
 $= (2^2)^{\log_2(\sin x)} + (3^2)^{\log_3(\cos x)}$
 $= (2)^{2\log_2(\sin x)} + (3)^{2\log_3(\cos x)}$
 $= (2)^{\log_2(\sin^2 x)} + (3)^{\log_3(\cos^2 x)} \quad [\because a^{\log_a f(x)} = f(x)]$
 $= \sin^2 x + \cos^2 x$

$\therefore y = 1$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (1) = 0$$

(vi) $y = a^{a^{\log_a(\cot x)}}$
 $y = a^{\cot x} \quad [\because a^{\log_a f(x)} = f(x)]$

Differentiate w. r. t. x

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} (a^{\cot x}) \\ &= a^{\cot x} \log a \cdot \frac{d}{dx} (\cot x) \\ &= a^{\cot x} \log a (-\operatorname{cosec}^2 x) \end{aligned}$$

$$\frac{dy}{dx} = -\operatorname{cosec}^2 x \cdot a^{\cot x} \log a$$

Ex. 4 : If $f(x) = \sqrt{7g(x) - 3}$, $g(3) = 4$ and $g'(3) = 5$, find $f'(3)$.

Solution : Given that : $f(x) = \sqrt{7g(x) - 3}$

Differentiate w. r. t. x

$$f'(x) = \frac{d}{dx} (\sqrt{7g(x) - 3}) = \frac{1}{2\sqrt{7g(x) - 3}} \frac{d}{dx} [7g(x) - 3]$$

$$\therefore f'(x) = \frac{7g'(x)}{2\sqrt{7g(x) - 3}}$$

For $x = 3$, we get

$$f'(3) = \frac{7g'(3)}{2\sqrt{7g(3) - 3}} = \frac{35}{2(5)} = \frac{7}{2} \quad [\text{Since } g(3) = 4 \text{ and } g'(3) = 5]$$

Ex. 5 : If $F(x) = G \{3G [5G(x)]\}$, $G(0) = 0$ and $G'(0) = 3$, find $F'(0)$.

Solution : Given that : $F(x) = G \{3G [5G(x)]\}$

Differentiate *w. r. t. x*

$$\begin{aligned} F'(x) &= \frac{d}{dx} G \{3G [5G(x)]\} \\ &= G' \{3G [5G(x)]\} \cdot 3 \cdot \frac{d}{dx} [G [5G(x)]] \\ &= G' \{3G [5G(x)]\} \cdot 3 \cdot G' [5G(x)] \cdot 5 \cdot \frac{d}{dx} [G(x)] \end{aligned}$$

$$F'(x) = 15 \cdot G' \{3G [5G(x)]\} G' [5G(x)] G'(x)$$

For $x = 0$, we get

$$\begin{aligned} F'(0) &= 15 \cdot G' \{3G [5G(0)]\} G' [5G(0)] G'(0) \\ &= 15 \cdot G' [3G(0)] G'(0) \cdot (3) \quad [\because G(0) = 0 \text{ and } G'(0) = 3] \\ &= 15 \cdot G'(0)(3)(3) = 15 \cdot (3)(3)(3) = 405 \end{aligned}$$

Ex. 6 : Select the appropriate hint from the hint basket and fill in the blank spaces in the following paragraph. [Activity]

"Let $f(x) = \sin x$ and $g(x) = \log x$ then $f[g(x)] = \dots\dots\dots$ and $g[f(x)] = \dots\dots\dots$. Now $f'(x) = \dots\dots\dots$ and $g'(x) = \dots\dots\dots$.

The derivative of $f[g(x)]$ *w. r. t. x* in terms of f and g is $\dots\dots\dots$.

Therefore $\frac{d}{dx} [f[g(x)]] = \dots\dots\dots$ and $\left[\frac{d}{dx} [f[g(x)]] \right]_{x=1} = \dots\dots\dots$.

The derivative of $g[f(x)]$ *w. r. t. x* in terms of f and g is $\dots\dots\dots$.

Therefore $\frac{d}{dx} [g[f(x)]] = \dots\dots\dots$ and $\left[\frac{d}{dx} [g[f(x)]] \right]_{x=\frac{\pi}{3}} = \dots\dots\dots$ "

Hint basket : $\left\{ f'[g(x)] \cdot g'(x), \frac{\cos(\log x)}{x}, 1, g'[f(x)] \cdot f'(x), \cot x, \sqrt{3}, \sin(\log x), \log(\sin x), \cos x, \frac{1}{x} \right\}$

Solution : $\sin(\log x), \log(\sin x), \cos x, \frac{1}{x}, f'[g(x)] \cdot g'(x), \frac{\cos(\log x)}{x}, 1, g'[f(x)] \cdot f'(x), \cot x, \sqrt{3}$.

EXERCISE 1.1

(1) Differentiate *w. r. t. x*.

- | | | |
|-----------------------------|---|---|
| (i) $(x^3 - 2x - 1)^5$ | (ii) $\left(2x^{\frac{3}{2}} - 3x^{\frac{4}{3}} - 5\right)^{\frac{5}{2}}$ | (v) $\frac{3}{5\sqrt{(2x^2 - 7x - 5)^5}}$ |
| (iii) $\sqrt{x^2 + 4x - 7}$ | (iv) $\sqrt{x^2 + \sqrt{x^2 + 1}}$ | (vi) $\left(\sqrt{3x - 5} - \frac{1}{\sqrt{3x - 5}}\right)^5$ |

(2) Differentiate the following *w.r.t. x*

- (i) $\cos(x^2 + a^2)$ (ii) $\sqrt{e^{(3x+2)} + 5}$
 (iii) $\log\left[\tan\left(\frac{x}{2}\right)\right]$ (iv) $\sqrt{\tan\sqrt{x}}$
 (v) $\cot^3[\log(x^3)]$ (vi) $5^{\sin^3 x + 3}$
 (vii) $\operatorname{cosec}(\sqrt{\cos x})$ (viii) $\log[\cos(x^3 - 5)]$
 (ix) $e^{3 \sin^2 x - 2 \cos^2 x}$ (x) $\cos^2[\log(x^2 + 7)]$
 (xi) $\tan[\cos(\sin x)]$ (xii) $\sec[\tan(x^4 + 4)]$
 (xiii) $e^{\log[(\log x)^2 - \log x^2]}$ (xiv) $\sin\sqrt{\sin\sqrt{x}}$
 (xv) $\log[\sec(e^{x^2})]$ (xvi) $\log_{e^2}(\log x)$
 (xvii) $[\log[\log(\log x)]]^2$
 (xviii) $\sin^2 x^2 - \cos^2 x^2$

(3) Differentiate the following *w.r.t. x*

- (i) $(x^2 + 4x + 1)^3 + (x^3 - 5x - 2)^4$
 (ii) $(1 + 4x)^5 (3 + x - x^2)^8$
 (iii) $\frac{x}{\sqrt{7-3x}}$ (iv) $\frac{(x^3 - 5)^5}{(x^3 + 3)^3}$
 (v) $(1 + \sin^2 x)^2 (1 + \cos^2 x)^3$
 (vi) $\sqrt{\cos x} + \sqrt{\cos\sqrt{x}}$
 (vii) $\log(\sec 3x + \tan 3x)$ (viii) $\frac{1 + \sin x^\circ}{1 - \sin x^\circ}$
 (ix) $\cot\left(\frac{\log x}{2}\right) - \log\left(\frac{\cot x}{2}\right)$
 (x) $\frac{e^{2x} - e^{-2x}}{e^{2x} + e^{-2x}}$ (xi) $\frac{e^{\sqrt{x}} + 1}{e^{\sqrt{x}} - 1}$
 (xii) $\log[\tan^3 x \cdot \sin^4 x \cdot (x^2 + 7)^7]$
 (xiii) $\log\left(\sqrt{\frac{1 - \cos 3x}{1 + \cos 3x}}\right)$
 (xiv) $\log\left(\sqrt{\frac{1 + \cos\left(\frac{5x}{2}\right)}{1 - \cos\left(\frac{5x}{2}\right)}}\right)$

- (xv) $\log\left(\sqrt{\frac{1 - \sin x}{1 + \sin x}}\right)$
 (xvi) $\log\left[4^{2x} \left(\frac{x^2 + 5}{\sqrt{2x^3 - 4}}\right)^{\frac{3}{2}}\right]$
 (xvii) $\log\left[\frac{e^{x^2} (5 - 4x)^{\frac{3}{2}}}{\sqrt[3]{7 - 6x}}\right]$
 (xviii) $\log\left(\frac{a^{\cos x}}{(x^2 - 3)^3 \log x}\right)$
 (xix) $y = (25)^{\log_5(\sec x)} - (16)^{\log_4(\tan x)}$
 (xx) $\frac{(x^2 + 2)^4}{\sqrt{x^2 - 5}}$

(4) A table of values of f, g, f' and g' is given

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
2	1	6	-3	4
4	3	4	5	-6
6	5	2	-4	7

- (i) If $r(x) = f[g(x)]$ find $r'(2)$.
 (ii) If $R(x) = g[3 + f(x)]$ find $R'(4)$.
 (iii) If $s(x) = f[9 - f(x)]$ find $s'(4)$.
 (iv) If $S(x) = g[g(x)]$ find $S'(6)$.
 (5) Assume that $f'(3) = -1, g'(2) = 5, g(2) = 3$ and $y = f[g(x)]$ then $\left[\frac{dy}{dx}\right]_{x=2} = ?$
 (6) If $h(x) = \sqrt{4f(x) + 3g(x)}, f(1) = 4, g(1) = 3, f'(1) = 3, g'(1) = 4$ find $h'(1)$.
 (7) Find the x co-ordinates of all the points on the curve $y = \sin 2x - 2 \sin x, 0 \leq x < 2\pi$ where $\frac{dy}{dx} = 0$.

(8) Select the appropriate hint from the hint basket and fill up the blank spaces in the following paragraph. [Activity]

"Let $f(x) = x^2 + 5$ and $g(x) = e^x + 3$ then

$f[g(x)] = \text{-----}$ and

$g[f(x)] = \text{-----}$.

Now $f'(x) = \text{-----}$ and

$g'(x) = \text{-----}$.

The derivative of $f[g(x)]$ w. r. t. x in terms

of f and g is ----- .

Therefore $\frac{d}{dx}[f[g(x)]] = \text{-----}$ and

$\left[\frac{d}{dx}[f[g(x)]]\right]_{x=0} = \text{-----}$.

The derivative of $g[f(x)]$ w. r. t. x in terms of f and g is ----- .

Therefore $\frac{d}{dx}[g[f(x)]] = \text{-----}$

and $\left[\frac{d}{dx}[g[f(x)]]\right]_{x=1} = \text{-----}$."

Hint basket : $\{f'[g(x)] \cdot g'(x), 2e^{2x} + 6e^x, 8, g'[f(x)] \cdot f'(x), 2xe^{x^2+5}, -2e^6, e^{2x} + 6e^x + 14, e^{x^2+5} + 3, 2x, e^x\}$

1.2.1 Geometrical meaning of Derivative :

Consider a point P on the curve $f(x)$. At $x = a$, the coordinates of P are $(a, f(a))$. Let Q be another point on the curve, a little to the right of P i.e. to the right of $x = a$, with a value increased by a small real number h . Therefore the coordinates of Q are $((a + h), f(a + h))$. Now we can calculate the slope of the secant line PQ i.e. slope of the secant line connecting the points P $(a, f(a))$ and Q $((a + h), f(a + h))$, by using formula for slope.

$$\begin{aligned} \text{Slope of secant PQ} &= \frac{f(a+h) - f(a)}{a+h-a} \\ &= \frac{f(a+h) - f(a)}{h} \end{aligned}$$




Fig. 1.2.1

Suppose we make h smaller and smaller then $a + h$ will approach a as h gets closer to zero, Q will approach P, that is as $h \rightarrow 0$, the secant covers to the tangent at P.

$$\therefore \lim_{Q \rightarrow P} (\text{Slope of secant PQ}) = \lim_{h \rightarrow 0} \left[\frac{f(a+h) - f(a)}{h} \right] = f'(a)$$

So we get, Slope of tangent at P = $f'(a)$ [If limit exists]

Thus the derivative of a function $y = f(x)$ at any point P (x_1, y_1) is the slope of the tangent at that point on the curve. If we consider the point $a - h$ to the left of a , $h > 0$, then with R = $((a - h), f(a - h))$ we will find the slope of PR which will also converge to the slope of tangent at P.

For Example : If $y = x^2 + 3x + 2$ then slope of the tangent at (2,3) is given by

$$\text{Slope } m = \left[\frac{dy}{dx} \right]_{(2,3)} = \left[\frac{d}{dx}(x^2 + 3x + 2) \right]_{(2,3)} = (2x + 3)_{(2,3)} = 2(2) + 3 \quad \therefore m = 7$$

1.2.2 Derivatives of Inverse Functions :

We know that if $y = f(x)$ is a one-one and onto function then $x = f^{-1}(y)$ exists. If $f^{-1}(y)$ is differentiable then we can find its derivative. In this section let us discuss the derivatives of some inverse functions and the derivatives of inverse trigonometric functions.

Example 1 : Consider $f(x) = 2x - 2$ then its inverse is $f^{-1}(x) = \frac{x+2}{2}$. Let $g(x) = f^{-1}(x)$.

If we find the derivatives of these functions we see that $\frac{d}{dx} [f(x)] = 2$ and $\frac{d}{dx} [g(x)] = \frac{1}{2}$.

These derivatives are reciprocals of one another.

Example 2 : Consider $y = f(x) = x^2$. Let $g = f^{-1}$.

$$\therefore g(y) = x = \sqrt{y}$$

$$\therefore g'(y) = \frac{1}{2\sqrt{y}} \text{ also } f'(x) = 2x$$

$$\text{Now } \frac{d}{dx} [g(f(x))] = \frac{f'(x)}{2\sqrt{f(x)}} = \frac{2x}{2\sqrt{x^2}} = 1 \text{ and } g[f(x)] = x \therefore \frac{d}{dx} [g(f(x))] = \frac{d}{dx} (x) = 1$$

$$\text{At a point } (x, x^2) \text{ on the curve, } f'(x) = 2x \text{ and } g'(y) = \frac{1}{2\sqrt{y}} = \frac{1}{2x} = \frac{1}{f'(x)}.$$

1.2.3 Theorem : Suppose $y = f(x)$ is a differentiable function of x on an interval I and y is One-one, onto and

$\frac{dy}{dx} \neq 0$ on I . Also if $f^{-1}(y)$ is differentiable on $f(I)$ then $\frac{d}{dy} [f^{-1}(y)] = \frac{1}{f'(x)}$ or $\frac{dx}{dy} = \frac{1}{\frac{dy}{dx}}$ where $\frac{dy}{dx} \neq 0$.

Proof : Given that $y = f(x)$ and $x = f^{-1}(y)$ are differentiable functions.

Let there be a small increment in the value of x say δx then correspondingly there will be an increment in the value of y say δy . As δx and δy are increments, $\delta x \neq 0$ and $\delta y \neq 0$.

$$\text{We have, } \frac{\delta x}{\delta y} \times \frac{\delta y}{\delta x} = 1$$

$$\therefore \frac{\delta x}{\delta y} = \frac{1}{\frac{\delta y}{\delta x}}, \text{ where } \frac{\delta y}{\delta x} \neq 0$$

Taking the limit as $\delta x \rightarrow 0$, we get,

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta x}{\delta y} \right) = \lim_{\delta x \rightarrow 0} \left(\frac{1}{\frac{\delta y}{\delta x}} \right)$$

as $\delta x \rightarrow 0$, $\delta y \rightarrow 0$,

$$\lim_{\delta y \rightarrow 0} \left(\frac{\delta x}{\delta y} \right) = \frac{1}{\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right)} \dots \dots (I)$$

Since $y = f(x)$ is a differentiable function of x .



we have, $\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \frac{dy}{dx}$ and $\frac{dy}{dx} \neq 0$ (II)

From (I) and (II), we get

$\lim_{\delta y \rightarrow 0} \left(\frac{\delta x}{\delta y} \right) = \frac{1}{\frac{dy}{dx}}$ (III)

As $\frac{dy}{dx} \neq 0$, $\frac{1}{\frac{dy}{dx}}$ exists and is finite. $\therefore \lim_{\delta y \rightarrow 0} \left(\frac{\delta x}{\delta y} \right) = \frac{dx}{dy}$ exists and is finite.

Hence, from (III) $\frac{dx}{dy} = \frac{1}{\frac{dy}{dx}}$ where $\frac{dy}{dx} \neq 0$

An alternative proof using derivatives of composite functions rule.

We know that $f^{-1} [f(x)] = x$ [Identity function]

Taking derivative on both sides we get,

$\frac{d}{dx} [f^{-1} [f(x)]] = \frac{d}{dx} (x)$

i.e. $(f^{-1})' [f(x)] \frac{d}{dx} [f(x)] = 1$

i.e. $(f^{-1})' [f(x)] f'(x) = 1$

$\therefore (f^{-1})' [f(x)] = \frac{1}{f'(x)}$ (I)

So, if $y = f(x)$ is a differentiable function of x and $x = f^{-1} (y)$ exists and is differentiable then

$(f^{-1})' [f(x)] = (f^{-1})' (y) = \frac{dx}{dy}$ and $f'(x) = \frac{dy}{dx}$

\therefore (I) becomes

$\frac{dx}{dy} = \frac{1}{\frac{dy}{dx}}$ where $\frac{dy}{dx} \neq 0$

 **SOLVED EXAMPLES**

Ex. 1 : Find the derivative of the function $y = f(x)$ using the derivative of the inverse function $x = f^{-1}(y)$ in the following

(i) $y = \sqrt[3]{x+4}$

(ii) $y = \sqrt{1+\sqrt{x}}$

(iii) $y = \ln x$

Solution :

(i) $y = \sqrt[3]{x+4}$

We first find the inverse of the function $y = f(x)$, i.e. x in term of y .

$y^3 = x + 4 \therefore x = y^3 - 4 \therefore x = f^{-1}(y) = y^3 - 4$

$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}} = \frac{1}{\frac{d}{dy}(y^3 - 4)} = \frac{1}{3y^2}$

$= \frac{1}{3(\sqrt[3]{x+4})^2} = \frac{1}{3\sqrt[3]{(x+4)^2}}$ for $x \neq -4$

(ii) $y = \sqrt{1 + \sqrt{x}}$

We first find the inverse of the function $y = f(x)$, i.e. x in term of y .

$$y^2 = 1 + \sqrt{x} \text{ i.e. } \sqrt{x} = y^2 - 1, \therefore x = f^{-1}(y) = (y^2 - 1)^2$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{1}{\frac{dx}{dy}} = \frac{1}{\frac{d}{dy}[(y^2 - 1)^2]} = \frac{1}{2(y^2 - 1)\frac{d}{dy}(y^2 - 1)} \\ &= \frac{1}{2(y^2 - 1)(2y)} = \frac{1}{4\sqrt{1 + \sqrt{x}}\left[\left(\sqrt{1 + \sqrt{x}}\right)^2 - 1\right]} \\ &= \frac{1}{4\sqrt{1 + \sqrt{x}}(1 + \sqrt{x} - 1)} = \frac{1}{4\sqrt{x}\sqrt{1 + \sqrt{x}}} \end{aligned}$$

(iii) $y = \log x$

We first find the inverse of the function $y = f(x)$, i.e. x in term of y .

$$y = \log x \therefore x = f^{-1}(y) = e^y$$

$$\frac{dy}{dx} = \frac{1}{\frac{dx}{dy}} = \frac{1}{\frac{d}{dy}(e^y)} = \frac{1}{e^y} = \frac{1}{e^{\ln x}} = \frac{1}{x}$$

Ex. 2 : Find the derivative of the inverse of function $y = 2x^3 - 6x$ and calculate its value at $x = -2$.

Solution : Given : $y = 2x^3 - 6x$

Diff. w. r. t. x we get,

$$\frac{dy}{dx} = 6x^2 - 6 = 6(x^2 - 1)$$

we have, $\frac{dx}{dy} = \frac{1}{\frac{dy}{dx}}$

$$\therefore \frac{dx}{dy} = \frac{1}{6(x^2 - 1)}$$

at $x = -2$,

$$\begin{aligned} \text{we get, } y &= 2(-2)^3 - 6(-2) \\ &= -16 + 12 = -4 \end{aligned}$$

$$\begin{aligned} \left[\frac{dx}{dy}\right]_{y=-4} &= \frac{1}{\left[\frac{dy}{dx}\right]_{x=-2}} \\ &= \frac{1}{6((-2)^2 - 1)} \\ &= \frac{1}{18} \end{aligned}$$

Ex. 3 : Let f and g be the inverse functions of each other. The following table lists a few values of f, g and f'

x	$f(x)$	$g(x)$	$f'(x)$
-4	2	1	$\frac{1}{3}$
1	-4	-2	4

find $g'(-4)$.

Solution : In order to find $g'(-4)$, we should first find an expression for $g'(x)$ for any input x . Since f and g are inverses we can use the following identify which holds for any two differentiable inverse functions.

$$\begin{aligned} g'(x) &= \frac{1}{f'[g(x)]} \quad \dots \text{ [check, how?]} \\ &\quad \dots \text{ [Hint : } f[g(x)] = x \text{]} \end{aligned}$$

$$\begin{aligned} \therefore g'(-4) &= \frac{1}{f'[g(-4)]} \\ &= \frac{1}{f'(1)} = \frac{1}{4} \end{aligned}$$

Ex. 4 : Let $f(x) = x^5 + 2x - 3$. Find $(f^{-1})'(-3)$.

Solution : Given : $f(x) = x^5 + 2x - 3$

Diff. w. r. t. x we get,

$$f'(x) = 5x^4 + 2$$

Note that $y = -3$ corresponds to $x = 0$.

$$\begin{aligned} \therefore (f^{-1})'(-3) &= \frac{1}{f'(0)} \\ &= \frac{1}{5(0) + 2} = \frac{1}{2} \end{aligned}$$

1.2.4 Derivatives of Standard Inverse trigonometric Functions :

We observe that inverse trigonometric functions are multi-valued functions and because of this, their derivatives depend on which branch of the function we are dealing with. We are not restricted to use these branches all the time. While solving the problems it is customary to select the branch of the inverse trigonometric function which is applicable to the kind of problem we are solving. We have to pay more attention towards the domain and range.

1. If $y = \sin^{-1} x$, $-1 \leq x \leq 1$, $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ then prove that $\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$, $|x| < 1$.

Proof : Given that $y = \sin^{-1} x$, $-1 \leq x \leq 1$, $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

$$\therefore x = \sin y \quad \dots \text{ (I)}$$

Differentiate w. r. t. y

$$\frac{dx}{dy} = \frac{d}{dy}(\sin y)$$

$$\frac{dx}{dy} = \cos y = \pm \sqrt{\cos^2 y} = \pm \sqrt{1 - \sin^2 y}$$

$$\therefore \frac{dx}{dy} = \pm \sqrt{1 - x^2} \quad \dots [\because \sin y = x]$$

But $\cos y$ is positive since y lies in 1st or 4th quadrant as $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

$$\therefore \frac{dx}{dy} = \sqrt{1 - x^2}$$

$$\text{We have } \frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}}, |x| < 1$$

2. If $y = \cos^{-1} x$, $-1 \leq x \leq 1$, $0 \leq y \leq \pi$ then prove that $\frac{dy}{dx} = -\frac{1}{\sqrt{1-x^2}}$.

[As home work for students to prove.]



3. If $y = \cot^{-1} x, x \in R, 0 < y < \pi$ then $\frac{dy}{dx} = -\frac{1}{1+x^2}$.

Proof: Given that $y = \cot^{-1} x, x \in R, 0 < y < \pi$

$$\therefore x = \cot y \quad \dots (I)$$

Differentiate w. r. t. y

$$\frac{dx}{dy} = \frac{d}{dy}(\cot y)$$

$$\frac{dx}{dy} = -\operatorname{cosec}^2 y = -(1 + \cot^2 y)$$

$$\therefore \frac{dx}{dy} = -(1 + x^2) \quad \dots [\because \cot y = x]$$

$$\text{We have } \frac{dy}{dx} = \frac{1}{\frac{dx}{dy}}$$

$$\therefore \frac{dy}{dx} = \frac{1}{-(1+x^2)} \quad \therefore \frac{dy}{dx} = -\frac{1}{1+x^2}$$

4. If $y = \tan^{-1} x, x \in R, -\frac{\pi}{2} < y < \frac{\pi}{2}$ then $\frac{dy}{dx} = \frac{1}{1+x^2}$. [left as home work for students to prove.]

5. If $y = \sec^{-1} x$, such that $|x| \geq 1$ and $0 \leq y \leq \pi, y \neq \frac{\pi}{2}$ then $\frac{dy}{dx} = \frac{1}{x\sqrt{x^2-1}}$ if $x > 1$
 $\frac{dy}{dx} = -\frac{1}{x\sqrt{x^2-1}}$ if $x < -1$

Proof: Given that $y = \sec^{-1} x, |x| \geq 1$ and $0 \leq y \leq \pi, y \neq \frac{\pi}{2}$

$$\therefore x = \sec y \quad \dots (I)$$

Differentiate w. r. t. y

$$\frac{dx}{dy} = \frac{d}{dy}(\sec y)$$

$$\frac{dx}{dy} = \sec y \cdot \tan y$$

$$\therefore \frac{dx}{dy} = \pm \sec y \cdot \sqrt{\tan^2 y} = \pm \sec y \cdot \sqrt{\sec^2 y - 1}$$

$$\therefore \frac{dx}{dy} = \pm x \sqrt{x^2 - 1} \quad \dots [\because \sec y = x]$$



We use the sign \pm because for y in 1st and 2nd quadrant. $\sec y \cdot \tan y > 0$.

Hence we choose $x \sqrt{x^2 - 1}$ if $x > 1$ and $-x \sqrt{x^2 - 1}$ if $x < -1$

In 1st quadrant both $\sec y$ and $\tan y$ are positive.

In 2nd quadrant both $\sec y$ and $\tan y$ are negative.

$\therefore \sec y \cdot \tan y$ is positive in both first and second quadrant.



Also, for $x > 0$, $x\sqrt{x^2-1} > 0$

and for $x < 0$, $-x\sqrt{x^2-1} > 0$

$$\frac{dy}{dx} = x\sqrt{x^2-1}, \quad \text{when } x > 0, |x| > 1 \quad \text{i.e. } x > 1$$

$$= -x\sqrt{x^2-1}, \quad \text{when } x < 0, |x| > 1 \quad \text{i.e. } x < -1$$

$$\frac{dy}{dx} = \frac{1}{x\sqrt{x^2-1}} \quad \text{if } x > 1$$

$$\frac{dy}{dx} = -\frac{1}{x\sqrt{x^2-1}} \quad \text{if } x < -1$$

Note 1 : A function is increasing if its derivative is positive and is decreasing if its derivative is negative.

Note 2 : The derivative of $\sec^{-1} x$ is always positive because the graph of $\sec^{-1} x$ is always increasing.

6. If $y = -\operatorname{cosec} x$, such that $|x| \geq 1$ and $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}, y \neq 0$ then

$$\frac{dy}{dx} = -\frac{1}{x\sqrt{x^2-1}} \quad \text{if } x > 1$$

$$\frac{dy}{dx} = \frac{1}{x\sqrt{x^2-1}} \quad \text{if } x < -1$$

[Left as home work for students to prove]

Note 3 : The derivative of $\operatorname{cosec}^{-1} x$ is always negative because the graph of $\operatorname{cosec}^{-1} x$ is always decreasing.

1.2.5 Derivatives of Standard Inverse trigonometric Functions :

y	$\frac{dy}{dx}$	Conditions	y	$\frac{dy}{dx}$	Conditions
$\sin^{-1} x$	$\frac{1}{\sqrt{1-x^2}}, x < 1$	$-1 \leq x \leq 1$ $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$	$\sec^{-1} x$	$\frac{1}{x\sqrt{x^2-1}}$ for $x > 1$ $-\frac{1}{x\sqrt{x^2-1}}$ for $x < -1$	$ x \geq 1$ $0 \leq y \leq \pi$ $y \neq \frac{\pi}{2}$
$\cos^{-1} x$	$-\frac{1}{\sqrt{1-x^2}}, x < 1$	$-1 \leq x \leq 1$ $0 \leq y \leq \pi$	$\operatorname{cosec}^{-1} x$	$-\frac{1}{x\sqrt{x^2-1}}$ for $x > 1$ $\frac{1}{x\sqrt{x^2-1}}$ for $x < -1$	$ x \geq 1$ $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$ $y \neq 0$
$\tan^{-1} x$	$\frac{1}{1+x^2}$	$x \in R$ $-\frac{\pi}{2} < y < \frac{\pi}{2}$			
$\cot^{-1} x$	$-\frac{1}{1+x^2}$	$x \in R$ $0 < y < \pi$			

Table 1.2.1

1.2.6 Derivatives of Standard Inverse trigonometric Composite Functions :

y	$\frac{dy}{dx}$	y	$\frac{dy}{dx}$
$\sin^{-1} [f(x)]$	$\frac{f'(x)}{\sqrt{1 - [f(x)]^2}}, f(x) < 1$	$\cot^{-1} [f(x)]$	$-\frac{f'(x)}{1 + [f(x)]^2}$
$\cos^{-1} [f(x)]$	$-\frac{f'(x)}{\sqrt{1 - [f(x)]^2}}, f(x) < 1$	$\sec^{-1} [f(x)]$	$\frac{f'(x)}{f(x) \sqrt{[f(x)]^2 - 1}}, \text{ for } f(x) > 1$
$\tan^{-1} [f(x)]$	$\frac{f'(x)}{1 + [f(x)]^2}$	$\operatorname{cosec}^{-1} [f(x)]$	$-\frac{f'(x)}{f(x) \sqrt{[f(x)]^2 - 1}}, \text{ for } f(x) > 1$

Table 1.2.2

Some Important Formulae for Inverse Trigonometric Functions :

(1) $\sin^{-1} (\sin \theta) = \theta, \sin(\sin^{-1} x) = x$	(2) $\cos^{-1} (\cos \theta) = \theta, \cos(\cos^{-1} x) = x$	
(3) $\tan^{-1} (\tan \theta) = \theta, \tan(\tan^{-1} x) = x$	(4) $\cot^{-1} (\cot \theta) = \theta, \cot(\cot^{-1} x) = x$	
(5) $\sec^{-1} (\sec \theta) = \theta, \sec(\sec^{-1} x) = x$	(6) $\operatorname{cosec}^{-1} (\operatorname{cosec} \theta) = \theta, \operatorname{cosec}(\operatorname{cosec}^{-1} x) = x$	
(7) $\sin^{-1} (\cos \theta) = \sin^{-1} \left[\sin \left(\frac{\pi}{2} - \theta \right) \right] = \frac{\pi}{2} - \theta$	(8) $\cos^{-1} (\sin \theta) = \cos^{-1} \left[\cos \left(\frac{\pi}{2} - \theta \right) \right] = \frac{\pi}{2} - \theta$	
(9) $\tan^{-1} (\cot \theta) = \tan^{-1} \left[\tan \left(\frac{\pi}{2} - \theta \right) \right] = \frac{\pi}{2} - \theta$	(10) $\cot^{-1} (\tan \theta) = \cot^{-1} \left[\cot \left(\frac{\pi}{2} - \theta \right) \right] = \frac{\pi}{2} - \theta$	
(11) $\sec^{-1} (\operatorname{cosec} \theta) = \sec^{-1} \left[\sec \left(\frac{\pi}{2} - \theta \right) \right] = \frac{\pi}{2} - \theta$		
(12) $\operatorname{cosec}^{-1} (\sec \theta) = \operatorname{cosec}^{-1} \left[\operatorname{cosec} \left(\frac{\pi}{2} - \theta \right) \right] = \frac{\pi}{2} - \theta$		
(13) $\sin^{-1} (x) = \operatorname{cosec}^{-1} \left(\frac{1}{x} \right)$	(14) $\operatorname{cosec}^{-1} (x) = \sin^{-1} \left(\frac{1}{x} \right)$	
(15) $\cos^{-1} (x) = \sec^{-1} \left(\frac{1}{x} \right)$	(16) $\sec^{-1} (x) = \cos^{-1} \left(\frac{1}{x} \right)$	
(17) $\tan^{-1} (x) = \cot^{-1} \left(\frac{1}{x} \right)$	(18) $\cot^{-1} (x) = \tan^{-1} \left(\frac{1}{x} \right)$	
(19) $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$	(20) $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$	(21) $\sec^{-1} x + \operatorname{cosec}^{-1} x = \frac{\pi}{2}$
(22) $\tan^{-1} x + \tan^{-1} y = \tan^{-1} \left(\frac{x + y}{1 - xy} \right)$		(23) $\tan^{-1} x - \tan^{-1} y = \tan^{-1} \left(\frac{x - y}{1 + xy} \right)$

In above tables, x is a real variable with restrictions. **Table 1.2.3**

Some Important Substitutions :

Expression	Substitutions	Expression	Substitutions
$\sqrt{1-x^2}$	$x = \sin \theta$ or $x = \cos \theta$	$\frac{2x}{1+x^2}$	$x = \tan \theta$
$\sqrt{1-x^2}$	$x = \tan \theta$ or $x = \cot \theta$	$\frac{1-x^2}{1+x^2}$	$x = \tan \theta$
$\sqrt{x^2+1}$	$x = \sec \theta$ or $x = \operatorname{cosec} \theta$	$3x-4x^3$ or $1-2x^2$	$x = \sin \theta$
$\sqrt{\frac{a+x}{a-x}}$ or $\sqrt{\frac{a-x}{a+x}}$	$x = a \cos 2\theta$ or $x = a \cos \theta$	$4x^3-3x$ or $2x^2-1$	$x = \cos \theta$
$\sqrt{\frac{1+x}{1-x}}$ or $\sqrt{\frac{1-x}{1+x}}$	$x = \cos 2\theta$ or $x = \cos \theta$	$\frac{3x-x^3}{1-3x^2}$	$x = \tan \theta$
$\sqrt{\frac{a+x^2}{a-x^2}}$ or $\sqrt{\frac{a-x^2}{a+x^2}}$	$x^2 = a \cos 2\theta$ or $x^2 = a \cos \theta$	$\frac{2f(x)}{1+[f(x)]^2}$ or $\frac{1-[f(x)]^2}{1+[f(x)]^2}$	$f(x) = \tan \theta$

Table 1.2.4**SOLVED EXAMPLES**

Ex. 1 : Using derivative prove that $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$.

Solution : Let $f(x) = \sin^{-1} x + \cos^{-1} x$ (I)

We have to prove that $f(x) = \frac{\pi}{2}$

Differentiate (I) w. r. t. x

$$\frac{d}{dx}[f(x)] = \frac{d}{dx}[\sin^{-1} x + \cos^{-1} x]$$

$$f'(x) = \frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-x^2}} = 0$$

$f'(x) = 0 \Rightarrow f(x)$ is a constant function.

Let $f(x) = c$. For any value of x , $f(x)$ must be c only. So conveniently we can choose $x = 0$,

\therefore from (I) we get,

$$f(0) = \sin^{-1}(0) + \cos^{-1}(0) = 0 + \frac{\pi}{2} = \frac{\pi}{2} \Rightarrow c = \frac{\pi}{2} \therefore f(x) = \frac{\pi}{2}$$

Hence, $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$.

Ex. 2 : Differentiate the following w. r. t. x .

(i) $\sin^{-1}(x^3)$

(ii) $\cos^{-1}(2x^2-x)$

(iii) $\sin^{-1}(2^x)$

(iv) $\cot^{-1}\left(\frac{1}{x^2}\right)$

(v) $\cos^{-1}\left(\sqrt{\frac{1+x}{2}}\right)$

(vi) $\sin^2(\sin^{-1}(x^2))$

Solution :

(i) Let $y = \sin^{-1}(x^3)$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\sin^{-1}(x^3)) \\ &= \frac{1}{\sqrt{1-(x^3)^2}} \cdot \frac{d}{dx}(x^3) \\ &= \frac{1}{\sqrt{1-x^6}} (3x^2) \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{3x^2}{\sqrt{1-x^6}}$$

(iii) Let $y = \sin^{-1}(2^x)$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\sin^{-1}(2^x)) \\ &= \frac{1}{\sqrt{1-(2^x)^2}} \cdot \frac{d}{dx}(2^x) \\ &= \frac{1}{\sqrt{1-2^{2x}}} (2^x \log 2) \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{2^x \log 2}{\sqrt{1-4^x}}$$

(iv) Let $y = \cot^{-1}\left(\frac{1}{x^2}\right) = \tan^{-1}(x^2)$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\tan^{-1}(x^2)) \\ &= \frac{1}{1+(x^2)^2} \cdot \frac{d}{dx}(x^2) \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{2x}{1+x^4}$$

(vi) Let $y = \sin^2(\sin^{-1}(x^2))$

$$= [\sin(\sin^{-1}(x^2))]^2 = (x^2)^2$$

$$\therefore y = x^4$$

Differentiate w. r. t. x.

$$\frac{dy}{dx} = \frac{d}{dx}(x^4) \qquad \therefore \frac{dy}{dx} = 4x^3$$

(ii) Let $y = \cos^{-1}(2x^2 - x)$

$$\text{Hence } \cos y = 2x^2 - x \qquad \dots \text{ (I)}$$

Differentiate w. r. t. x.

$$-\sin y \cdot \frac{dy}{dx} = 4x - 1$$

$$\frac{dy}{dx} = \frac{1-4x}{\sin y} = \frac{1-4x}{\sqrt{1-\cos^2 y}}$$

$$\therefore \frac{dy}{dx} = \frac{1-4x}{\sqrt{1-x^2(2x-1)^2}} \qquad \dots \text{ from (I)}$$

Alternate Method :

If $y = \cos^{-1}(2x^2 - x)$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx}(\cos^{-1}(2x^2 - x)) \\ &= \frac{-1}{\sqrt{1-(2x^2-x)^2}} \cdot \frac{d}{dx}(2x^2 - x) \\ &= \frac{-1}{\sqrt{1-x^2(2x-1)^2}} \cdot (4x - 1) \end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{1-4x}{\sqrt{1-x^2(2x-1)^2}}$$

(v) Let $y = \cos^{-1}\left(\sqrt{\frac{1+x}{2}}\right)$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} \left[\cos^{-1} \left(\sqrt{\frac{1+x}{2}} \right) \right] \\ &= - \frac{1}{\sqrt{1-\left(\sqrt{\frac{1+x}{2}}\right)^2}} \cdot \frac{d}{dx} \left(\sqrt{\frac{1+x}{2}} \right) \\ &= - \frac{1}{\sqrt{1-\frac{1+x}{2}}} \times \frac{1}{2\sqrt{\frac{1+x}{2}}} \times \frac{d}{dx} \left(\frac{1+x}{2} \right) \\ &= - \frac{\sqrt{2}}{\sqrt{1-x}} \times \frac{1}{\sqrt{2}\sqrt{1+x}} \times \frac{1}{2} \end{aligned}$$

$$\therefore \frac{dy}{dx} = - \frac{1}{2\sqrt{1-x^2}}$$

Ex. 3 : Differentiate the following w. r. t. x.

(i) $\cos^{-1}(4\cos^3 x - 3\cos x)$ (ii) $\cos^{-1}[\sin(4^x)]$ (iii) $\sin^{-1}\left(\sqrt{\frac{1-\cos x}{2}}\right)$

(iv) $\tan^{-1}\left(\frac{1-\cos 3x}{\sin 3x}\right)$ (v) $\cot^{-1}\left(\frac{\cos x}{1+\sin x}\right)$

Solution :

(i) Let $y = \cos^{-1}(4\cos^3 x - 3\cos x)$
 $= \cos^{-1}(\cos 3x)$
 $\therefore y = 3x$

Differentiate w. r. t. x.

$$\frac{dy}{dx} = \frac{d}{dx}(3x)$$

$\therefore \frac{dy}{dx} = 3$

(iii) Let $y = \sin^{-1}\left(\sqrt{\frac{1-\cos x}{2}}\right)$
 $= \sin^{-1}\left(\sqrt{\frac{2\sin^2\left(\frac{x}{2}\right)}{2}}\right)$

$$= \sin^{-1}\left[\sin\left(\frac{x}{2}\right)\right]$$

$\therefore y = \frac{x}{2}$

Differentiate w. r. t. x.

$$\frac{dy}{dx} = \frac{d}{dx}\left(\frac{x}{2}\right) \quad \therefore \frac{dy}{dx} = \frac{1}{2}$$

(v) Let $y = \cot^{-1}\left(\frac{\cos x}{1+\sin x}\right) = \tan^{-1}\left(\frac{1+\sin x}{\cos x}\right)$

$$= \tan^{-1}\left(\frac{[\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)]^2}{\cos^2\left(\frac{x}{2}\right) - \sin^2\left(\frac{x}{2}\right)}\right) = \tan^{-1}\left(\frac{[\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)]^2}{[\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)][\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)]}\right)$$

$$= \tan^{-1}\left(\frac{\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)}\right) = \tan^{-1}\left(\frac{1 + \tan\left(\frac{x}{2}\right)}{1 - \tan\left(\frac{x}{2}\right)}\right) \dots \text{Divide Numerator \& Denominator by } \cos\left(\frac{x}{2}\right)$$

$$= \tan^{-1}\left[\tan\left(\frac{\pi}{4} + \frac{x}{2}\right)\right] \quad \therefore y = \frac{\pi}{4} + \frac{x}{2}$$

Differentiate w. r. t. x.

$$\frac{dy}{dx} = \frac{d}{dx}\left(\frac{\pi}{4} + \frac{x}{2}\right) = 0 + \frac{1}{2} \quad \therefore \frac{dy}{dx} = \frac{1}{2}$$

(ii) Let $y = \cos^{-1}[\sin(4^x)]$

$$= \cos^{-1}\left[\cos\left(\frac{\pi}{2} - 4^x\right)\right]$$

$\therefore y = \frac{\pi}{2} - 4^x$

Differentiate w. r. t. x.

$$\frac{dy}{dx} = \frac{d}{dx}\left(\frac{\pi}{2} - 4^x\right) = 0 - 4^x \log 4$$

$\therefore \frac{dy}{dx} = -4^x \log 4$

(iv) Let $y = \tan^{-1}\left(\frac{1-\cos 3x}{\sin 3x}\right)$
 $= \tan^{-1}\left(\frac{2\sin^2\left(\frac{3x}{2}\right)}{2\sin\left(\frac{3x}{2}\right)\cos\left(\frac{3x}{2}\right)}\right)$

$$= \tan^{-1}\left[\tan\left(\frac{3x}{2}\right)\right]$$

$\therefore y = \frac{3x}{2}$

Differentiate w. r. t. x.

$$\frac{dy}{dx} = \frac{d}{dx}\left(\frac{3x}{2}\right) \quad \therefore \frac{dy}{dx} = \frac{3}{2}$$

Ex. 4 : Differentiate the following w. r. t. x .

$$(i) \sin^{-1}\left(\frac{2 \cos x + 3 \sin x}{\sqrt{13}}\right) \quad (ii) \cos^{-1}\left(\frac{3 \sin x^2 + 4 \cos x^2}{5}\right) \quad (iii) \sin^{-1}\left(\frac{a \cos x - b \sin x}{\sqrt{a^2 + b^2}}\right)$$

Solution :

$$(i) \text{ Let } y = \sin^{-1}\left(\frac{2 \cos x + 3 \sin x}{\sqrt{13}}\right)$$

$$= \sin^{-1}\left(\frac{2}{\sqrt{13}} \cos x + \frac{3}{\sqrt{13}} \sin x\right)$$

$$\text{Put } \frac{2}{\sqrt{13}} = \sin \alpha, \frac{3}{\sqrt{13}} = \cos \alpha$$

$$\text{Also, } \sin^2 \alpha + \cos^2 \alpha = \frac{4}{13} + \frac{9}{13} = 1$$

$$\text{And } \tan \alpha = \frac{2}{3} \therefore \alpha = \tan^{-1}\left(\frac{2}{3}\right)$$

$$y = \sin^{-1}(\sin \alpha \cos x + \cos \alpha \sin x)$$

$$y = \sin^{-1}(\sin x \cos \alpha + \cos x \sin \alpha)$$

$$y = \sin^{-1}[\sin(x + \alpha)]$$

$$\therefore y = x + \tan^{-1}\left(\frac{2}{3}\right)$$

Differentiate w. r. t. x .

$$\frac{dy}{dx} = \frac{d}{dx} \left[x + \tan^{-1}\left(\frac{2}{3}\right) \right] = 1 + 0$$

$$\therefore \frac{dy}{dx} = 1$$

$$(ii) \text{ Let } y = \cos^{-1}\left(\frac{3 \sin x^2 + 4 \cos x^2}{5}\right)$$

$$= \cos^{-1}\left(\frac{3}{5} \sin x^2 + \frac{4}{5} \cos x^2\right)$$

$$\text{Put } \frac{3}{5} = \sin \alpha, \frac{4}{5} = \cos \alpha$$

$$\text{Also, } \sin^2 \alpha + \cos^2 \alpha = \frac{9}{25} + \frac{16}{25} = 1$$

$$\text{And } \tan \alpha = \frac{3}{4} \therefore \alpha = \tan^{-1}\left(\frac{3}{4}\right)$$

$$y = \cos^{-1}(\sin \alpha \sin x^2 + \cos \alpha \cos x^2)$$

$$y = \cos^{-1}(\cos x^2 \cos \alpha + \sin x^2 \sin \alpha)$$

$$y = \cos^{-1}[\cos(x^2 - \alpha)]$$

$$\therefore y = x^2 - \tan^{-1}\left(\frac{3}{4}\right)$$

Differentiate w. r. t. x .

$$\frac{dy}{dx} = \frac{d}{dx} \left[x^2 - \tan^{-1}\left(\frac{3}{4}\right) \right] = 2x - 0$$

$$\therefore \frac{dy}{dx} = 2x$$

$$(iii) \text{ Let } y = \sin^{-1}\left(\frac{a \cos x - b \sin x}{\sqrt{a^2 + b^2}}\right) = \sin^{-1}\left(\frac{a}{\sqrt{a^2 + b^2}} \cos x - \frac{b}{\sqrt{a^2 + b^2}} \sin x\right)$$

$$\text{Put } \frac{a}{\sqrt{a^2 + b^2}} = \sin \alpha, \frac{b}{\sqrt{a^2 + b^2}} = \cos \alpha$$

$$\text{Also, } \sin^2 \alpha + \cos^2 \alpha = \frac{a^2}{a^2 + b^2} + \frac{b^2}{a^2 + b^2} = 1 \quad \text{And } \tan \alpha = \frac{a}{b} \therefore \alpha = \tan^{-1}\left(\frac{a}{b}\right)$$

$$y = \sin^{-1}(\sin \alpha \cos x - \cos \alpha \sin x)$$

But $\sin(\alpha - x) = \sin \alpha \cos x - \cos \alpha \sin x$

$$y = \sin^{-1}[\sin(\alpha - x)]$$

$$\therefore y = \tan^{-1}\left(\frac{a}{b}\right) - x$$

Differentiate w. r. t. x .

$$\frac{dy}{dx} = \frac{d}{dx} \left[\tan^{-1}\left(\frac{a}{b}\right) - x \right] = 0 - 1 \quad \therefore \frac{dy}{dx} = -1$$

Ex. 5 : Differentiate the following *w. r. t. x*.

(i) $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$

(ii) $\cos^{-1}(2x\sqrt{1-x^2})$

(iii) $\operatorname{cosec}^{-1}\left(\frac{1}{3x-4x^3}\right)$

(iv) $\tan^{-1}\left(\frac{2e^x}{1-e^{2x}}\right)$

(v) $\cos^{-1}\left(\frac{1-9x^2}{1+9x^2}\right)$

(vi) $\cos^{-1}\left(\frac{2^x-2^{-x}}{2^x+2^{-x}}\right)$

(vii) $\tan^{-1}\left(\sqrt{\frac{3-x}{3+x}}\right)$

(viii) $\sin^{-1}\left(\frac{5\sqrt{1-x^2}-12x}{13}\right)$

(ix) $\sin^{-1}\left(\frac{2^{x+1}}{1+4^x}\right)$

Solution :

(i) Let $y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$

Put $x = \tan \theta \therefore \theta = \tan^{-1} x$

$\therefore y = \sin^{-1}\left(\frac{2 \tan \theta}{1 + \tan^2 \theta}\right)$

$y = \sin^{-1}(\sin 2\theta) = 2\theta$

$\therefore y = 2 \tan^{-1} x$

Differentiate *w. r. t. x*.

$$\frac{dy}{dx} = 2 \frac{d}{dx} (\tan^{-1} x)$$

$\therefore \frac{dy}{dx} = \frac{2}{1+x^2}$

(ii) Let $y = \cos^{-1}(2x\sqrt{1-x^2})$

Put $x = \sin \theta \therefore \theta = \sin^{-1} x$

$\therefore y = \cos^{-1}(2 \sin \theta \sqrt{1 - \sin^2 \theta})$

$y = \cos^{-1}(2 \sin \theta \sqrt{\cos^2 \theta})$

$y = \cos^{-1}(2 \sin \theta \cos \theta) = \cos^{-1}(\sin 2\theta)$

$y = \cos^{-1}\left[\cos\left(\frac{\pi}{2} - 2\theta\right)\right] = \frac{\pi}{2} - 2\theta$

$\therefore y = \frac{\pi}{2} - 2 \sin^{-1} x$

Differentiate *w. r. t. x*.

$$\frac{dy}{dx} = \frac{d}{dx} \left(\frac{\pi}{2} - 2 \sin^{-1} x \right)$$

$$\frac{dy}{dx} = 0 - \frac{2 \times 1}{\sqrt{1-x^2}}$$

$\therefore \frac{dy}{dx} = -\frac{2}{\sqrt{1-x^2}}$

(iii) Let $y = \operatorname{cosec}^{-1}\left(\frac{1}{3x-4x^3}\right)$

$y = \sin^{-1}(3x-4x^3)$

Put $x = \sin \theta \therefore \theta = \sin^{-1} x$

$y = \sin^{-1}(3 \sin \theta - 4 \sin^3 \theta)$

$y = \sin^{-1}(\sin 3\theta) = 3\theta$

$\therefore y = 3 \sin^{-1} x$

Differentiate *w. r. t. x*.

$$\frac{dy}{dx} = 3 \frac{d}{dx} (\sin^{-1} x)$$

$\therefore \frac{dy}{dx} = \frac{3}{\sqrt{1-x^2}}$

(iv) Let $y = \tan^{-1}\left(\frac{2e^x}{1-e^{2x}}\right)$

Put $e^x = \tan \theta \therefore \theta = \tan^{-1}(e^x)$

$y = \tan^{-1}\left(\frac{2 \tan \theta}{1 + \tan^2 \theta}\right)$

$y = \tan^{-1}(\tan 2\theta) = 2\theta$

$\therefore y = 2 \tan^{-1}(e^x)$

Differentiate *w. r. t. x*.

$$\frac{dy}{dx} = 2 \frac{d}{dx} [\tan^{-1}(e^x)]$$

$\therefore \frac{dy}{dx} = \frac{2}{1+(e^x)^2} \frac{d}{dx}(e^x) = \frac{2e^x}{1+e^{2x}}$

(v) Let $y = \cos^{-1} \left(\frac{1 - 9x^2}{1 + 9x^2} \right)$

$$y = \cos^{-1} \left(\frac{1 - (3x)^2}{1 + (3x)^2} \right)$$

Put $3x = \tan \theta \therefore \theta = \tan^{-1}(3x)$

$$\therefore y = \cos^{-1} \left[\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right]$$

$$y = \cos^{-1}(\cos 2\theta) = 2\theta$$

$$y = 2 \tan^{-1}(3x)$$

Differentiate *w. r. t. x.*

$$\frac{dy}{dx} = 2 \frac{d}{dx} [\tan^{-1}(3x)]$$

$$\frac{dy}{dx} = \frac{2}{1 + (3x)^2} \frac{d}{dx} (3x)$$

$$\therefore \frac{dy}{dx} = \frac{6}{1 + 9x^2}$$

(vi) Let $y = \cos^{-1} \left(\frac{2^x - 2^{-x}}{2^x + 2^{-x}} \right)$

$$y = \cos^{-1} \left(\frac{2^x(2^x - 2^{-x})}{2^x(2^x + 2^{-x})} \right) \dots \text{[Multiply \& Divide by } 2^x]$$

$$y = \cos^{-1} \left(\frac{2^{2x} - 1}{2^{2x} + 1} \right) = \cos^{-1} \left[-\frac{1 - (2^x)^2}{1 + (2^x)^2} \right]$$

Put $2^x = \tan \theta \therefore \theta = \tan^{-1}(2^x)$

$$\therefore y = \cos^{-1} \left[-\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right] = \cos^{-1}[-\cos 2\theta]$$

$$y = \cos^{-1}[\cos(\pi - 2\theta)] = \pi - 2\theta$$

$$y = \pi - 2 \tan^{-1}(2^x)$$

Differentiate *w. r. t. x.*

$$\frac{dy}{dx} = \frac{d}{dx} [\pi - 2 \tan^{-1}(2^x)]$$

$$\frac{dy}{dx} = 0 - \frac{2}{1 + (2^x)^2} \frac{d}{dx} (2^x) = -\frac{2 \cdot 2^x \cdot \log 2}{1 + 2^{2x}}$$

$$\therefore \frac{dy}{dx} = -\frac{2^{x+1} \log 2}{1 + 2^{2x}}$$

(vii) Let $y = \tan^{-1} \left(\sqrt{\frac{3-x}{3+x}} \right)$

Put $x = 3 \cos 2\theta \therefore \theta = \frac{1}{2} \cos^{-1} \left(\frac{x}{3} \right)$

$$\therefore y = \tan^{-1} \left[\sqrt{\frac{3 - 3 \cos 2\theta}{3 + 3 \cos 2\theta}} \right] = \tan^{-1} \left[\sqrt{\frac{3(1 - \cos 2\theta)}{3(1 + \cos 2\theta)}} \right] = \tan^{-1} \left[\sqrt{\frac{2 \sin^2 \theta}{2 \cos^2 \theta}} \right]$$

$$y = \tan^{-1}(\sqrt{\tan^2 \theta}) = \tan^{-1}(\tan \theta)$$

$$y = \theta = \frac{1}{2} \cos^{-1} \left(\frac{x}{3} \right)$$

Differentiate *w. r. t. x.*

$$\frac{dy}{dx} = \frac{1}{2} \cdot \frac{d}{dx} \left[\cos^{-1} \left(\frac{x}{3} \right) \right]$$

$$= \frac{1}{2} \left[-\frac{1}{\sqrt{1 - \left(\frac{x}{3}\right)^2}} \right] \frac{d}{dx} \left(\frac{x}{3} \right) = -\frac{1}{2} \times \frac{1}{\sqrt{\frac{9-x^2}{9}}} \times \frac{1}{3}$$

$$= -\frac{1}{2} \times \frac{1}{\frac{\sqrt{9-x^2}}{3}} \times \frac{1}{3}$$

$$\therefore \frac{dy}{dx} = -\frac{1}{2\sqrt{9-x^2}}$$

(viii) Let $y = \sin^{-1} \left(\frac{5\sqrt{1-x^2} - 12x}{13} \right)$

Put $x = \sin \theta \therefore \theta = \sin^{-1} x$

$\therefore y = \sin^{-1} \left(\frac{5\sqrt{1-\sin^2\theta} - 12\sin\theta}{13} \right) = \sin^{-1} \left(\frac{5\sqrt{\cos^2\theta} - 12\sin\theta}{13} \right) = \sin^{-1} \left(\frac{5\cos\theta - 12\sin\theta}{13} \right)$

$\therefore y = \sin^{-1} \left(\frac{5}{13} \cos\theta - \frac{12}{13} \sin\theta \right)$

Put $\frac{5}{13} = \sin \alpha, \frac{12}{13} = \cos \alpha$

Also, $\sin^2 \alpha + \cos^2 \alpha = \frac{25}{169} + \frac{144}{169} = 1$

And $\tan \alpha = \frac{5}{12} \therefore \alpha = \tan^{-1} \left(\frac{5}{12} \right)$

$y = \sin^{-1} (\sin \alpha \cos \theta - \cos \alpha \sin \theta) = \sin^{-1} [\sin (\alpha - \theta)] = (\alpha - \theta)$

$\therefore y = \tan^{-1} \left(\frac{5}{12} \right) - \sin^{-1} x$

Differentiate *w. r. t. x.*

$\frac{dy}{dx} = \frac{d}{dx} \left[\tan^{-1} \left(\frac{5}{12} \right) - \sin^{-1} x \right] = 0 - \frac{1}{\sqrt{1-x^2}}$

$\therefore \frac{dy}{dx} = -\frac{1}{\sqrt{1-x^2}}$

(ix) Let $y = \sin^{-1} \left(\frac{2^{x+1}}{1+4^x} \right) = \sin^{-1} \left(\frac{2 \cdot 2^x}{1+(2^x)^2} \right)$

Put $2^x = \tan \theta \therefore \theta = \tan^{-1} (2^x)$

$\therefore y = \sin^{-1} \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) = \sin^{-1} (\sin 2\theta) = 2\theta = 2 \tan^{-1} (2^x)$

Differentiate *w. r. t. x.*

$\frac{dy}{dx} = 2 \frac{d}{dx} [\tan^{-1} (2^x)] = \frac{2}{1+(2^x)^2} \cdot \frac{d}{dx} (2^x) = \frac{2}{1+2^{2x}} (2^x \cdot \log 2)$

$\therefore \frac{dy}{dx} = \frac{2^{x+1} \log 2}{1+4^x}$

Ex. 6 : Differentiate the following *w. r. t. x.*

(i) $\tan^{-1} \left(\frac{4x}{1+21x^2} \right)$

(ii) $\tan^{-1} \left(\frac{7x}{1-12x^2} \right)$

(iii) $\cot^{-1} \left(\frac{b \sin x - a \cos x}{a \sin x + b \cos x} \right)$

(iv) $\tan^{-1} \left(\frac{5x+1}{3-x-6x^2} \right)$

Solution :

$$\begin{aligned} \text{(i) Let } y &= \tan^{-1}\left(\frac{4x}{1+21x^2}\right) \\ &= \tan^{-1}\left(\frac{7x-3x}{1+(7x)(3x)}\right) \\ y &= \tan^{-1}(7x) - \tan^{-1}(3x) \end{aligned}$$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} [\tan^{-1}(7x) - \tan^{-1}(3x)] \\ &= \frac{d}{dx} [\tan^{-1}(7x)] - \frac{d}{dx} [\tan^{-1}(3x)] \\ &= \frac{1}{1+(7x)^2} \cdot \frac{d}{dx}(7x) - \frac{1}{1+(3x)^2} \cdot \frac{d}{dx}(3x) \\ \therefore \frac{dy}{dx} &= \frac{7}{1+49x^2} - \frac{3}{1+9x^2} \end{aligned}$$

$$\begin{aligned} \text{(ii) Let } y &= \tan^{-1}\left(\frac{7x}{1-12x^2}\right) \\ &= \tan^{-1}\left(\frac{3x+4x}{1-(3x)(4x)}\right) \\ y &= \tan^{-1}(3x) + \tan^{-1}(4x) \end{aligned}$$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} [\tan^{-1}(3x) + \tan^{-1}(4x)] \\ &= \frac{d}{dx} [\tan^{-1}(3x)] + \frac{d}{dx} [\tan^{-1}(4x)] \\ &= \frac{1}{1+(3x)^2} \cdot \frac{d}{dx}(3x) + \frac{1}{1+(4x)^2} \cdot \frac{d}{dx}(4x) \\ \therefore \frac{dy}{dx} &= \frac{3}{1+9x^2} + \frac{4}{1+16x^2} \end{aligned}$$

$$\begin{aligned} \text{(iii) Let } y &= \cot^{-1}\left(\frac{b \sin x - a \cos x}{a \sin x + b \cos x}\right) = \tan^{-1}\left(\frac{a \sin x + b \cos x}{b \sin x - a \cos x}\right) = \tan^{-1}\left(\frac{\frac{a}{b} + \cot x}{1 - \left(\frac{a}{b}\right)(\cot x)}\right) \\ &= \tan^{-1}\left(\frac{a}{b}\right) + \tan^{-1}(\cot x) = \tan^{-1}\left(\frac{a}{b}\right) + \tan^{-1}\left[\tan\left(\frac{\pi}{2} - x\right)\right] \\ y &= \tan^{-1}\left(\frac{a}{b}\right) + \frac{\pi}{2} - x \end{aligned}$$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} \left[\tan^{-1}\left(\frac{a}{b}\right) + \frac{\pi}{2} - x \right] \\ &= \frac{d}{dx} \left[\tan^{-1}\left(\frac{a}{b}\right) \right] + \frac{d}{dx} \left(\frac{\pi}{2} \right) - \frac{d}{dx}(x) \\ &= 0 + 0 - 1 \end{aligned}$$

$$\therefore \frac{dy}{dx} = -1$$

$$\begin{aligned} \text{(iv) Let } y &= \tan^{-1}\left(\frac{5x+1}{3-x-6x^2}\right) = \tan^{-1}\left(\frac{5x+1}{1+2-x-6x^2}\right) = \tan^{-1}\left(\frac{5x+1}{1-(6x^2+x-2)}\right) \\ &= \tan^{-1}\left(\frac{5x+1}{1-(6x^2+4x-3x-2)}\right) = \tan^{-1}\left(\frac{5x+1}{1-[2x(3x+2)-(3x+2)]}\right) \\ &= \tan^{-1}\left(\frac{5x+1}{1-(3x+2)(2x-1)}\right) = \tan^{-1}\left(\frac{(3x+2)+(2x-1)}{1-(3x+2)(2x-1)}\right) \end{aligned}$$

$$y = \tan^{-1}(3x + 2) + \tan^{-1}(2x - 1)$$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} [\tan^{-1}(3x + 2) + \tan^{-1}(2x - 1)] \\ &= \frac{d}{dx} [\tan^{-1}(3x + 2)] + \frac{d}{dx} [\tan^{-1}(2x - 1)] \\ &= \frac{1}{1 + (3x + 2)^2} \cdot \frac{d}{dx} (3x + 2) + \frac{1}{1 + (2x - 1)^2} \cdot \frac{d}{dx} (2x - 1) \\ \therefore \frac{dy}{dx} &= \frac{3}{1 + (3x + 2)^2} + \frac{2}{1 + (2x - 1)^2} \end{aligned}$$

EXERCISE 1.2

- (1) Find the derivative of the function $y = f(x)$ using the derivative of the inverse function $x = f^{-1}(y)$ in the following
- (i) $y = \sqrt{x}$ (ii) $y = \sqrt{2 - \sqrt{x}}$
 (iii) $y = \sqrt[3]{x-2}$ (iv) $y = \log(2x - 1)$
 (v) $y = 2x + 3$ (vi) $y = e^x - 3$
 (vii) $y = e^{2x-3}$ (viii) $y = \log_2\left(\frac{x}{2}\right)$
- (2) Find the derivative of the inverse function of the following
- (i) $y = x^2 \cdot e^x$ (ii) $y = x \cos x$
 (iii) $y = x \cdot 7^x$ (iv) $y = x^2 + \log x$
 (v) $y = x \log x$
- (3) Find the derivative of the inverse of the following functions, and also find their value at the points indicated against them.
- (i) $y = x^5 + 2x^3 + 3x$, at $x = 1$
 (ii) $y = e^x + 3x + 2$, at $x = 0$
 (iii) $y = 3x^2 + 2 \log x^3$, at $x = 1$
 (iv) $y = \sin(x - 2) + x^2$, at $x = 2$
- (4) If $f(x) = x^3 + x - 2$, find $(f^{-1})'(0)$.
- (5) Using derivative prove
- (i) $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$
 (ii) $\sec^{-1} x + \operatorname{cosec}^{-1} x = \frac{\pi}{2} \dots$ [for $|x| \geq 1$]
- (6) Differentiate the following w. r. t. x.
- (i) $\tan^{-1}(\log x)$ (ii) $\operatorname{cosec}^{-1}(e^{-x})$
 (iii) $\cot^{-1}(x^3)$ (iv) $\cot^{-1}(4^x)$
 (v) $\tan^{-1}(\sqrt{x})$ (vi) $\sin^{-1}\left(\sqrt{\frac{1+x^2}{2}}\right)$
 (vii) $\cos^{-1}(1 - x^2)$ (viii) $\sin^{-1}(x^{\frac{3}{2}})$
 (ix) $\cos^3[\cos^{-1}(x^3)]$ (x) $\sin^4[\sin^{-1}(\sqrt{x})]$
- (7) Differentiate the following w. r. t. x.
- (i) $\cot^{-1}[\cot(e^{x^2})]$
 (ii) $\operatorname{cosec}^{-1}\left(\frac{1}{\cos(5^x)}\right)$
 (iii) $\cos^{-1}\left(\sqrt{\frac{1 + \cos x}{2}}\right)$
 (iv) $\cos^{-1}\left(\sqrt{\frac{1 - \cos(x^2)}{2}}\right)$
 (v) $\tan^{-1}\left(\frac{1 - \tan(\frac{x}{2})}{1 + \tan(\frac{x}{2})}\right)$
 (vi) $\operatorname{cosec}^{-1}\left(\frac{1}{4 \cos^3 2x - 3 \cos 2x}\right)$
 (vii) $\tan^{-1}\left(\frac{1 + \cos(\frac{x}{3})}{\sin(\frac{x}{3})}\right)$
 (viii) $\cot^{-1}\left(\frac{\sin 3x}{1 + \cos 3x}\right)$

(ix) $\tan^{-1}\left(\frac{\cos 7x}{1 + \sin 7x}\right)$

(x) $\tan^{-1}\left(\frac{\sqrt{1 + \cos x}}{\sqrt{1 - \cos x}}\right)$

(xi) $\tan^{-1}(\operatorname{cosec} x + \cot x)$

(xii) $\cot^{-1}\left(\frac{\sqrt{1 + \sin\left(\frac{4x}{3}\right)} + \sqrt{1 - \sin\left(\frac{4x}{3}\right)}}{\sqrt{1 + \sin\left(\frac{4x}{3}\right)} - \sqrt{1 - \sin\left(\frac{4x}{3}\right)}}\right)$

(iii) $\sin^{-1}\left(\frac{1 - x^2}{1 + x^2}\right)$ (iv) $\sin^{-1}(2x\sqrt{1 - x^2})$

(v) $\cos^{-1}(3x - 4x^3)$ (vi) $\cos^{-1}\left(\frac{e^x - e^{-x}}{e^x + e^{-x}}\right)$

(vii) $\cos^{-1}\left(\frac{1 - 9^x}{1 + 9^x}\right)$ (viii) $\sin^{-1}\left(\frac{4^{x+\frac{1}{2}}}{1 + 2^{4x}}\right)$

(ix) $\sin^{-1}\left(\frac{1 - 25x^2}{1 + 25x^2}\right)$ (x) $\sin^{-1}\left(\frac{1 - x^3}{1 + x^3}\right)$

(xi) $\tan^{-1}\left(\frac{2x^{\frac{5}{2}}}{1 - x^5}\right)$ (xii) $\cot^{-1}\left(\frac{1 - \sqrt{x}}{1 + \sqrt{x}}\right)$

(8) Differentiate the following w. r. t. x.

(i) $\sin^{-1}\left(\frac{4 \sin x + 5 \cos x}{\sqrt{41}}\right)$

(ii) $\cos^{-1}\left(\frac{\sqrt{3} \cos x - \sin x}{2}\right)$

(iii) $\sin^{-1}\left(\frac{\cos \sqrt{x} + \sin \sqrt{x}}{\sqrt{2}}\right)$

(iv) $\cos^{-1}\left(\frac{3 \cos 3x - 4 \sin 3x}{5}\right)$

(v) $\cos^{-1}\left(\frac{3 \cos(e^x) + 2 \sin(e^x)}{\sqrt{13}}\right)$

(vi) $\operatorname{cosec}^{-1}\left(\frac{10}{6 \sin(2^x) - 8 \cos(2^x)}\right)$

(9) Differentiate the following w. r. t. x.

(i) $\cos^{-1}\left(\frac{1 - x^2}{1 + x^2}\right)$ (ii) $\tan^{-1}\left(\frac{2x}{1 - x^2}\right)$

(10) Differentiate the following w. r. t. x.

(i) $\tan^{-1}\left(\frac{8x}{1 - 15x^2}\right)$ (ii) $\cot^{-1}\left(\frac{1 + 35x^2}{2x}\right)$

(iii) $\tan^{-1}\left(\frac{2\sqrt{x}}{1 + 3x}\right)$ (iv) $\tan^{-1}\left(\frac{2^{x+2}}{1 - 3(4^x)}\right)$

(v) $\tan^{-1}\left(\frac{2^x}{1 + 2^{2x+1}}\right)$ (vi) $\cot^{-1}\left(\frac{a^2 - 6x^2}{5ax}\right)$

(vii) $\tan^{-1}\left(\frac{a + b \tan x}{b - a \tan x}\right)$

(viii) $\tan^{-1}\left(\frac{5 - x}{6x^2 - 5x - 3}\right)$

(ix) $\cot^{-1}\left(\frac{4 - x - 2x^2}{3x + 2}\right)$

1.3.1 Logarithmic Differentiation

The complicated functions given by formulas that involve products, quotients and powers can often be simplified more quickly by taking the natural logarithms on both the sides. This enables us to use the laws of logarithms to simplify the functions and differentiate easily. Especially when the functions are of the form $y = [f(x)]^{g(x)}$ it is recommended to take logarithms on both the sides which simplifies to $\log y = g(x) \cdot \log [f(x)]$, now it becomes convenient to find the derivative. This process of finding the derivative is called logarithmic differentiation.



SOLVED EXAMPLES

Ex. 1 : Differentiate the following w. r. t. x.

(i) $\left(\frac{(x^2 + 3)^2 \sqrt[3]{(x^3 + 5)^2}}{\sqrt{(2x^2 + 1)^3}}\right)$

(ii) $\frac{e^{x^2} (\tan x)^{\frac{x}{2}}}{(1 + x^2)^{\frac{3}{2}} \cos^3 x}$

(iii) $(x + 1)^{\frac{3}{2}} (2x + 3)^{\frac{5}{2}} (3x + 4)^{\frac{2}{3}}$ for $x \geq 0$ (iv) $x^a + x^x + a^x$ (v) $(\sin x)^{\tan x} - x^{\log x}$

Solution :

(i) Let $y = \left(\frac{(x^2 + 3)^2 \sqrt[3]{(x^3 + 5)^2}}{\sqrt{(2x^2 + 1)^3}} \right)$

Taking log of both the sides we get,

$$\log y = \log \left(\frac{(x^2 + 3)^2 \sqrt[3]{(x^3 + 5)^2}}{\sqrt{(2x^2 + 1)^3}} \right) = \log \left[\frac{(x^2 + 3)^2 (x^3 + 5)^{\frac{2}{3}}}{(2x^2 + 1)^{\frac{3}{2}}} \right]$$

$$= \log [(x^2 + 3)^2 (x^3 + 5)^{\frac{2}{3}}] - \log (2x^2 + 1)^{\frac{3}{2}}$$

$$= \log (x^2 + 3)^2 + \log (x^3 + 5)^{\frac{2}{3}} - \log (2x^2 + 1)^{\frac{3}{2}}$$

$$\log y = 2 \log (x^2 + 3) + \frac{2}{3} \log (x^3 + 5) - \frac{3}{2} \log (2x^2 + 1)$$

Differentiate w. r. t. x.

$$\frac{d}{dx} (\log y) = \frac{d}{dx} \left[2 \log (x^2 + 3) + \frac{2}{3} \log (x^3 + 5) - \frac{3}{2} \log (2x^2 + 1) \right]$$

$$\frac{1}{y} \frac{dy}{dx} = 2 \cdot \frac{d}{dx} [\log (x^2 + 3)] + \frac{2}{3} \cdot \frac{d}{dx} [\log (x^3 + 5)] - \frac{3}{2} \cdot \frac{d}{dx} [\log (2x^2 + 1)]$$

$$= \frac{2}{x^2 + 3} \cdot \frac{d}{dx} (x^2 + 3) + \frac{2}{3(x^3 + 5)} \cdot \frac{d}{dx} (x^3 + 5) - \frac{3}{2(2x^2 + 1)} \cdot \frac{d}{dx} (2x^2 + 1)$$

$$\frac{dy}{dx} = y \left[\frac{2}{x^2 + 3} (2x) + \frac{2}{3(x^3 + 5)} (3x^2) - \frac{3}{2(2x^2 + 1)} (4x) \right]$$

$$\therefore \frac{dy}{dx} = \frac{(x^2 + 3)^2 \sqrt[3]{(x^3 + 5)^2}}{\sqrt{(2x^2 + 1)^3}} \left[\frac{4x}{x^2 + 3} + \frac{2x^2}{(x^3 + 5)} - \frac{6x}{2x^2 + 1} \right]$$

(ii) Let $y = \frac{e^{x^2} (\tan x)^{\frac{x}{2}}}{(1 + x^2)^{\frac{3}{2}} \cos^3 x}$

Taking log of both the sides we get,

$$\log y = \log \left(\frac{e^{x^2} (\tan x)^{\frac{x}{2}}}{(1 + x^2)^{\frac{3}{2}} (\cos x)^3} \right) = \log [e^{x^2} (\tan x)^{\frac{x}{2}}] - \log [(1 + x^2)^{\frac{3}{2}} (\cos x)^3]$$

$$= \log e^{x^2} + \log (\tan x)^{\frac{x}{2}} - [\log (1 + x^2)^{\frac{3}{2}} + \log (\cos x)^3]$$

$$= x^2 \log e + \frac{x}{2} \log (\tan x) - \frac{3}{2} \log (1 + x^2) - 3 \log (\cos x)$$

$$\therefore \log y = x^2 + \frac{x}{2} \log (\tan x) - \frac{3}{2} \log (1 + x^2) - 3 \log (\cos x)$$



Differentiate w. r. t. x.

$$\begin{aligned} \frac{d}{dx} (\log y) &= \frac{d}{dx} \left[x^2 + \frac{x}{2} \log (\tan x) - \frac{3}{2} \log (1+x^2) - 3 \log (\cos x) \right] \\ \frac{1}{y} \frac{dy}{dx} &= \frac{d}{dx} (x^2) + \frac{x}{2} \cdot \frac{d}{dx} [\log (\tan x)] + \log (\tan x) \frac{d}{dx} \left(\frac{x}{2} \right) - \frac{3}{2} \frac{d}{dx} [\log (1+x^2)] - 3 \frac{d}{dx} [\log (\cos x)] \\ &= 2x + \frac{x}{2} \cdot \frac{1}{\tan x} \cdot \frac{d}{dx} (\tan x) + \log (\tan x) \frac{1}{2} - \frac{3}{2(1+x^2)} \cdot \frac{d}{dx} (1+x^2) - \frac{3}{\cos x} \cdot \frac{d}{dx} (\cos x) \\ &= 2x + \frac{x}{2} \cdot (\cot x) (\sec^2 x) + \frac{1}{2} \log (\tan x) - \frac{3}{2(1+x^2)} \cdot (2x) - \frac{3}{\cos x} (-\sin x) \\ &= 2x + \frac{x}{2} \times \frac{\cos x}{\sin x} \times \frac{1}{\cos^2 x} + \frac{1}{2} \log (\tan x) - \frac{3x}{1+x^2} + 3 \tan x \\ \frac{dy}{dx} &= y \left[2x + \frac{x}{2 \sin x \cos x} + \frac{1}{2} \log (\tan x) - \frac{3x}{1+x^2} + 3 \tan x \right] \\ \therefore \frac{dy}{dx} &= \frac{e^{x^2} (\tan x)^{\frac{x}{2}}}{(1+x^2)^{\frac{3}{2}} \cos^3 x} \left[2x + x \operatorname{cosec} 2x + \frac{1}{2} \log (\tan x) - \frac{3x}{1+x^2} + 3 \tan x \right] \end{aligned}$$

(iii) Let $y = (x+1)^{\frac{3}{2}} (2x+3)^{\frac{5}{2}} (3x+4)^{\frac{2}{3}}$

Taking log of both the sides we get,

$$\begin{aligned} \log y &= \log \left[(x+1)^{\frac{3}{2}} (2x+3)^{\frac{5}{2}} (3x+4)^{\frac{2}{3}} \right] \\ &= \log (x+1)^{\frac{3}{2}} + \log (2x+3)^{\frac{5}{2}} + \log (3x+4)^{\frac{2}{3}} \\ \log y &= \frac{3}{2} \log (x+1) + \frac{5}{2} \log (2x+3) + \frac{2}{3} \log (3x+4) \end{aligned}$$

Differentiate w. r. t. x.

$$\begin{aligned} \frac{d}{dx} (\log y) &= \frac{d}{dx} \left[\frac{3}{2} \log (x+1) + \frac{5}{2} \log (2x+3) + \frac{2}{3} \log (3x+4) \right] \\ \frac{1}{y} \cdot \frac{dy}{dx} &= \frac{3}{2} \cdot \frac{d}{dx} [\log (2x+1)] + \frac{5}{2} \cdot \frac{d}{dx} [\log (3x+2)] + \frac{2}{3} \cdot \frac{d}{dx} [\log (3x+4)] \\ &= \frac{3}{2(2x+1)} \cdot \frac{d}{dx} (2x+1) + \frac{5}{2(3x+1)} \cdot \frac{d}{dx} (3x+2) + \frac{2}{3(3x+4)} \cdot \frac{d}{dx} (3x+4) \\ \frac{dy}{dx} &= y \left[\frac{3}{2(2x+1)} (2) + \frac{5}{2(3x+1)} (3) + \frac{2}{3(3x+4)} (3) \right] \\ \therefore \frac{dy}{dx} &= (x+1)^{\frac{3}{2}} (2x+3)^{\frac{5}{2}} (3x+4)^{\frac{2}{3}} \left[\frac{3}{2x+1} + \frac{15}{2(3x+1)} + \frac{2}{3x+4} \right] \end{aligned}$$

(iv) Let $y = x^a + x^x + a^x$

Here the derivatives of x^a and a^x can be found directly but we can not find the derivative of x^x without the use of logarithm. So the given function is split in to two functions, find their derivatives and then add them.

Let $u = x^a + a^x$ and $v = x^x$

$\therefore y = u + v$, where u and v are differentiable functions of x .

$$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \quad \dots \dots (I)$$

Now, $u = x^a + a^x$

Differentiate *w. r. t. x*.

$$\begin{aligned} \frac{du}{dx} &= \frac{d}{dx}(x^a) + \frac{d}{dx}(a^x) \\ \frac{du}{dx} &= ax^{a-1} + a^x \log a \quad \dots \dots (II) \end{aligned}$$

And, $v = x^x$

Taking log of both the sides we get,

$$\log v = \log x^x$$

$$\log v = x \log x$$

Differentiate *w. r. t. x*.

$$\begin{aligned} \frac{d}{dx}(\log v) &= \frac{d}{dx}(x \log x) \\ \frac{1}{v} \frac{dv}{dx} &= x \frac{d}{dx}(\log x) + \log x \frac{d}{dx}(x) \\ \frac{dv}{dx} &= v \left[x \times \frac{1}{x} + \log x (1) \right] \\ \frac{dv}{dx} &= x^x [1 + \log x] \quad \dots \dots (III) \end{aligned}$$

Substituting (II) and (III) in (I) we get,

$$\frac{dy}{dx} = ax^{a-1} + a^x \log a + x^x [1 + \log x]$$

(v) Let $y = (\sin x)^{\tan x} - x^{\log x}$

Let $u = (\sin x)^{\tan x}$ and $v = x^{\log x}$

$\therefore y = u - v$, where u and v are differentiable functions of x .

$$\frac{dy}{dx} = \frac{du}{dx} - \frac{dv}{dx} \quad \dots \dots (I)$$

Now, $u = (\sin x)^{\tan x}$, taking log of both the sides we get,

$$\log u = \log (\sin x)^{\tan x} \quad \therefore \log u = \tan x \log (\sin x)$$

Differentiate *w. r. t. x*.

$$\begin{aligned} \frac{d}{dx}(\log u) &= \frac{d}{dx} [\tan x \log (\sin x)] \\ \frac{1}{u} \frac{du}{dx} &= \tan x \frac{d}{dx} [\log (\sin x)] + \log (\sin x) \frac{d}{dx} (\tan x) \\ &= \tan x \cdot \frac{1}{\sin x} \cdot \frac{d}{dx} (\sin x) + \log (\sin x) \cdot (\sec^2) \end{aligned}$$

$$\begin{aligned} \frac{du}{dx} &= u \left[\tan x \cdot \frac{1}{\sin x} \cdot (\cos x) + \sec^2 x \cdot \log(\sin x) \right] \\ \frac{du}{dx} &= (\sin x)^{\tan x} [\tan x \cdot \cot x + \sec^2 x \cdot \log(\sin x)] \\ \frac{du}{dx} &= (\sin x)^{\tan x} [1 + \sec^2 x \cdot \log(\sin x)] \end{aligned} \dots\dots \text{(II)}$$

And, $v = x^{\log x}$

Taking log on both the sides we get,

$$\begin{aligned} \log v &= \log(x^{\log x}) \\ \log v &= \log x \log x = (\log x)^2 \end{aligned}$$

Differentiate *w. r. t. x*.

$$\begin{aligned} \frac{d}{dx}(\log v) &= \frac{d}{dx}[(\log x)^2] \\ \frac{1}{v} \frac{dv}{dx} &= 2 \log x \frac{d}{dx}(\log x) \\ \frac{dv}{dx} &= u \left[\frac{2 \log x}{x} \right] = \frac{2x^{\log x} \log x}{x} \end{aligned} \dots\dots \text{(III)}$$

Substituting (II) and (III) in (I) we get,

$$\frac{dy}{dx} = (\sin x)^{\tan x} [1 + \sec^2 x \cdot \log(\sin x)] - \frac{2x^{\log x} \log x}{x}$$


1.3.2 Implicit Functions

Functions can be represented in a variety of ways. Most of the functions we have dealt with so far have been described by an equation of the form $y = f(x)$ that expresses y solely in terms of the variable x . It is not always possible to solve for one variable explicitly in terms of another. Those cases where it is possible to solve for one variable in terms of another to obtain $y = f(x)$ or $x = g(y)$ are said to be in **explicit** form.

If an equation in x and y is given but x is not an explicit function of y and y is not an explicit function of x then either of the variables is an **Implicit function** of the other.

1.3.3 Derivatives of Implicit Functions

1. Differentiate both sides of the equation with respect to x (independent variable), treating y as a differentiable function of x .
2. Collect the terms containing $\frac{dy}{dx}$ on one side of the equation and solve for $\frac{dy}{dx}$.



SOLVED EXAMPLES

Ex. 1 : Find $\frac{dy}{dx}$ if

(i) $x^5 + xy^3 + x^2y + y^4 = 4$

(ii) $y^3 + \cos(xy) = x^2 - \sin(x + y)$

(iii) $x^2 + e^{xy} = y^2 + \log(x + y)$

Solution :

(i) Given that : $x^5 + xy^3 + x^2y + y^4 = 4$

Differentiate w. r. t. x.

$$\frac{d}{dx}(x^5) + \frac{d}{dx}(xy^3) + \frac{d}{dx}(x^2y) + \frac{d}{dx}(y^4) = \frac{d}{dx}(4)$$

$$5x^4 + x \frac{d}{dx}(y^3) + y^3 \frac{d}{dx}(x) + x^2 \frac{d}{dx}(y) + y \frac{d}{dx}(x^2) + 4y^3 \frac{d}{dx}(y) = 0$$

$$5x^4 + x(3y^2) \frac{dy}{dx} + y^3(1) + x^2 \frac{dy}{dx} + y(2x) + 4y^3 \frac{dy}{dx} = 0$$

$$x^2 \frac{dy}{dx} + 3xy^2 \frac{dy}{dx} + 4y^3 \frac{dy}{dx} = -5x^4 - 2xy - y^3$$

$$(x^2 + 3xy^2 + 4y^3) \frac{dy}{dx} = -(5x^4 + 2xy + y^3)$$

$$\therefore \frac{dy}{dx} = -\frac{5x^4 + 2xy + y^3}{x^2 + 3xy^2 + 4y^3}$$

(ii) Given that : $y^3 + \cos(xy) = x^2 - \sin(x + y)$

Differentiate w. r. t. x.

$$\frac{d}{dx}(y^3) + \frac{d}{dx}[\cos(xy)] = \frac{d}{dx}(x^2) - \frac{d}{dx}[\sin(x + y)]$$

$$3y^2 \frac{d}{dx}(y) - \sin(xy) \frac{d}{dx}(xy) = 2x - \cos(x + y) \frac{d}{dx}(x + y)$$

$$3y^2 \frac{dy}{dx} - \sin(xy) \left[x \frac{dy}{dx} + y(1) \right] = 2x - \cos(x + y) \left[1 + \frac{dy}{dx} \right]$$

$$3y^2 \frac{dy}{dx} - x \sin(xy) \frac{dy}{dx} - y \sin(xy) = 2x - \cos(x + y) - \cos(x + y) \frac{dy}{dx}$$

$$3y^2 \frac{dy}{dx} - x \sin(xy) \frac{dy}{dx} + \cos(x + y) \frac{dy}{dx} = 2x + y \sin(xy) - \cos(x + y)$$

$$[3y^2 - x \sin(xy) + \cos(x + y)] \frac{dy}{dx} = 2x + y \sin(xy) - \cos(x + y)$$

$$\therefore \frac{dy}{dx} = \frac{2x + y \sin(xy) - \cos(x + y)}{3y^2 - x \sin(xy) + \cos(x + y)}$$

(iii) Given that : $x^2 + e^{xy} = y^2 + \log(x + y)$

Recall that $\frac{d}{dx} g(f(x)) = g'(f(x)) \cdot \frac{d}{dx} f(x)$

Differentiate w. r. t. x .

$$\frac{d}{dx} (x^2) + \frac{d}{dx} [e^{xy}] = \frac{d}{dx} (y^2) + \frac{d}{dx} [\log(x + y)]$$

$$2x + e^{xy} \frac{d}{dx} (xy) = 2y \frac{dy}{dx} + \frac{1}{x + y} \frac{d}{dx} (x + y)$$

$$2x + e^{xy} \left[x \frac{dy}{dx} + y(1) \right] = 2y \frac{dy}{dx} + \frac{1}{x + y} \left[1 + \frac{dy}{dx} \right]$$

$$2x + xe^{xy} \frac{dy}{dx} + ye^{xy} = 2y \frac{dy}{dx} + \frac{1}{x + y} + \frac{1}{x + y} \cdot \frac{dy}{dx}$$

$$2x + ye^{xy} - \frac{1}{x + y} = 2y \frac{dy}{dx} - xe^{xy} \frac{dy}{dx} + \frac{1}{x + y} \cdot \frac{dy}{dx}$$

$$2x + ye^{xy} - \frac{1}{x + y} = \left[2y - xe^{xy} + \frac{1}{x + y} \right] \frac{dy}{dx}$$

$$\frac{2x(x + y) + ye^{xy}(x + y) - 1}{x + y} = \left[\frac{2y(x + y) - xe^{xy}(x + y) + 1}{x + y} \right] \frac{dy}{dx}$$

$$\therefore \frac{dy}{dx} = \frac{2x(x + y) + ye^{xy}(x + y) - 1}{2y(x + y) - xe^{xy}(x + y) + 1}$$

Ex. 2 : Find $x^m \cdot y^n = (x + y)^{m+n}$, then prove that $\frac{dy}{dx} = \frac{y}{x}$.

Solution : Given that : $x^m \cdot y^n = (x + y)^{m+n}$

Taking log on both the sides, we get

$$\log [x^m \cdot y^n] = \log [(x + y)^{m+n}]$$

$$m \log x + n \log y = (m + n) \log (x + y)$$

Differentiate w. r. t. x .

$$m \frac{d}{dx} (\log x) + n \frac{d}{dx} (\log y) = (m + n) \frac{d}{dx} [\log (x + y)]$$

$$\frac{m}{x} + \frac{n}{y} \cdot \frac{dy}{dx} = \frac{m + n}{x + y} \cdot \frac{d}{dx} (x + y)$$

$$\frac{m}{x} + \frac{n}{y} \cdot \frac{dy}{dx} = \frac{m + n}{x + y} \cdot \left[1 + \frac{dy}{dx} \right]$$

$$\frac{m}{x} + \frac{n}{y} \cdot \frac{dy}{dx} = \frac{m + n}{x + y} + \frac{m + n}{x + y} \cdot \frac{dy}{dx}$$

$$\frac{n}{y} \cdot \frac{dy}{dx} - \frac{m + n}{x + y} \cdot \frac{dy}{dx} = \frac{m + n}{x + y} - \frac{m}{x}$$

$$\left[\frac{n}{y} - \frac{m+n}{x+y} \right] \frac{dy}{dx} = \frac{m+n}{x+y} - \frac{m}{x}$$

$$\left[\frac{n(x+y) - (m+n)y}{y(x+y)} \right] \frac{dy}{dx} = \left[\frac{(m+n)x - m(x+y)}{x(x+y)} \right]$$

$$\left[\frac{nx + ny - my - ny}{y} \right] \frac{dy}{dx} = \frac{mx + nx - mx - my}{x}$$

$$\left[\frac{nx - my}{y} \right] \frac{dy}{dx} = \frac{nx - my}{x}$$

$$\therefore \frac{dy}{dx} = \frac{y}{x}$$

Ex. 3 : If $\sin\left(\frac{px^m - qy^m}{px^m + qy^m}\right) = r$, then show that $\frac{dy}{dx} = \frac{y}{x}$, where r is a constant.

Solution : Given that : $\sin\left(\frac{px^m - qy^m}{px^m + qy^m}\right) = r$

$$\frac{px^m - qy^m}{px^m + qy^m} = \sin^{-1} r$$

$$\frac{px^m - qy^m}{px^m + qy^m} = t \quad \dots\dots [\text{Let } t = \sin^{-1} r]$$

$$px^m - qy^m = ptx^m + qty^m$$

$$px^m - ptx^m = qy^m + qty^m$$

$$p(1-t)x^m = q(1+t)y^m$$

$$y^m = \left(\frac{p(1-t)}{q(1+t)} \right) x^m$$

$$y^m = s \cdot x^m \quad \dots\dots (I) \quad \dots\dots \left[\text{Let } s = \left(\frac{p(1-t)}{q(1+t)} \right) \right]$$

Differentiate w. r. t. x

$$\frac{d}{dx} (y^m) = s \frac{d}{dx} (x^m)$$

$$my^{m-1} \frac{dy}{dx} = s \cdot mx^{m-1}$$

$$\frac{dy}{dx} = s \cdot \frac{x^{m-1}}{y^{m-1}} x^{m-1}$$

$$\frac{dy}{dx} = \frac{y^m}{x^m} \times \frac{x^{m-1}}{y^{m-1}} \quad \dots\dots [\text{From (I)}]$$

$$\therefore \frac{dy}{dx} = \frac{y}{x}$$

Ex. 4 : If $\sec^{-1} \left(\frac{x^3 + y^3}{x^3 - y^3} \right) = 2a$, then show that $\frac{dy}{dx} = \frac{x^2 \tan^2 a}{y^2}$, where a is a constant.

Solution : Given that : $\sec^{-1} \left(\frac{x^3 + y^3}{x^3 - y^3} \right) = 2a$ [We will not eliminate a , as answer contains a]

$$\therefore \cos^{-1} \left(\frac{x^3 - y^3}{x^3 + y^3} \right) = 2a$$

$$\frac{x^3 - y^3}{x^3 + y^3} = \cos 2a$$

$$x^3 - y^3 = x^3 \cos 2a + y^3 \cos 2a$$

$$x^3 - x^3 \cos 2a = y^3 \cos 2a + y^3$$

$$x^3 (1 - \cos 2a) = y^3 (1 + \cos 2a)$$

$$y^3 = \left(\frac{1 - \cos 2a}{1 + \cos 2a} \right) x^3$$

$$y^3 = \left(\frac{2 \sin^2 a}{2 \cos^2 a} \right) x^3$$

$$y^3 = (\tan^2 a) x^3 \quad \dots \dots \dots \text{(I)}$$

Differentiate *w. r. t. x*

$$\frac{d}{dx} (y^3) = (\tan^2 a) \frac{d}{dx} (x^3)$$

$$3y^2 \frac{dy}{dx} = (\tan^2 a) 3x^2$$

$$\therefore \frac{dy}{dx} = \frac{x^2 \tan^2 a}{y^2}$$

Ex. 5 : If $y = \sqrt{\tan x + \sqrt{\tan x + \sqrt{\tan x + \dots \infty}}}$, then show that $\frac{dy}{dx} = \frac{\sec^2 x}{2y - 1}$.

Solution : Given that : $y = \sqrt{\tan x + \sqrt{\tan x + \sqrt{\tan x + \dots \infty}}}$ (I)

Squaring both sides, we get


$$y^2 = \tan x + \sqrt{\tan x + \sqrt{\tan x + \dots \infty}}, \text{ which is same as}$$

$$y^2 = \tan x + \sqrt{\tan x + \sqrt{\tan x + \sqrt{\tan x + \dots \infty}}}$$

$$y^2 = \tan x + y \quad \dots \dots \dots \text{ [From (I)]}$$

Differentiate *w. r. t. x*

$$\frac{d}{dx} (y^2) = \frac{d}{dx} (\tan x) + \frac{dy}{dx}$$



$$2y \frac{dy}{dx} - \frac{dy}{dx} = \sec^2 x$$

$$(2y - 1) \frac{dy}{dx} = \sec^2 x$$

$$\therefore \frac{dy}{dx} = \frac{\sec^2 x}{2y - 1}$$

Ex. 6 : If $\sqrt{1 - x^2} + \sqrt{1 - y^2} = a(x - y)$, then show that $\frac{dy}{dx} = \sqrt{\frac{1 - y^2}{1 - x^2}}$.

Solution : Given that : $\sqrt{1 - x^2} + \sqrt{1 - y^2} = a(x - y)$ (I)

Put $x = \sin \alpha, y = \sin \beta$

$\therefore \alpha = \sin^{-1} x, \beta = \sin^{-1} y$

Equation (I) becomes,

$$\sqrt{1 - \sin^2 \alpha} + \sqrt{1 - \sin^2 \beta} = a(\sin \alpha - \sin \beta)$$

$$\cos \alpha + \cos \beta = a(\sin \alpha - \sin \beta)$$

$$2 \cos \left(\frac{\alpha + \beta}{2} \right) \cos \left(\frac{\alpha - \beta}{2} \right) = 2a \cos \left(\frac{\alpha + \beta}{2} \right) \sin \left(\frac{\alpha - \beta}{2} \right)$$

$$\cos \left(\frac{\alpha - \beta}{2} \right) = a \sin \left(\frac{\alpha + \beta}{2} \right) \Rightarrow \cot \left(\frac{\alpha - \beta}{2} \right) = a$$

$$\frac{\alpha - \beta}{2} = \cot^{-1} a \quad \therefore \alpha - \beta = 2 \cot^{-1} a$$

$$\therefore \sin^{-1} x - \sin^{-1} y = 2 \cot^{-1} a$$

Differentiate w. r. t. x

$$\frac{d}{dx} (\sin^{-1} x) - \frac{d}{dx} (\sin^{-1} y) = \frac{d}{dx} (2 \cot^{-1} a)$$

$$\frac{1}{\sqrt{1 - x^2}} - \frac{1}{\sqrt{1 - y^2}} \cdot \frac{dy}{dx} = 0$$

$$\therefore \frac{dy}{dx} = \sqrt{\frac{1 - y^2}{1 - x^2}}$$

EXERCISE 1.3

(1) Differentiate the following w. r. t. x

(i) $\frac{(x + 1)^2}{(x + 2)^3 (x + 3)^4}$

(ii) $\sqrt[3]{\frac{4x - 1}{(2x + 3)(5 - 2x)^2}}$

(iii) $(x^2 + 3)^{\frac{3}{2}} \cdot \sin^3 2x \cdot 2^{x^2}$

(iv) $\frac{(x^2 + 2x + 2)^{\frac{3}{2}}}{(\sqrt{x} + 3)^3 (\cos x)^x}$

(v) $\frac{x^5 \cdot \tan^3 4x}{\sin^2 3x}$

(vii) $(\sin x)^x$

(vi) $x^{\tan^{-1} x}$

(viii) $\sin x^x$

(2) Differentiate the following w. r. t. x .

- (i) $x^e + x^x + e^x + e^e$ (ii) $x^{x^x} + e^{x^x}$
 (iii) $(\log x)^x - (\cos x)^{\cot x}$
 (iv) $x^{e^x} + (\log x)^{\sin x}$ (v) $e^{\tan x} + (\log x)^{\tan x}$
 (vi) $(\sin x)^{\tan x} + (\cos x)^{\cot x}$
 (vii) $10^{x^x} + x^{x^{10}} + x^{10^x}$
 (viii) $[(\tan x)^{\tan x}]^{\tan x}$ at $x = \frac{\pi}{4}$

(3) Find $\frac{dy}{dx}$ if

- (i) $\sqrt{x} + \sqrt{y} = \sqrt{a}$ (ii) $x\sqrt{x} + y\sqrt{y} = a\sqrt{a}$
 (iii) $x + \sqrt{xy} + y = 1$
 (iv) $x^3 + x^2y + xy^2 + y^3 = 81$
 (v) $x^2y^2 - \tan^{-1}\sqrt{x^2 + y^2} = \cot^{-1}\sqrt{x^2 + y^2}$
 (vi) $xe^y + ye^x = 1$ (vii) $e^{x+y} = \cos(x-y)$
 (viii) $\cos(xy) = x + y$ (ix) $e^{e^{x-y}} = \frac{x}{y}$
 (x) $x + \sin(x+y) = y - \cos(x-y)$

(4) Show that $\frac{dy}{dx} = \frac{y}{x}$ in the following,

where a and p are constants.

- (i) $x^7y^5 = (x+y)^{12}$
 (ii) $x^p y^4 = (x+y)^{p+4}, p \in \mathbb{N}$
 (iii) $\sec\left(\frac{x^5 + y^5}{x^5 - y^5}\right) = a^2$
 (iv) $\tan^{-1}\left(\frac{3x^2 - 4y^2}{3x^2 + 4y^2}\right) = a^2$
 (v) $\cos^{-1}\left(\frac{7x^4 + 5y^4}{7x^4 - 5y^4}\right) = \tan^{-1} a$
 (vi) $\log\left(\frac{x^{20} - y^{20}}{x^{20} + y^{20}}\right) = 20$
 (vii) $e^{\frac{x^7 - y^7}{x^7 + y^7}} = a$
 (viii) $\sin\left(\frac{x^3 - y^3}{x^3 + y^3}\right) = a^3$

(5) (i) If $\log(x+y) = \log(xy) + p$, where p is constant then prove that $\frac{dy}{dx} = -\frac{y^2}{x^2}$.

(ii) If $\log_{10}\left(\frac{x^3 - y^3}{x^3 + y^3}\right) = 2$,
 show that $\frac{dy}{dx} = -\frac{99x^2}{101y^2}$.

(iii) If $\log_5\left(\frac{x^4 + y^4}{x^4 - y^4}\right) = 2$,
 show that $\frac{dy}{dx} = -\frac{12x^3}{13y^3}$.

(iv) If $e^x + e^y = e^{x+y}$, then
 show that $\frac{dy}{dx} = -e^{y-x}$.

(v) If $\sin^{-1}\left(\frac{x^5 - y^5}{x^5 + y^5}\right) = \frac{\pi}{6}$,
 show that $\frac{dy}{dx} = \frac{x^4}{3y^4}$.

(vi) If $x^y = e^{x-y}$, then
 show that $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$.

(vii) If $y = \sqrt{\cos x + \sqrt{\cos x + \sqrt{\cos x + \dots \infty}}}$,
 then show that $\frac{dy}{dx} = \frac{\sin x}{1 - 2y}$.

(viii) If $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots \infty}}}$,
 then show that $\frac{dy}{dx} = \frac{1}{x(2y - 1)}$.

(ix) If $y = x^{x^{\dots \infty}}$, then
 show that $\frac{dy}{dx} = \frac{y^2}{x(1 - \log y)}$.

(x) If $e^y = y^x$, then
 show that $\frac{dy}{dx} = \frac{(\log y)^2}{\log y - 1}$.

1.4.1 Derivatives of Parametric Functions

Consider the equations $x = f(t), y = g(t)$. These equations may imply a functional relation between the variables x and y . Given the value of t in some domain $[a, b]$, we can find x and y .

For example $x = a \cos t$ and $y = a \sin t$. The functional relation between these two functions is that, $x^2 + y^2 = a^2 \cos^2 t + a^2 \sin^2 t = a^2 (\cos^2 t + \sin^2 t) = a^2$ represents the equation of a circle of radius a with center at the origin. And the domain of t is $[0, 2\pi]$. We can find x and y for any $t \in [0, 2\pi]$.

If two variables x and y are defined separately as functions by an inter mediating variable t , then that inter mediating variable is known as parameter. Let us discuss the derivatives of parametric functions.

1.4.2 Theorem : If $x = f(t)$ and $y = g(t)$ are differentiable functions of t so that y is a differentiable function of x and if $\frac{dx}{dt} \neq 0$ then $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$.

Proof : Given that $x = f(t)$ and $y = g(t)$.

Let there be a small increment in the value of t say δt then δx and δy are the corresponding increments in x and y respectively.

As $\delta t, \delta x, \delta y$ are small increments in t, x and y respectively such that $\delta t \neq 0$ and $\delta x \neq 0$.

Consider, the incrementary ratio $\frac{\delta y}{\delta x}$, and note that $\delta x \rightarrow 0 \Rightarrow \delta t \rightarrow 0$.

i.e. $\frac{\delta y}{\delta x} = \frac{\frac{\delta y}{\delta t}}{\frac{\delta x}{\delta t}}$, since $\frac{\delta x}{\delta t} \neq 0$

Taking the limit as $\delta t \rightarrow 0$ on both sides we get,

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \lim_{\delta t \rightarrow 0} \left(\frac{\frac{\delta y}{\delta t}}{\frac{\delta x}{\delta t}} \right)$$

As $\delta t \rightarrow 0, \delta x \rightarrow 0$

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \frac{\lim_{\delta t \rightarrow 0} \left(\frac{\delta y}{\delta t} \right)}{\lim_{\delta t \rightarrow 0} \left(\frac{\delta x}{\delta t} \right)} \dots\dots (I)$$

Since x and y are differentiable function of t . we have,

$$\lim_{\delta t \rightarrow 0} \left(\frac{\delta x}{\delta t} \right) = \frac{dx}{dt} \text{ and } \lim_{\delta t \rightarrow 0} \left(\frac{\delta y}{\delta t} \right) = \frac{dy}{dt} \text{ exist and are finite } \dots\dots (II)$$

From (I) and (II), we get

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} \dots\dots (III)$$

The R.H.S. of (III) exists and is finite, implies L.H.S.of (III) also exist and finite

$$\therefore \lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \frac{dy}{dx}$$

Thus the equation (III) becomes,

$$\boxed{\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}} \text{ where } \frac{dx}{dt} \neq 0$$





SOLVED EXAMPLES

Ex. 1 : Find $\frac{dy}{dx}$ if

(i) $x = at^4, y = 2at^2$

(ii) $x = t - \sqrt{t}, y = t + \sqrt{t}$

(iii) $x = \cos (\log t), y = \log (\cos t)$

(iv) $x = a (\theta + \sin \theta), y = a (1 - \cos \theta)$

(v) $x = \sqrt{1 - t^2}, y = \sin^{-1} t$

Solution :

(i) Given, $y = 2at^2$

Differentiate w. r. t. t

$$\frac{dy}{dt} = 2a \frac{d}{dt} (t^2) = 2a (2t) = 4at. \dots (I)$$

And, $x = at^4$

Differentiate w. r. t. t

$$\frac{dx}{dt} = a \frac{d}{dt} (t^4) = a (4t^3) = 4at^3. \dots (II)$$

Now, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{4at}{4at^3} \dots$ [From (I) and (II)]

$\therefore \frac{dy}{dx} = \frac{1}{t^2}$

(ii) Given, $y = t + \sqrt{t}$

Differentiate w. r. t. t

$$\frac{dy}{dt} = \frac{d}{dt} (t + \sqrt{t}) = 1 + \frac{1}{2\sqrt{t}}$$

$$\frac{dy}{dt} = \frac{2\sqrt{t} + 1}{2\sqrt{t}} \dots (I)$$

And, $x = t - \sqrt{t}$

Differentiate w. r. t. t

$$\frac{dx}{dt} = \frac{d}{dt} (t - \sqrt{t}) = 1 - \frac{1}{2\sqrt{t}}$$

$$\frac{dx}{dt} = \frac{2\sqrt{t} - 1}{2\sqrt{t}} \dots (II)$$

Now, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{2\sqrt{t} + 1}{2\sqrt{t}}}{\frac{2\sqrt{t} - 1}{2\sqrt{t}}} \dots$ [From (I) and (II)]

$\therefore \frac{dy}{dx} = \frac{2\sqrt{t} + 1}{2\sqrt{t} - 1}$

(iii) Given, $y = \log (\cos t)$

Differentiate w. r. t. t

$$\frac{dy}{dt} = \frac{d}{dt} [\log (\cos t)] = \frac{1}{\cot t} \cdot \frac{d}{dt} (\cos t) = \frac{1}{\cot t} (-\sin t) \quad \therefore \frac{dy}{dt} = -\tan t \quad \dots (I)$$

And, $x = \cos (\log t)$

Differentiate w. r. t. t

$$\frac{dx}{dt} = \frac{d}{dt} [\cos (\log t)] = -\sin (\log t) \cdot \frac{d}{dt} (\log t) = -\frac{\sin (\log t)}{t} \quad \therefore \frac{dx}{dt} = -\frac{\sin (\log t)}{t} \quad \dots (II)$$

Now, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-\tan t}{-\frac{\sin (\log t)}{t}} \dots$ [From (I) and (II)]

$\therefore \frac{dy}{dx} = \frac{t \cdot \tan t}{\sin (\log t)}$



(ii) Given, $y = \frac{1}{t^2}$

Differentiate w. r. t. t

$$\frac{dy}{dt} = \frac{d}{dt} \left(\frac{1}{t^2} \right)$$

$$\frac{dy}{dt} = -\frac{2}{t^3} \quad \dots\dots (I)$$

And, $x = t + \frac{1}{t}$

Differentiate w. r. t. t

$$\frac{dx}{dt} = \frac{d}{dt} \left(t + \frac{1}{t} \right) = 1 - \frac{1}{t^2}$$

$$\frac{dx}{dt} = -\frac{t^2 - 1}{t^2} \quad \dots\dots (II)$$

Now, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-\frac{2}{t^3}}{\frac{t^2-1}{t^2}} \dots$ [From (I) and (II)]

$$\therefore \frac{dy}{dx} = -\frac{2}{t(t^2 - 1)}$$

At $t = \frac{1}{2}$, we get

$$\left(\frac{dy}{dx} \right)_{t=\frac{1}{2}} = -\frac{2}{\left(\frac{1}{2}\right) \left[\left(\frac{1}{2}\right)^2 - 1 \right]}$$

$$= -\frac{2}{\left(\frac{1}{2}\right) \left(\frac{1}{4} - 1\right)}$$

$$\left(\frac{dy}{dx} \right)_{t=\frac{1}{2}} = -\frac{2}{\left(\frac{1}{2}\right) \left(-\frac{3}{4}\right)}$$

$$\left(\frac{dy}{dx} \right)_{t=\frac{1}{2}} = \frac{16}{3}$$

(iii) Given, $y = 3 \sin t - 2 \sin^3 t$

Differentiate w. r. t. t

$$\frac{dy}{dt} = \frac{d}{dt} (3 \sin t - 2 \sin^3 t)$$

$$= 3 \frac{d}{dt} (\sin t) - 2 (\sin t)^3$$

$$= 3 \cos t - 2(3) \sin^2 t \frac{d}{dt} (\sin t)$$

$$= 3 \cos t - 6 \sin^2 t (\cos t)$$

$$= 3 \cos t (1 - 2 \sin^2 t)$$

$$\frac{dy}{dt} = 3 \cos t \cos 2t \quad \dots\dots (I)$$

And, $x = 3 \cos t - 2 \cos^3 t$

Differentiate w. r. t. t

$$\frac{dx}{dt} = \frac{d}{dt} (3 \cos t - 2 \cos^3 t)$$

$$= 3 \frac{d}{dt} (\cos t) - 2 \frac{d}{dt} (\cos^3 t)$$

$$= 3(-\sin t) - 2(3) \cos^2 t \frac{d}{dt} (\cos t)$$

$$= -3 \sin t - 6 \cos^2 t (-\sin t)$$

$$= 6 \cos^2 t \sin t - 3 \sin t$$

$$= 3 \sin t (2 \cos^2 t - 1)$$

$$\frac{dx}{dt} = 3 \sin t \cos 2t \quad \dots\dots (II)$$

Now, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{3 \cos t \cos 2t}{3 \sin t \cos 2t} \dots\dots$ [From (I) and (II)]

$$\therefore \frac{dy}{dx} = -\cot t$$

At $t = \frac{\pi}{6}$, we get

$$\left(\frac{dy}{dx} \right)_{t=\frac{\pi}{6}} = -\cot \left(\frac{\pi}{6} \right) = -\sqrt{3}$$

Ex. 3 : If $x^2 + y^2 = t + \frac{1}{t}$ and $x^4 + y^4 = t^2 + \frac{1}{t^2}$, then show that $x^3y \frac{dy}{dx} = 1$.

Solution :

Given that, $x^4 + y^4 = t^2 + \frac{1}{t^2}$... (I)

And $x^2 + y^2 = t + \frac{1}{t}$

Squaring both sides,

$$(x^2 + y^2)^2 = \left(t + \frac{1}{t}\right)^2$$

$$x^4 + 2x^2y^2 + y^4 = t^2 + 2 + \frac{1}{t^2}$$

$$x^4 + 2x^2y^2 + y^4 = x^4 + y^4 + 2 \quad \dots \text{ [From (I)]}$$

$$2x^2y^2 = 2 \quad \therefore x^2y^2 = 1 \quad \dots \text{ (II)}$$

Differentiate w. r. t. x

$$\frac{d}{dx}(x^2y^2) = \frac{d}{dx}(1)$$

$$x^2 \frac{d}{dx}(y^2) + y^2 \frac{d}{dx}(x^2) = 0$$

$$x^2(2y) \frac{dy}{dx} + y^2(2x) = 0$$

$$2x^2y \frac{dy}{dx} = -2xy^2 \Rightarrow \frac{dy}{dx} = -\frac{2xy^2}{2x^2y}$$

$$\frac{dy}{dx} = -\frac{x\left(-\frac{1}{x^2}\right)}{x^2y} \quad \dots \text{ [From (II)]}$$

$$\therefore \frac{dy}{dx} = \frac{1}{x^3y} \quad \therefore x^3y \frac{dy}{dx} = 1$$

Ex. 4 : If $x = a\left(t - \frac{1}{t}\right)$ and $y = b\left(t + \frac{1}{t}\right)$, then show that $\frac{dy}{dx} = \frac{b^2x}{a^2y}$.

Solution :

Given that, $x = a\left(t - \frac{1}{t}\right)$ and $y = b\left(t + \frac{1}{t}\right)$

i.e. $\frac{x}{a} = t - \frac{1}{t}$... (I) and $\frac{y}{b} = t + \frac{1}{t}$... (II)

Square of (I) - Square of (II) gives,

$$\begin{aligned} \frac{x^2}{a^2} - \frac{y^2}{b^2} &= \left(t - \frac{1}{t}\right)^2 - \left(t + \frac{1}{t}\right)^2 \\ &= t^2 - 2 + \frac{1}{t^2} - t^2 - 2 - \frac{1}{t^2} \end{aligned}$$

$$\therefore \frac{x^2}{a^2} - \frac{y^2}{b^2} = -4$$

Differentiate w. r. t. x

$$\frac{1}{a^2} \cdot \frac{d}{dx}(x^2) - \frac{1}{b^2} \cdot \frac{d}{dx}(y^2) = \frac{d}{dx}(-4)$$

$$\frac{1}{a^2}(2x) - \frac{1}{b^2}(2y) \cdot \frac{dy}{dx} = 0$$

$$\frac{1}{a^2}(2x) - \frac{1}{b^2}(2y) \cdot \frac{dy}{dx} = 0$$

$$\frac{2y}{b^2} \cdot \frac{dy}{dx} = \frac{2x}{a^2} \Rightarrow \frac{dy}{dx} = \frac{b^2x}{a^2y}$$

$$\therefore \frac{dy}{dx} = \frac{b^2x}{a^2y}$$

Ex. 5 : If $x = \sqrt{a^{\sin^{-1}t}}$ and $y = \sqrt{a^{\cos^{-1}t}}$, then show that $\frac{dy}{dx} = -\frac{y}{x}$.

Solution : Given that, $x = \sqrt{a^{\sin^{-1}t}}$ and $y = \sqrt{a^{\cos^{-1}t}}$

i.e. $x = \sqrt{a^{\sin^{-1}t}}$... (I) and $y = \sqrt{a^{\cos^{-1}t}}$... (II)

Differentiate (I) w. r. t. t

$$\frac{dx}{dt} = \frac{d}{dt}\left(\sqrt{a^{\sin^{-1}t}}\right) = \frac{1}{2\sqrt{a^{\sin^{-1}t}}} \cdot \frac{d}{dt}(a^{\sin^{-1}t})$$

$$= \frac{1}{2\sqrt{a^{\sin^{-1}t}}} \cdot a^{\sin^{-1}t} \cdot \log a \frac{d}{dt}(\sin^{-1}t)$$

$$= \frac{a^{\sin^{-1} t} \cdot \log a}{2\sqrt{a^{\sin^{-1} t}} \cdot \sqrt{1-x^2}} \cdot \frac{1}{\sqrt{1-x^2}}$$

$$\frac{dx}{dt} = \frac{\sqrt{a^{\sin^{-1} t}} \cdot \log a}{2\sqrt{1-x^2}} = \frac{x \log a}{2\sqrt{1-x^2}} \dots \text{(III)} \dots \text{[From (I)]}$$

Now $y = \sqrt{a^{\cos^{-1} t}}$

Differentiate (II) w. r. t. t

$$\frac{dy}{dt} = \frac{d}{dt} \left(\sqrt{a^{\cos^{-1} t}} \right) = \frac{1}{2\sqrt{a^{\cos^{-1} t}}} \cdot \frac{d}{dt} (a^{\cos^{-1} t})$$

$$= \frac{1}{2\sqrt{a^{\cos^{-1} t}}} \cdot a^{\cos^{-1} t} \cdot \log a \cdot \frac{d}{dt} (\cos^{-1} t)$$

$$= \frac{a^{\cos^{-1} t} \cdot \log a}{2\sqrt{a^{\cos^{-1} t}}} \left(-\frac{1}{\sqrt{1-x^2}} \right)$$

$$\frac{dy}{dt} = \frac{-\sqrt{a^{\cos^{-1} t}} \cdot \log a}{2\sqrt{1-x^2}} = -\frac{y \log a}{2\sqrt{1-x^2}} \dots \text{(IV)} \dots \text{[From (II)]}$$

Now, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-\frac{y \log a}{2\sqrt{1-x^2}}}{\frac{x \log a}{2\sqrt{1-x^2}}} \dots \dots \text{[From (III) and (IV)]}$

$\therefore \frac{dy}{dx} = -\frac{y}{x}$

1.4.3 Differentiation of one function with respect to another function :

If y is differentiable function of x , then the derivative of y with respect to x is $\frac{dy}{dx}$.

Similarly, if $u = f(x)$, $v = g(x)$ differentiable function of x , such that $\frac{du}{dx} = f'(x)$ and $\frac{dv}{dx} = g'(x)$

then the derivative of u with respect to v is $\frac{du}{dv} = \frac{\frac{du}{dx}}{\frac{dv}{dx}} = \frac{f'(x)}{g'(x)}$.



SOLVED EXAMPLES

Ex. 1 : Find the derivative of 7^x w. r. t. x^7 .

Solution : Let : $u = 7^x$ and $v = x^7$, then we have to find $\frac{du}{dv}$.

$$\therefore \frac{du}{dv} = \frac{\frac{du}{dx}}{\frac{dv}{dx}} \dots \text{(I)}$$

Now, $u = 7^x$

Differentiate w. r. t. x

$$\frac{du}{dx} = \frac{d}{dx} (7^x) = 7^x \log 7 \dots \text{(II)}$$

And, $v = x^7$

Differentiate w. r. t. x

$$\frac{dv}{dx} = \frac{d}{dx} (x^7) = 7x^6 \dots \text{(III)}$$

Substituting (II) and (III) in (I) we get,

$$\therefore \frac{du}{dv} = \frac{7^x \log 7}{7x^6}$$

Ex. 2 : Find the derivative of $\cos^{-1} x$ w. r. t. $\sqrt{1-x^2}$.

Solution : Let $u = \cos^{-1} x$ and $v = \sqrt{1-x^2}$, then we have to find $\frac{du}{dv}$.

$$\text{i.e. } \frac{du}{dv} = \frac{\frac{du}{dx}}{\frac{dv}{dx}} \quad \dots \text{ (I)}$$

Now, $u = \cos^{-1} x$

Differentiate w. r. t. x

$$\frac{du}{dx} = \frac{d}{dx} (\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}} \quad \dots \text{ (II)}$$

And, $v = \sqrt{1-x^2}$

Differentiate w. r. t. x

$$\frac{dv}{dx} = \frac{d}{dx} (\sqrt{1-x^2}) = \frac{1}{2\sqrt{1-x^2}} \cdot \frac{d}{dx} (1-x^2) = \frac{1}{2\sqrt{1-x^2}} \cdot (-2x)$$

$$\frac{dv}{dx} = -\frac{x}{\sqrt{1-x^2}} \quad \dots \text{ (III)}$$

Substituting (II) and (III) in (I) we get,

$$\frac{du}{dv} = \frac{-\frac{1}{\sqrt{1-x^2}}}{-\frac{x}{\sqrt{1-x^2}}} \quad \therefore \quad \frac{du}{dv} = \frac{1}{x}$$

Ex. 3 : Find the derivative of $\tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$ w. r. t. $\sin^{-1} \left(\frac{2x}{1+x^2} \right)$.

Solution : Let $u = \tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$ and $v = \sin^{-1} \left(\frac{2x}{1+x^2} \right)$, then we have to find $\frac{du}{dv}$.

$$\text{i.e. } \frac{du}{dv} = \frac{\frac{du}{dx}}{\frac{dv}{dx}} \quad \dots \text{ (I)}$$

$$\text{Now, } u = \tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$$

Put $x = \tan \theta \therefore \theta = \tan^{-1} x$

$$u = \tan^{-1} \left(\frac{\sqrt{1+\tan^2 \theta}-1}{\tan \theta} \right) = \tan^{-1} \left(\frac{\sec \theta - 1}{\tan \theta} \right) = \tan^{-1} \left(\frac{\frac{1}{\cos \theta} - 1}{\frac{\sin \theta}{\cos \theta}} \right) = \tan^{-1} \left(\frac{1 - \cos \theta}{\sin \theta} \right)$$

$$= \tan^{-1} \left[\frac{2 \sin^2 \left(\frac{\theta}{2} \right)}{2 \sin \left(\frac{\theta}{2} \right) \cos \left(\frac{\theta}{2} \right)} \right] = \tan^{-1} \left[\tan \left(\frac{\theta}{2} \right) \right]$$

$$u = \frac{\theta}{2} = \frac{1}{2} \tan^{-1} x$$

Differentiate w. r. t. x

$$\frac{du}{dx} = \frac{1}{2} \frac{d}{dx} (\tan^{-1} x) = \frac{1}{2(1+x^2)} \quad \dots \text{ (II)}$$

And, $v = \sin^{-1}\left(\frac{2x}{1+x^2}\right) = \sin^{-1}\left(\frac{2 \tan \theta}{1 + \tan^2 \theta}\right) = \sin^{-1}(\sin 2\theta) = 2\theta$

$v = 2 \tan^{-1} x$

Differentiate w. r. t. x

$\frac{dv}{dx} = 2 \frac{d}{dx}(\tan^{-1} x) = \frac{2}{1+x^2} \dots (III)$

Substituting (II) and (III) in (I) we get,

$\frac{du}{dv} = \frac{\frac{1}{2(1+x^2)}}{\frac{2}{1+x^2}} = \frac{1}{4}$

EXERCISE 1.4

(1) Find $\frac{dy}{dx}$ if

- (i) $x = at^2, y = 2at$
- (ii) $x = a \cot \theta, y = b \operatorname{cosec} \theta$
- (iii) $x = \sqrt{a^2 + m^2}, y = \log(a^2 + m^2)$
- (iv) $x = \sin \theta, y = \tan \theta$
- (v) $x = a(1 - \cos \theta), y = b(\theta - \sin \theta)$
- (vi) $x = \left(t + \frac{1}{t}\right)^a, y = a^{t + \frac{1}{t}}$,

where $a > 0, a \neq 1$ and $t \neq 0$.

- (vii) $x = \cos^{-1}\left(\frac{2t}{1+t^2}\right), y = \sec^{-1}(\sqrt{1+t^2})$
- (viii) $x = \cos^{-1}(4t^3 - 3t), y = \tan^{-1}\left(\frac{\sqrt{1-t^2}}{t}\right)$

(2) Find $\frac{dy}{dx}$ if

- (i) $x = \operatorname{cosec}^2 \theta, y = \cot^3 \theta$, at $\theta = \frac{\pi}{6}$
- (ii) $x = a \cos^3 \theta, y = a \sin^3 \theta$, at $\theta = \frac{\pi}{3}$
- (iii) $x = t^2 + t + 1, y = \sin\left(\frac{\pi t}{2}\right) + \cos\left(\frac{\pi t}{2}\right)$,
at $t = 1$
- (iv) $x = 2 \cos t + \cos 2t, y = 2 \sin t - \sin 2t$,
at $t = \frac{\pi}{4}$
- (v) $x = t + 2 \sin(\pi t), y = 3t - \cos(\pi t)$,
at $t = \frac{1}{2}$

(3) (i) If $x = a\sqrt{\sec \theta - \tan \theta}, y = a\sqrt{\sec \theta + \tan \theta}$,

then show that $\frac{dy}{dx} = -\frac{y}{x}$.

(ii) If $x = e^{\sin 3t}, y = e^{\cos 3t}$, then

show that $\frac{dy}{dx} = -\frac{y \log x}{x \log y}$.

(iii) If $x = \frac{t+1}{t-1}, y = \frac{t-1}{t+1}$, then

show that $y^2 + \frac{dy}{dx} = 0$.

(iv) If $x = a \cos^3 t, y = a \sin^3 t$, then

show that $\frac{dy}{dx} = -\left(\frac{y}{x}\right)^{\frac{1}{3}}$.

(v) If $x = 2 \cos^4(t+3), y = 3 \sin^4(t+3)$,

show that $\frac{dy}{dx} = -\sqrt{\frac{3y}{2x}}$.

(vi) If $x = \log(1+t^2), y = t - \tan^{-1} t$,

show that $\frac{dy}{dx} = \frac{\sqrt{e^x - 1}}{2}$.

(vii) If $x = \sin^{-1}(e^t), y = \sqrt{1 - e^{2t}}$,

show that $\sin x + \frac{dy}{dx} = 0$.

(viii) If $x = \frac{2bt}{1+t^2}, y = a\left(\frac{1-t^2}{1+t^2}\right)$,

show that $\frac{dx}{dy} = -\frac{b^2 y}{a^2 x}$.

(4) (i) Differentiate $x \sin x$ w. r. t. $\tan x$.

(ii) Differentiate $\sin^{-1} \left(\frac{2x}{1+x^2} \right)$

w. r. t. $\cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)$.

(iii) Differentiate $\tan^{-1} \left(\frac{x}{\sqrt{1-x^2}} \right)$

w. r. t. $\sec^{-1} \left(\frac{1}{2x^2-1} \right)$.

(iv) Differentiate $\cos^{-1} \left(\frac{1-x^2}{1+x^2} \right)$ w. r. t. $\tan^{-1} x$.

(v) Differentiate 3^x w. r. t. $\log_x 3$.

(vi) Differentiate $\tan^{-1} \left(\frac{\cos x}{1+\sin x} \right)$

w. r. t. $\sec^{-1} x$.

(vii) Differentiate x^x w. r. t. $x^{\sin x}$.

(viii) Differentiate $\tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$

w. r. t. $\tan^{-1} \left(\frac{2x\sqrt{1-x^2}}{1-2x^2} \right)$.

1.5.1 Higher order derivatives :

If $f(x)$ is differentiable function of x on an open interval I , then its derivative $f'(x)$ is also a function on I , so $f'(x)$ may have a derivative of its own, denoted as $(f'(x))' = f''(x)$. This new function $f''(x)$ is called the **second derivative** of $f(x)$. By Leibniz notation, we write the second derivative of

$$y = f(x) \text{ as } y'' = f''(x) = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d^2y}{dx^2}$$

By method of first principle

$$f'(x) = \lim_{h \rightarrow 0} \left(\frac{f(x+h) - f(x)}{h} \right) = \frac{dy}{dx} \text{ and}$$

$$f''(x) = \lim_{h \rightarrow 0} \left(\frac{f'(x+h) - f'(x)}{h} \right) = \frac{d^2y}{dx^2}$$

Further if $f''(x)$ is a differentiable function of x then its derivative is denoted as $\frac{d}{dx}[f''(x)] = f'''(x)$.

Now the new function $f'''(x)$ is called the **third derivative** of $f(x)$. We write the third of $y = f(x)$ as

$$y''' = f'''(x) = \frac{d}{dx} \left(\frac{d^2y}{dx^2} \right) = \frac{d^3y}{dx^3}. \text{ The fourth derivative, is usually denoted by } f^{(4)}(x). \text{ Therefore}$$

$$f^{(4)}(x) = \frac{d^4y}{dx^4}.$$

In general, the n^{th} derivative of $f(x)$, is denoted by $f^{(n)}(x)$ and it obtained by differentiating $f(x)$, n times. So, we can write the n^{th} derivative of $y = f(x)$ as $y^{(n)} = f^{(n)}(x) = \frac{d^n y}{dx^n}$. These are called higher order derivatives.

Note : The higher order derivatives may also be denoted by y_2, y_3, \dots, y_n .

For example: Consider $f(x) = x^3 - x$

Differentiate w. r. t. x

$$f'(x) = \frac{d}{dx} [f(x)] = 3x^2 - 1$$

Differentiate w. r. t. x


$$f''(x) = \frac{d}{dx} [f'(x)] = 6x$$

Differentiate w. r. t. x

$$f'''(x) = \frac{d}{dx} [f''(x)] = 6$$

Which is the slope of the line represented by $f''(x)$. Hence forth all its next derivatives are zero.

Note : From the above example we can deduce one important result that, if $f(x)$ is a polynomial of degree n , then its n^{th} order derivative is a constant and all the onward derivatives are zeros.



SOLVED EXAMPLES

Ex. 1 : Find the second order derivative of the following :

(i) $x^3 + 7x^2 - 2x - 9$

(ii) $x^2 e^x$

(iii) $e^{2x} \sin 3x$

(iv) $x^2 \log x$

(v) $\sin (\log x)$

Solution :

(i) Let $y = x^3 + 7x^2 - 2x - 9$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (x^3 + 7x^2 - 2x - 9)$$

$$\frac{dy}{dx} = 3x^2 + 14x - 2$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} (3x^2 + 14x - 2)$$

$$\frac{d^2y}{dx^2} = 6x + 14$$

(ii) Let $y = x^2 e^x$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (x^2 e^x)$$

$$\frac{dy}{dx} = x^2 \frac{d}{dx} (e^x) + e^x \frac{d}{dx} (x^2)$$

$$\frac{dy}{dx} = x^2 e^x + 2x e^x = e^x (x^2 + 2x)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} [e^x (x^2 + 2x)]$$

$$\frac{d^2y}{dx^2} = e^x \frac{d}{dx} (x^2 + 2x) + (x^2 + 2x) \frac{d}{dx} (e^x)$$

$$= e^x (2x + 2) + (x^2 + 2x) (e^x)$$

$$= (x^2 + 4x + 2) e^x$$

$$\frac{d^2y}{dx^2} = (x^2 + 4x + 2) e^x$$

(iii) Let $y = e^{2x} \sin 3x$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (e^{2x} \sin 3x) = e^{2x} \frac{d}{dx} (\sin 3x) + \sin 3x \frac{d}{dx} (e^{2x})$$

$$\frac{dy}{dx} = e^{2x} (\cos 3x) (3) + \sin 3x (e^{2x}) (2)$$

$$\frac{dy}{dx} = e^{2x} (3 \cos 3x + 2 \sin 3x)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} [e^{2x} (3 \cos 3x + 2 \sin 3x)]$$

$$\frac{d^2y}{dx^2} = e^{2x} \frac{d}{dx} (3 \cos 3x + 2 \sin 3x) + (3 \cos 3x + 2 \sin 3x) \frac{d}{dx} (e^{2x})$$

$$= e^{2x} [3 (-\sin 3x) (3) + 2 (\cos 3x)(3)] + (3 \cos 3x + 2 \sin 3x) e^{2x} (2)$$

$$= e^{2x} [-9 \sin 3x + 6 \cos 3x + 6 \cos 3x + 4 \sin 3x]$$

$$\frac{d^2y}{dx^2} = e^{2x} [12 \cos 3x - 5 \sin 3x]$$

(iv) Let $y = x^2 \log x$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (x^2 \log x)$$

$$\frac{dy}{dx} = x^2 \frac{d}{dx} (\log x) + \log x \frac{d}{dx} (x^2)$$

$$\frac{dy}{dx} = x^2 \cdot \frac{1}{x} + \log x (2x)$$

$$\frac{dy}{dx} = x(1 + 2 \log x)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} [x(1 + 2 \log x)]$$

$$\frac{d^2y}{dx^2} = x \frac{d}{dx} (1 + 2 \log x) + (1 + 2 \log x) \frac{d}{dx} (x)$$

$$= x \cdot \frac{2}{x} + (1 + 2 \log x) (1)$$

$$\frac{d^2y}{dx^2} = 3 + 2 \log x$$

(v) Let $y = \sin (\log x)$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} [\sin (\log x)]$$

$$\frac{dy}{dx} = \cos (\log x) \frac{d}{dx} (\log x)$$

$$\frac{dy}{dx} = \frac{\cos (\log x)}{x}$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} \left[\frac{\cos (\log x)}{x} \right]$$

$$\frac{d^2y}{dx^2} = \frac{x \frac{d}{dx} [\cos (\log x)] - \cos (\log x) \frac{d}{dx} (x)}{x^2}$$

$$= \frac{x [-\sin (\log x)] \frac{d}{dx} (\log x) - \cos (\log x)(1)}{x^2}$$

$$= \frac{-\frac{x \sin (\log x)}{x} - \cos (\log x)}{x^2}$$

$$\frac{d^2y}{dx^2} = -\frac{\sin (\log x) + \cos (\log x)}{x^2}$$

Ex. 2 : Find $\frac{d^2y}{dx^2}$ if, (i) $x = \cot^{-1} \left(\frac{\sqrt{1-t^2}}{t} \right)$ and $x = \operatorname{cosec}^{-1} \left(\frac{1+t^2}{2t} \right)$ (ii) $x = a \cos^3 \theta, y = b \sin^3 \theta$ at $\theta = \frac{\pi}{4}$

Solution :

(i) $x = \cot^{-1} \left(\frac{\sqrt{1-t^2}}{t} \right)$ and $x = \operatorname{cosec}^{-1} \left(\frac{1+t^2}{2t} \right)$

Put $t = \sin \theta \quad \therefore \theta = \sin^{-1} t$

$$x = \cot^{-1} \left(\frac{\sqrt{1-\sin^2 \theta}}{\sin \theta} \right) = \cot^{-1} \left(\frac{\sqrt{\sin^2 \theta}}{\sin \theta} \right)$$

$$x = \cot^{-1} (\cot \theta) = \theta \quad \therefore x = \sin^{-1} t$$

Differentiate w. r. t. t

$$\frac{dx}{dt} = \frac{d}{dt} (\sin^{-1} t) = \left(\frac{1}{\sqrt{1-t^2}} \right) \dots \text{(I)}$$

We know that,

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{2}{1+t^2} \dots [\text{From (I) and (II)}] \quad \therefore \frac{dy}{dx} = \left(\frac{2\sqrt{1-t^2}}{1+t^2} \right)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \cdot \frac{dy}{dx} = \frac{d}{dx} \cdot \left(\frac{2\sqrt{1-t^2}}{1+t^2} \right)$$

$$\frac{d^2y}{dx^2} = 2 \frac{d}{dt} \cdot \left(\frac{\sqrt{1-t^2}}{1+t^2} \right) \times \frac{dt}{dx}$$

$$= 2 \times \left[\frac{(1+t^2) \frac{d}{dt} (\sqrt{1-t^2}) - \sqrt{1-t^2} \frac{d}{dt} (1+t^2)}{(1+t^2)^2} \right] \times \frac{1}{\frac{dx}{dt}}$$

$$= 2 \times \left[\frac{(1+t^2) \frac{1}{2\sqrt{1-t^2}} \frac{d}{dt} (\sqrt{1-t^2}) - \sqrt{1-t^2} (2t)}{(1+t^2)^2} \right] \times \frac{1}{\frac{1}{\sqrt{1-t^2}}} [\text{From (I)}]$$

$$= 2 \times \left[\frac{(1+t^2) \frac{1}{2\sqrt{1-t^2}} (-2t) - 2t (\sqrt{1-t^2})}{(1+t^2)^2} \right] \times \sqrt{1-t^2}$$

$$= 2 \times \left[\frac{(1+t^2) \frac{-t}{\sqrt{1-t^2}} - 2t (\sqrt{1-t^2})}{(1+t^2)^2} \right] \times \sqrt{1-t^2}$$

$$= 2 \times \left[\frac{-t(1+t^2) - 2t(1-t^2)}{(1+t^2)^2} \right] = 2 \times \left[\frac{-t-t^3-2t+2t^3}{(1+t^2)^2} \right]$$

$$= 2 \times \left[\frac{t^3-3t}{(1+t^2)^2} \right]$$

$$\frac{d^2y}{dx^2} = \frac{2t(t^2-3)}{(1+t^2)^2}$$

$$y = \operatorname{cosec}^{-1} \left(\frac{1+t^2}{2t} \right) = \sin^{-1} \left(\frac{2t}{1+t^2} \right)$$

Put $t = \tan \theta \quad \therefore \theta = \tan^{-1} t$

$$y = \sin^{-1} \left(\frac{2 \tan \theta}{1 + \tan^2 \theta} \right) = \sin^{-1} (\sin 2\theta) = 2\theta$$

$$\therefore y = 2 \tan^{-1} t$$

Differentiate w. r. t. t

$$\frac{dy}{dt} = 2 \frac{d}{dt} (\tan^{-1} t) = \frac{2}{1+t^2} \dots \text{(II)}$$

(ii) $x = a \cos^3 \theta, y = b \sin^3 \theta$ at $\theta = \frac{\pi}{4}$

Solution :

Given that : $x = a \cos^3 \theta$

Differentiate w. r. t. θ

$$\frac{dx}{d\theta} = \frac{d}{d\theta} (a \cos^3 \theta) = a (3) (\cos^2 \theta) \frac{d}{d\theta} (\cos \theta)$$

$$\frac{dx}{d\theta} = -3a \cos^2 \theta \sin \theta \quad \dots \text{ (I)}$$

$y = b \sin^3 \theta$

Differentiate w. r. t. θ

$$\frac{dy}{d\theta} = \frac{d}{d\theta} (b \sin^3 \theta) = b (3) (\sin^2 \theta) \frac{d}{d\theta} (\sin \theta)$$

$$\frac{dy}{d\theta} = 3b \sin^2 \theta \cos \theta \quad \dots \text{ (II)}$$

We know that,

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{3b \sin^2 \theta \cos \theta}{-3a \cos^2 \theta \sin \theta} \quad \dots \text{ [From (I) and (II)]}$$

$$\therefore \frac{dy}{dx} = -\frac{b}{a} \cdot \tan \theta$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = -\frac{b}{a} \cdot \frac{d}{dx} (\tan \theta)$$

$$\frac{d^2 y}{dx^2} = -\frac{b}{a} \cdot \frac{d}{d\theta} (\tan \theta) \times \frac{d\theta}{dx}$$

$$= -\frac{b}{a} (\sec^2 \theta) \times \frac{1}{\frac{dx}{d\theta}}$$

$$= -\frac{b}{a} (\sec^2 \theta) \times \frac{1}{-3a \cos^2 \theta \sin \theta} \quad \dots \text{ [From (I)]}$$

$$\frac{d^2 y}{dx^2} = \frac{b}{3a^2} \times \frac{\sec^2 \theta}{\cos^2 \theta \sin \theta}$$

$$\frac{d^2 y}{dx^2} = \frac{b \sec^4 \theta}{3a^2 \sin \theta}$$

When $\theta = \frac{\pi}{4}$

$$\left(\frac{d^2 y}{dx^2} \right)_{\theta = \frac{\pi}{4}} = \frac{b \sec^4 \left(\frac{\pi}{4} \right)}{3a^2 \sin \left(\frac{\pi}{4} \right)} = \frac{b (\sqrt{2})^4}{3a^2 \left(\frac{1}{\sqrt{2}} \right)}$$

$$\left(\frac{d^2 y}{dx^2} \right)_{\theta = \frac{\pi}{4}} = \frac{4 \sqrt{2} b}{3a^2}$$

Ex. 3 : If $ax^2 + 2hxy + by^2 = 0$ then show that $\frac{d^2y}{dx^2} = 0$.

Solution : Given that $ax^2 + 2hxy + by^2 = 0 \dots$ (I)

$$ax^2 + hxy + hxy + by^2 = 0$$

$$x(ax + hy) + y(hx + by) = 0$$

$$y(hx + by) = -x(ax + hy)$$

$$\frac{y}{x} = -\frac{ax + hy}{hx + by} \dots$$
 (II)

Differentiate (I) w. r. t. x

$$a \frac{d}{dx} (x^2) + 2h \frac{d}{dx} (xy) + b \frac{d}{dx} (y^2) = 0$$

$$a (2x) + 2h \left[a \frac{dy}{dx} + y(1) \right] + b (2y) \frac{dy}{dx} = 0$$

$$2 \left[ax + hx \frac{dy}{dx} + hy + by \frac{dy}{dx} \right] = 0$$

$$(hx + by) \frac{dy}{dx} = -ax - hy$$

$$\frac{dy}{dx} = -\frac{ax + hy}{hx + by}$$

From (II), we get

$$\therefore \frac{dy}{dx} = \frac{y}{x} \dots$$
 (III)

Differentiate (III), w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} \left(\frac{y}{x} \right)$$

$$\frac{d^2y}{dx^2} = \frac{x \frac{dy}{dx} - y(1)}{x^2} = \frac{x \left(\frac{y}{x} \right) - y}{x^2} \dots$$
 [From (II)]

$$\therefore \frac{d^2y}{dx^2} = \frac{y - y}{x^2} = 0$$

Ex. 4 : If $y = \cos (m \cos^{-1} x)$ then show that $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + m^2y = 0$.

Solution : Given that $y = \cos (m \cos^{-1} x)$

$$\therefore \cos^{-1} y = m \cos^{-1} x$$

Differentiate (I) w. r. t. x

$$\frac{d}{dx} (\cos^{-1} y) = m \frac{d}{dx} (\cos^{-1} x)$$

$$-\frac{1}{\sqrt{1-y^2}} \cdot \frac{dy}{dx} = -\frac{m}{\sqrt{1-x^2}}$$

$$\sqrt{1-x^2} \cdot \frac{dy}{dx} = m \sqrt{1-y^2}$$

Squaring both sides

$$(1-x^2) \cdot \left(\frac{dy}{dx}\right)^2 = m^2 (1-y^2)$$

Differentiate w. r. t. x

$$(1-x^2) \frac{d}{dx} \left(\frac{dy}{dx}\right)^2 + \left(\frac{dy}{dx}\right)^2 \frac{d}{dx} (1-x^2) = m^2 \frac{d}{dx} (1-y^2)$$

$$(1-x^2) \cdot 2 \left(\frac{dy}{dx}\right) \cdot \frac{d}{dx} \left(\frac{dy}{dx}\right) + \left(\frac{dy}{dx}\right)^2 (-2x) = m^2 (-2y) \frac{dy}{dx}$$

$$2(1-x^2) \cdot \frac{dy}{dx} \cdot \frac{d^2y}{dx^2} - 2x \left(\frac{dy}{dx}\right)^2 = -2m^2y \frac{dy}{dx}$$

Dividing throughout by $2 \frac{dy}{dx}$ we get,

$$(1-x^2) \cdot \frac{d^2y}{dx^2} - x \frac{dy}{dx} = -m^2y$$

$$\therefore (1-x^2) \cdot \frac{d^2y}{dx^2} - x \frac{dy}{dx} + m^2y = 0$$

Ex. 5 : If $x = \sin t$, $y = e^{mt}$ then show that $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - m^2y = 0$.

Solution : Given that $x = \sin t \quad \therefore \quad t = \sin^{-1} x$
and $y = e^{mt} \quad \therefore \quad y = e^{m \sin^{-1} x} \quad \dots (I)$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (e^{m \sin^{-1} x}) = e^{m \sin^{-1} x} \cdot m \frac{d}{dx} (\sin^{-1} x)$$

$$\frac{dy}{dx} = \frac{m \cdot e^{m \sin^{-1} x}}{\sqrt{1-x^2}}$$

$$\sqrt{1-x^2} \frac{dy}{dx} = my \quad \dots [\text{From (I)}]$$

Squaring both sides

$$(1-x^2) \cdot \left(\frac{dy}{dx}\right)^2 = m^2 y^2$$

Differentiate w. r. t. x

$$(1-x^2) \frac{d}{dx} \left(\frac{dy}{dx}\right)^2 + \left(\frac{dy}{dx}\right)^2 \frac{d}{dx} (1-x^2) = m^2 \frac{d}{dx} (y^2)$$

$$(1-x^2) \cdot 2 \left(\frac{dy}{dx}\right) \cdot \frac{d}{dx} \left(\frac{dy}{dx}\right) + \left(\frac{dy}{dx}\right)^2 (-2x) = m^2 (2y) \frac{dy}{dx}$$

$$2(1-x^2) \cdot \frac{dy}{dx} \cdot \frac{d^2y}{dx^2} - 2x \left(\frac{dy}{dx} \right)^2 = 2m^2y \frac{dy}{dx}$$

Dividing throughout by $2 \frac{dy}{dx}$ we get,

$$(1-x^2) \cdot \frac{d^2y}{dx^2} - x \frac{dy}{dx} = m^2y$$

$$\therefore (1-x^2) \cdot \frac{d^2y}{dx^2} - x \frac{dy}{dx} - m^2y = 0$$

1.5.2 Successive differentiation (or n^{th} order derivative) of some standard functions :

Successive Differentiation is the process of differentiating a given function successively for n times and the results of such differentiation are called successive derivatives. The higher order derivatives are of utmost importance in scientific and engineering applications.

There is no general formula to find n^{th} derivative of a function. Because each and every function has it's own specific general formula for it's n^{th} derivative. But there are algorithms to find it.

So, here is the algorithm, for some standard functions.

Let us use the method of mathematical induction wherever applicable.

Step 1 :- Use simple differentiation to get 1st, 2nd and 3rd order derivatives.

Step 2 :- Observe the changes in the coefficients, the angles, the power of the function and the signs of each term etc.

Step 3 :- Express the n^{th} derivative with the help of the patterns of changes that you have observed. This will be your general formula for the n^{th} derivative of the given standard function.



SOLVED EXAMPLES

Ex. 1 : Find the n^{th} derivative of the following :

(i) x^m

(ii) $\frac{1}{ax+b}$

(iii) $\log x$

(iv) $\sin x$

(v) $\cos(ax+b)$

(vi) $e^{ax} \sin(bx+c)$

Solution :

(i) Let $y = x^m$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (x^m) = mx^{m-1}$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = m \frac{d}{dx} x^{m-1}$$

$$\frac{d^2y}{dx^2} = m \cdot (m-1) x^{m-2}$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{d^2y}{dx^2} \right) = m \cdot (m-1) \frac{d}{dx} (x^{m-2})$$

$$\frac{d^3y}{dx^3} = m \cdot (m-1) \cdot (m-2) x^{m-3}$$

In general n^{th} order derivative will be

$$\frac{d^n y}{dx^n} = m \cdot (m-1) \cdot (m-2) \dots [m - (n-1)] x^{m-n}$$

$$\frac{d^n y}{dx^n} = m \cdot (m-1) \cdot (m-2) \dots [m - n + 1] x^{m-n}$$

case (i) :- If $m > 0$ and $m > n$, then

$$\frac{d^n y}{dx^n} = \frac{m \cdot (m-1) \cdot (m-2) \dots [m - (n-1)] \cdot (m-n) \dots 2 \cdot 1}{(m-n) \cdot [m-n-1] \dots 2 \cdot 1} x^{m-n}$$

$$\frac{d^n y}{dx^n} = \frac{m! \cdot x^{m-n}}{(m-n)!}$$

case (ii) :- If $m > 0$ and $m = n$, then

$$\frac{d^n y}{dx^n} = \frac{n! \cdot x^{m-n}}{(n-n)!} = \frac{n! \cdot x^0}{0!} = n!$$

case (iii) :- If $m > 0$ and $m < n$, then

$$\frac{d^n y}{dx^n} = 0$$

(ii) Let $y = \frac{1}{ax+b}$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} \left(\frac{1}{ax+b} \right) = \frac{-1}{(ax+b)^2} \cdot \frac{d}{dx} (ax+b)$$

$$\frac{dy}{dx} = \frac{(-1) \cdot a}{(ax+b)^2}$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = (-1)(a) \frac{d}{dx} \left(\frac{1}{(ax+b)^2} \right)$$

$$\frac{d^2 y}{dx^2} = (-1)(a) \frac{-2}{(ax+b)^3} \cdot \frac{d}{dx} (ax+b)$$

$$\frac{d^2 y}{dx^2} = \frac{(-1)^2 \cdot 2 \cdot 1 \cdot a^2}{(ax+b)^3}$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{d^2 y}{dx^2} \right) = (-1)^2 \cdot 2 \cdot 1 \cdot a^2 \cdot \frac{d}{dx} \left(\frac{1}{(ax+b)^3} \right)$$

$$\frac{d^3 y}{dx^3} = (-1)^2 \cdot 2 \cdot 1 \cdot a^2 \cdot \frac{-3}{(ax+b)^4} \cdot \frac{d}{dx} (ax+b)$$

$$\frac{d^3 y}{dx^3} = \frac{(-1)^3 \cdot 3 \cdot 2 \cdot 1 \cdot a^3}{(ax+b)^4}$$

In general n^{th} order derivative will be

$$\frac{d^n y}{dx^n} = \frac{(-1)^n \cdot n \cdot (n-1) \dots 2 \cdot 1 \cdot a^n}{(ax+b)^{n+1}}$$

$$\frac{d^n y}{dx^n} = \frac{(-1)^n \cdot n! \cdot a^n}{(ax+b)^{n+1}}$$

(iii) Let $y = \log x$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (\log x) = \frac{1}{x}$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} \left(\frac{1}{x} \right)$$

$$\frac{d^2 y}{dx^2} = \frac{-1}{x^2} = \frac{(-1)^1}{x^2}$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{d^2 y}{dx^2} \right) = (-1)^1 \frac{d}{dx} \left(\frac{1}{x^2} \right)$$

$$\frac{d^3 y}{dx^3} = (-1)^1 \left(\frac{-2}{x^3} \right) = \frac{(-1)^2 \cdot 1 \cdot 2}{x^3}$$

In general n^{th} order derivative will be

$$\frac{d^n y}{dx^n} = \frac{(-1)^{n-1} \cdot 1 \cdot 2 \cdot 3 \dots (n-1)}{x^n}$$

$$\frac{d^n y}{dx^n} = \frac{(-1)^{n-1} \cdot (n-1)!}{x^n}$$

(iv) Let $y = \sin x$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (\sin x) = \cos x$$

$$\frac{dy}{dx} = \sin \left(\frac{\pi}{2} + x \right)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} \left[\sin \left(\frac{\pi}{2} + x \right) \right]$$

$$\frac{d^2y}{dx^2} = \cos \left(\frac{\pi}{2} + x \right) \frac{d}{dx} \left(\frac{\pi}{2} + x \right)$$

$$\frac{d^2y}{dx^2} = \sin \left(\frac{\pi}{2} + \frac{\pi}{2} + x \right) (1)$$

$$\frac{d^2y}{dx^2} = \sin \left(\frac{2\pi}{2} + x \right)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{d^2y}{dx^2} \right) = \frac{d}{dx} \left[\sin \left(\frac{2\pi}{2} + x \right) \right]$$

$$\frac{d^3y}{dx^3} = \cos \left(\frac{2\pi}{2} + x \right) \frac{d}{dx} \left(\frac{2\pi}{2} + x \right)$$

$$= \sin \left(\frac{\pi}{2} + \frac{2\pi}{2} + x \right) (1)$$

$$\frac{d^3y}{dx^3} = \sin \left(\frac{3\pi}{2} + x \right)$$

In general n^{th} order derivative will be

$$\frac{d^n y}{dx^n} = \sin \left(\frac{n\pi}{2} + x \right)$$

(v) Let $y = \cos (ax + b)$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} [\cos (ax + b)]$$

$$= -\sin (ax + b) \frac{d}{dx} (ax + b)$$

$$= \cos \left(\frac{\pi}{2} + ax + b \right) (a)$$

$$\frac{dy}{dx} = a \cos \left(\frac{\pi}{2} + ax + b \right)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{dx} \left[a \cos \left(\frac{\pi}{2} + ax + b \right) \right]$$

$$\frac{d}{dx} \left(\frac{dy}{dx} \right) = a \frac{d}{dx} \left[\cos \left(\frac{\pi}{2} + ax + b \right) \right]$$

$$\frac{d^2y}{dx^2} = a \left[-\sin \left(\frac{\pi}{2} + ax + b \right) \right] \frac{d}{dx} \left(\frac{\pi}{2} + ax + b \right)$$

$$= a \cos \left(\frac{\pi}{2} + \frac{\pi}{2} + ax + b \right) (a)$$

$$\frac{d^2y}{dx^2} = a^2 \cos \left(\frac{2\pi}{2} + ax + b \right)$$

Differentiate w. r. t. x

$$\frac{d}{dx} \left(\frac{d^2y}{dx^2} \right) = \frac{d}{dx} \left[a^2 \cos \left(\frac{2\pi}{2} + ax + b \right) \right]$$

$$\frac{d}{dx} \left(\frac{d^2y}{dx^2} \right) = a^2 \frac{d}{dx} \left[\cos \left(\frac{2\pi}{2} + ax + b \right) \right]$$


$$\frac{d^3y}{dx^3} = a^2 \left[-\sin \left(\frac{2\pi}{2} + ax + b \right) \right] \frac{d}{dx} \left(\frac{2\pi}{2} + ax + b \right)$$

$$= a^2 \cos \left(\frac{\pi}{2} + \frac{2\pi}{2} + ax + b \right) (a)$$

$$\frac{d^3y}{dx^3} = a^3 \cos \left(\frac{3\pi}{2} + ax + b \right)$$

In general n^{th} order derivative will be

$$\frac{d^n y}{dx^n} = a^n \cos \left(\frac{n\pi}{2} + ax + b \right)$$



(vi) Let $y = e^{ax} \sin (bx + c)$

Differentiate w. r. t. x

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} [e^{ax} \sin (bx + c)] = e^{ax} \frac{d}{dx} [\sin (bx + c)] + [\sin (bx + c)] \frac{d}{dx} (e^{ax}) \\ &= e^{ax} \cos (bx + c) \frac{d}{dx} (bx + c) + \sin (bx + c) \cdot e^{ax} \cdot \frac{d}{dx} (ax) \\ &= e^{ax} [b \cos (bx + c) + a \sin (bx + c)] \\ &= e^{ax} \sqrt{a^2 + b^2} \left[\frac{b}{\sqrt{a^2 + b^2}} \cos (bx + c) + \frac{a}{\sqrt{a^2 + b^2}} \sin (bx + c) \right] \end{aligned}$$

$$\text{Let } \frac{b}{\sqrt{a^2 + b^2}} = \sin \alpha, \frac{a}{\sqrt{a^2 + b^2}} = \cos \alpha, \alpha = \tan^{-1} \left(\frac{b}{a} \right) \quad \dots \text{(I)}$$

$$\frac{dy}{dx} = e^{ax} \sqrt{a^2 + b^2} [\sin \alpha \cdot \cos (bx + c) + \sin (bx + c) \cdot \cos \alpha]$$

$$\frac{dy}{dx} = e^{ax} (a^2 + b^2)^{\frac{1}{2}} \cdot \sin (bx + c + \alpha)$$

Differentiate w. r. t. x

$$\begin{aligned} \frac{d}{dx} \left(\frac{dy}{dx} \right) &= \frac{d}{dx} \left[e^{ax} (a^2 + b^2)^{\frac{1}{2}} \cdot \sin (bx + c + \alpha) \right] \\ &= (a^2 + b^2)^{\frac{1}{2}} \cdot \frac{d}{dx} [e^{ax} \cdot \sin (bx + c + \alpha)] \\ &= (a^2 + b^2)^{\frac{1}{2}} \left[e^{ax} \frac{d}{dx} [\sin (bx + c + \alpha)] + [\sin (bx + c + \alpha)] \frac{d}{dx} [e^{ax}] \right] \\ &= (a^2 + b^2)^{\frac{1}{2}} \left[e^{ax} \cos (bx + c + \alpha) \frac{d}{dx} (bx + c + \alpha) + \sin (bx + c + \alpha) \cdot e^{ax} \frac{d}{dx} (ax) \right] \\ &= e^{ax} (a^2 + b^2)^{\frac{1}{2}} [b \cos (bx + c + \alpha) + a \sin (bx + c + \alpha)] \\ &= e^{ax} (a^2 + b^2)^{\frac{1}{2}} \sqrt{a^2 + b^2} \left[\frac{b}{\sqrt{a^2 + b^2}} \cos (bx + c + \alpha) + \frac{a}{\sqrt{a^2 + b^2}} \sin (bx + c + \alpha) \right] \\ \frac{d^2 y}{dx^2} &= e^{ax} (a^2 + b^2)^{\frac{2}{2}} [\sin \alpha \cos (bx + c + \alpha) + \sin (bx + c + \alpha) \cos \alpha] \quad \dots \text{[from (I)]} \\ \frac{d^2 y}{dx^2} &= e^{ax} (a^2 + b^2)^{\frac{2}{2}} \cdot \sin (bx + c + 2\alpha) \end{aligned}$$

Similarly,

$$\frac{d^3 y}{dx^3} = e^{ax} (a^2 + b^2)^{\frac{3}{2}} \cdot \sin (bx + c + 3\alpha)$$

In general n^{th} order derivative will be

$$\frac{d^n y}{dx^n} = e^{ax} (a^2 + b^2)^{\frac{n}{2}} \cdot \sin (bx + c + n\alpha) \text{ where } \alpha = \tan^{-1} \left(\frac{b}{a} \right).$$

EXERCISE 1.5

(1) Find the second order derivative of the following :

(i) $2x^5 - 4x^3 - \frac{2}{x^2} - 9$ (ii) $e^{2x} \cdot \tan x$

(iii) $e^{4x} \cdot \cos 5x$ (iv) $x^3 \log x$

(v) $\log(\log x)$ (iv) x^x

(2) Find $\frac{d^2y}{dx^2}$ of the following :

(i) $x = a(\theta - \sin \theta), y = a(1 - \cos \theta)$

(ii) $x = 2at^2, y = 4at$

(iii) $x = \sin \theta, y = \sin^3 \theta$ when $\theta = \frac{\pi}{2}$

(iv) $x = a \cos \theta, y = b \sin \theta$ at $\theta = \frac{\pi}{4}$

(3) (i) If $x = at^2$ and $y = 2at$ then show that

$$xy \frac{d^2y}{dx^2} + a = 0$$

(ii) If $y = e^{m \tan^{-1} x}$, show that

$$(1 + x^2) \frac{d^2y}{dx^2} + (2x - m) \frac{dy}{dx} = 0$$

(iii) If $x = \cos t, y = e^{mt}$ show that

$$(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - m^2y = 0$$

(iv) If $y = x + \tan x$, show that

$$\cos^2 x \cdot \frac{d^2y}{dx^2} - 2y + 2x = 0$$

(v) If $y = e^{ax} \cdot \sin(bx)$, show that

$$y_2 - 2ay_1 + (a^2 + b^2)y = 0$$

(vi) If $\sec^{-1} \left(\frac{7x^3 - 5y^3}{7x^3 + 5y^3} \right) = m$,

show that $\frac{d^2y}{dx^2} = 0$.

(vii) If $2y = \sqrt{x+1} + \sqrt{x-1}$,

show that $4(x^2 - 1)y_2 + 4xy_1 - y = 0$.

(viii) If $y = \log(x + \sqrt{x^2 + a^2})^m$,

show that $(x^2 + a^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 0$

(ix) If $y = \sin(m \cos^{-1} x)$ then show that

$$(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + m^2y = 0$$

(x) If $y = \log(\log 2x)$, show that

$$xy_2 + y_1(1 + xy_1) = 0.$$

(xi) If $x^2 + 6xy + y^2 = 10$, show that

$$\frac{d^2y}{dx^2} = \frac{80}{(3x+y)^3}.$$

(xii) If $x = a \sin t - b \cos t, y = a \cos t + b \sin t$,

show that $\frac{d^2y}{dx^2} = -\frac{x^2 + y^2}{y^3}$.

(4) Find the n^{th} derivative of the following :

(i) $(ax + b)^m$ (ii) $\frac{1}{x}$

(iii) e^{ax+b} (iv) a^{px+q}

(v) $\log(ax + b)$ (vi) $\cos x$

(vii) $\sin(ax + b)$ (viii) $\cos(3 - 2x)$

(ix) $\log(2x + 3)$

(x) $\frac{1}{3x - 5}$

(xi) $y = e^{ax} \cdot \cos(bx + c)$

(xii) $y = e^{8x} \cdot \cos(6x + 7)$



Let us Remember

✿ If a function $f(x)$ is differentiable at $x = a$ then it is continuous at $x = a$, but the converse is not true.

✿ **Chain Rule :** If y is differentiable function of u and u is differentiable function of x then y is differentiable function of x and $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

✿ If $y = f(x)$ is a differentiable of x such that the inverse function $x = f^{-1}(y)$ exists then

$$\frac{dy}{dx} = \frac{1}{\frac{dy}{dx}}, \quad \text{where } \frac{dy}{dx} \neq 0$$

✿ **Derivatives of Inverse Trigonometric functions :**

$f(x)$	$\sin^{-1} x$	$\cos^{-1} x$	$\tan^{-1} x$	$\cot^{-1} x$	$\sec^{-1} x$	$\operatorname{cosec}^{-1} x$
$f'(x)$	$\frac{1}{\sqrt{1-x^2}},$ $ x < 1$	$-\frac{1}{\sqrt{1-x^2}},$ $ x < 1$	$\frac{1}{1+x^2}$ $x \in \mathbb{R}$	$-\frac{1}{1+x^2}$ $x \in \mathbb{R}$	$\frac{1}{x\sqrt{x^2-1}}$ $ x < 1$	$-\frac{1}{x\sqrt{x^2-1}}$ $ x < 1$

✿ This is a simple shortcut to find the derivative of (function)^(function)

$$\frac{d}{dx} f^g = f^g \left[\frac{g}{f} \cdot f' + (\log f) \cdot g' \right]$$

✿ If $y = f(t)$ and $y = g(t)$ is a differentiable of t such that y is a function of x then

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}, \quad \text{where } \frac{dx}{dt} \neq 0$$

✿ Implicit function of the form $x^m y^n = (x + y)^{m+n}$, $m, n \in \mathbb{R}$ always have the first order derivative

$$\frac{dy}{dx} = \frac{y}{x} \text{ and second order derivative } \frac{d^2y}{dx^2} = 0$$

MISCELLANEOUS EXERCISE 1

(I) **Choose the correct option from the given alternatives :**

(1) Let $f(1) = 3, f'(1) = -\frac{1}{3}, g(1) = -4$ and $g'(1) = -\frac{8}{3}$. The derivative of $\sqrt{[f(x)]^2 + [g(x)]^2}$

w. r. t. x at $x = 1$ is

(A) $-\frac{29}{15}$

(B) $\frac{7}{3}$

(C) $\frac{31}{15}$

(D) $\frac{29}{15}$

- (2) If $y = \sec(\tan^{-1} x)$ then $\frac{dy}{dx}$ at $x = 1$, is equal to :
 (A) $\frac{1}{2}$ (B) 1 (C) $\frac{1}{\sqrt{2}}$ (D) $\sqrt{2}$
- (3) If $f(x) = \sin^{-1}\left(\frac{4^{x+\frac{1}{2}}}{1+2^{4x}}\right)$, which of the following is not the derivative of $f(x)$
 (A) $\frac{2 \cdot 4^x \log 4}{1+4^{2x}}$ (B) $\frac{4^{x+1} \log 2}{1+4^{2x}}$ (C) $\frac{4^{x+1} \log 4}{1+4^{4x}}$ (D) $\frac{2^{2(x+1)} \log 2}{1+2^{4x}}$
- (4) If $x^y = y^x$, then $\frac{dy}{dx} = \dots$
 (A) $\frac{x(x \log y - y)}{y(y \log x - x)}$ (B) $\frac{y(y \log x - x)}{x(x \log y - y)}$ (C) $\frac{y^2(1 - \log x)}{x^2(1 - \log y)}$ (D) $\frac{y(1 - \log x)}{x(1 - \log y)}$
- (5) If $y = \sin(2 \sin^{-1} x)$, then $\frac{dy}{dx} = \dots$
 (A) $\frac{2-4x^2}{\sqrt{1-x^2}}$ (B) $\frac{2+4x^2}{\sqrt{1-x^2}}$ (C) $\frac{4x^2-1}{\sqrt{1-x^2}}$ (D) $\frac{1-2x^2}{\sqrt{1-x^2}}$
- (6) If $y = \tan^{-1}\left(\frac{x}{1+\sqrt{1-x^2}}\right) + \sin\left[2 \tan^{-1}\left(\sqrt{\frac{1-x}{1+x}}\right)\right]$, then $\frac{dy}{dx} = \dots$
 (A) $\frac{x}{\sqrt{1-x^2}}$ (B) $\frac{1-2x}{\sqrt{1-x^2}}$ (C) $\frac{1-2x}{2\sqrt{1-x^2}}$ (D) $\frac{1-2x^2}{\sqrt{1-x^2}}$
- (7) If y is a function of x and $\log(x+y) = 2xy$, then the value of $y'(0) = \dots$
 (A) 2 (B) 0 (C) -1 (D) 1
- (8) If g is the inverse of a function f and $f'(x) = \frac{1}{1+x^7}$, then the value of $g'(x)$ is equal to :
 (A) $1+x^7$ (B) $\frac{1}{1+[g(x)]^7}$ (C) $1+[g(x)]^7$ (D) $7x^6$
- (9) If $x\sqrt{y+1} + y\sqrt{x+1} = 0$ and $x \neq y$ then $\frac{dy}{dx} = \dots$
 (A) $\frac{1}{(1+x)^2}$ (B) $-\frac{1}{(1+x)^2}$ (C) $(1+x)^2$ (D) $-\frac{x}{x+1}$
- (10) If $y = \tan^{-1}\left(\sqrt{\frac{a-x}{a+x}}\right)$, where $-a < x < a$ then $\frac{dy}{dx} = \dots$
 (A) $\frac{x}{\sqrt{a^2-x^2}}$ (B) $\frac{a}{\sqrt{a^2-x^2}}$ (C) $-\frac{1}{2\sqrt{a^2-x^2}}$ (D) $\frac{1}{2\sqrt{a^2-x^2}}$

(11) If $x = a(\cos \theta + \theta \sin \theta)$, $y = a(\sin \theta - \theta \cos \theta)$ then $\left[\frac{d^2y}{dx^2}\right]_{\theta=\frac{\pi}{4}} = \dots$

- (A) $\frac{8\sqrt{2}}{a\pi}$ (B) $-\frac{8\sqrt{2}}{a\pi}$ (C) $\frac{a\pi}{8\sqrt{2}}$ (D) $\frac{4\sqrt{2}}{a\pi}$

(12) If $y = a \cos(\log x)$ and $A\frac{d^2y}{dx^2} + B\frac{dy}{dx} + C = 0$, then the values of A, B, C are ...

- (A) $x^2, -x, -y$ (B) x^2, x, y (C) $x^2, x, -y$ (D) $x^2, -x, y$

(II) Solve the following :

- (1) $f(x) = -x,$ for $-2 \leq x < 0$ $g(x) = 6 - 3x,$ for $0 \leq x \leq 2$
 $= 2x,$ for $0 \leq x \leq 2$ $= \frac{2x - 4}{3},$ for $2 < x \leq 7$
 $= \frac{18 - x}{4},$ for $2 < x \leq 7$

Let $u(x) = f[g(x)], v(x) = g[f(x)]$ and $w(x) = g[g(x)]$.

Find each derivative at $x = 1$, if it exists i.e. find $u'(1), v'(1)$ and $w'(1)$. if it doesn't exist then explain why ?

(2) The values of $f(x), g(x), f'(x)$ and $g'(x)$ are given in the following table.

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
-1	3	2	-3	4
2	2	-1	-5	-4

Match the following.

A Group - Function	B Group - Derivative
(A) $\frac{d}{dx} [f(g(x))] \text{ at } x = -1$	1. -16
(B) $\frac{d}{dx} [g(f(x) - 1)] \text{ at } x = -1$	2. 20
(C) $\frac{d}{dx} [f(f(x) - 3)] \text{ at } x = 2$	3. -20
(D) $\frac{d}{dx} [g(g(x))] \text{ at } x = 2$	4. 15
	5. 12

(3) Suppose that the functions f and g and their derivatives with respect to x have the following values at $x = 0$ and $x = 1$.

x	$f(x)$	$g(x)$	$f'(x)$	$g'(x)$
0	1	1	5	$\frac{1}{3}$
1	3	-4	$-\frac{1}{3}$	$-\frac{8}{3}$

- (i) The derivative of $f[g(x)]$ w. r. t. x at $x = 0$ is
- (ii) The derivative of $g[f(x)]$ w. r. t. x at $x = 0$ is
- (iii) The value of $\left[\frac{d}{dx} [x^{10} + f(x)]^{-2}\right]_{x=1}$ is
- (iv) The derivative of $f[(x + g(x))]$ w. r. t. x at $x = 0$ is

(4) Differentiate the following *w. r. t. x*

(i) $\sin \left[2 \tan^{-1} \left(\sqrt{\frac{1-x}{1+x}} \right) \right]$

(ii) $\sin^2 \left[\cot^{-1} \left(\sqrt{\frac{1+x}{1-x}} \right) \right]$

(iii) $\tan^{-1} \left[\frac{\sqrt{x}(3-x)}{1-3x} \right]$

(iv) $\cos^{-1} \left(\frac{\sqrt{1+x} - \sqrt{1-x}}{2} \right)$

(v) $\tan^{-1} \left(\frac{x}{1+6x^2} \right) + \cot^{-1} \left(\frac{1-10x^2}{7x} \right)$

(vi) $\tan^{-1} \left[\frac{\sqrt{1+x^2+x}}{\sqrt{1+x^2-x}} \right]$

(5) (i) If $\sqrt{y+x} + \sqrt{y-x} = c$, then show that $\frac{dy}{dx} = \frac{y}{x} - \sqrt{\frac{y^2}{x^2} - 1}$.

(ii) If $x\sqrt{1-y^2} + y\sqrt{1-x^2} = 1$, then show that $\frac{dy}{dx} = -\sqrt{\frac{1-y^2}{1-x^2}}$.

(iii) If $x \sin (a+y) + \sin a \cos (a+y) = 0$, then show that $\frac{dy}{dx} = \frac{\sin^2 (a+y)}{\sin a}$.

(iv) If $\sin y = x \sin (a+y)$, then show that $\frac{dy}{dx} = \frac{\sin^2 (a+y)}{\sin a}$.

(v) If $x = e^{\frac{x}{y}}$, then show that $\frac{dy}{dx} = \frac{x-y}{x \log x}$.

(vi) If $y = f(x)$ is a differentiable function then show that $\frac{d^2x}{dy^2} = -\left(\frac{dy}{dx}\right)^{-3} \cdot \frac{d^2y}{dx^2}$.

(6) (i) Differentiate $\tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$ *w. r. t.* $\tan^{-1} \left(\frac{2x\sqrt{1-x^2}}{1-2x^2} \right)$.

(ii) Differentiate $\log \left(\frac{\sqrt{1+x^2+x}}{\sqrt{1+x^2-x}} \right)$ *w. r. t.* $\cos (\log x)$.

(iii) Differentiate $\tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$ *w. r. t.* $\cos^{-1} \left(\sqrt{\frac{1+\sqrt{1+x^2}}{2\sqrt{1+x^2}}} \right)$.

(7) (i) If $y^2 = a^2 \cos^2 x + b^2 \sin^2 x$, show that $y + \frac{d^2y}{dx^2} = \frac{a^2b^2}{y^3}$.

(ii) If $\log y = \log (\sin x) - x^2$, show that $\frac{d^2y}{dx^2} + 4x \frac{dy}{dx} + (4x^2 + 3)y = 0$.

(iii) If $x = a \cos \theta, y = b \sin \theta$, show that $a^2 \left[y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx} \right)^2 \right] + b^2 = 0$.

(iv) If $y = A \cos (\log x) + B \sin (\log x)$, show that $x^2 y_2 + x y_1 + y = 0$.

(v) If $y = A e^{mx} + B e^{nx}$, show that $y_2 - (m+n)y_1 + (mn)y = 0$.



2. APPLICATIONS OF DERIVATIVES



Let us Study

- Applications of Derivatives to Tangents and Normals
- Derivative as a rate measure
- Approximations
- Rolle's Theorem and Lagrange's Mean Value Theorem.
- Increasing and Decreasing Functions
- Maxima and Minima



Let us Recall

- Continuous functions.
- Derivatives of Composite, Inverse Trigonometric, Logarithmic, Parametric functions.
- Relation between derivative and slope.
- Higher Order Derivatives.

2.1.1 Introduction :

In the previous chapter we have studied the derivatives of various functions such as composite functions, Inverse Trigonometric functions, Logarithmic functions etc. and also the relation between Derivative and slope of the tangent. In this chapter we are going to study various applications of differentiation such as application to (i) Geometry, (ii) Rate measure (iii) Approximations (iv) Rolle's Theorem and Lagrange's Mean Value Theorem (v) Increasing and Decreasing functions and (vi) Maxima and Minima.



Let us Learn

2.1.2 Application of Derivative in Geometry :

In the previous chapter we have studied the relation between derivative and slope of a line or slope of a tangent to the curve at a given point on it.

Let $y = f(x)$ be a continuous function of x representing a curve in XY- plane and $P(x_1, y_1)$ be any point on the curve.

Then $\left[\frac{dy}{dx} \right]_{(x_1, y_1)} = [f'(x)]_{(x_1, y_1)}$ represents slope, also called gradient, of the tangent to the curve at

$P(x_1, y_1)$. The normal is perpendicular to the tangent. Hence, the slope of the normal at P will be the negative of reciprocal of the slope of tangent at P . Let m and m' be the slopes of tangent and normal respectively,

then $m = \left[\frac{dy}{dx} \right]_{(x_1, y_1)}$ and $m' = -\frac{1}{\left[\frac{dy}{dx} \right]_{(x_1, y_1)}}$ if $\left[\frac{dy}{dx} \right]_{(x_1, y_1)} \neq 0$.

Equation of tangent at $P(x_1, y_1)$ is given by $y - y_1 = m(x - x_1)$ i.e. $y - y_1 = \left[\frac{dy}{dx} \right]_{(x_1, y_1)} (x - x_1)$

and equation of normal at $P(x_1, y_1)$ is given by

$$y - y_1 = m'(x - x_1) \text{ where } m' = -\frac{1}{\left[\frac{dy}{dx} \right]_{(x_1, y_1)}}$$



SOLVED EXAMPLES

Ex. 1 : Find the equations of tangent and normal to the curve at the given point on it.

(i) $y = 2x^3 - x^2 + 2$ at $\left(\frac{1}{2}, 2\right)$

(ii) $x^3 + 2x^2y - 9xy = 0$ at $(2, 1)$

(iii) $x = 2 \sin^3 \theta, y = 3 \cos^3 \theta$ at $\theta = \frac{\pi}{4}$

Solution :

(i) Given that : $y = 2x^3 - x^2 + 2$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (2x^3 - x^2 + 2) = 6x^2 - 2x$$

$$\text{Slope of tangent at } \left(\frac{1}{2}, 2\right) = m = 6\left(\frac{1}{2}\right)^2 - 2\left(\frac{1}{2}\right)$$

$$\therefore m = \frac{1}{2}$$

$$\text{Slope of normal at } \left(\frac{1}{2}, 2\right) = m' = -2$$

Equation of tangent is given by

$$y - 2 = \frac{1}{2} \left(x - \frac{1}{2}\right) \Rightarrow 2y - 4 = \frac{2x - 1}{2}$$

$$4y - 8 = 2x - 1 \Rightarrow 2x - 4y + 7 = 0$$

Equation of normal is given by

$$y - 2 = -2 \left(x - \frac{1}{2}\right) \Rightarrow y - 2 = -2x + 1$$

$$2x + y - 3 = 0$$

(ii) Given that : $x^3 + 2x^2y - 9xy = 0$

Differentiate w. r. t. x

$$3x^2 + 2 \left(x^2 \frac{dy}{dx} + y \frac{d}{dx} (x^2)\right) - 9 \left(x \frac{dy}{dx} + y \frac{d}{dx} (x)\right) = 0$$

$$3x^2 + 2x^2 \frac{dy}{dx} + 4xy - 9x \frac{dy}{dx} - 9y = 0$$

$$(2x^2 - 9x) \frac{dy}{dx} = 9y - 4xy - 3x^2 \therefore \frac{dy}{dx} = \frac{9y - 4xy - 3x^2}{2x^2 - 9x}$$

Slope of tangent at $(2, 1)$

$$\left(\frac{dy}{dx}\right)_{(1, 2)} = m = \frac{9(1) - 4(2)(1) - 3(4)}{2(4) - 9(1)} = \frac{9 - 8 - 12}{8 - 9}$$

$$m = \frac{-11}{-1} \therefore m = 11$$

$$\text{Slope of normal at } (2, 1) = m' = -\frac{1}{11}$$

Equation of tangent is given by

$$y - 1 = 11(x - 2) \Rightarrow 11x - y - 21 = 0$$

Equation of normal is given by

$$y - 1 = -\frac{1}{11} (x - 2) \Rightarrow 11y - 11 = -x + 2$$

$$x + 11y - 13 = 0$$

(iii) Given that : $y = 3 \cos^3 \theta$

Differentiate w. r. t. θ

$$\frac{dy}{d\theta} = 3 \frac{d}{d\theta} (\cos \theta)^3 = 9 \cos^2 \theta \frac{d}{d\theta} (\cos \theta)$$

$$\therefore \frac{dy}{d\theta} = -9 \cos^2 \theta \sin \theta$$

Now, $x = 2 \sin^3 \theta$

Differentiate w. r. t. θ

$$\frac{dx}{d\theta} = 2 \frac{d}{d\theta} (\sin \theta)^3 = 6 \sin^2 \theta \frac{d}{d\theta} (\sin \theta)$$

$$\therefore \frac{dx}{d\theta} = 6 \sin^2 \theta \cos \theta$$

We know that

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = -\frac{9 \cos^2 \theta \sin \theta}{6 \sin^2 \theta \cos \theta} = -\frac{3}{2} \cot \theta$$

Slope of tangent at $\theta = \frac{\pi}{4}$ is

$$\left(\frac{dy}{dx}\right)_{\theta=\frac{\pi}{4}} = m = -\frac{3}{2} \cot\left(\frac{\pi}{4}\right) = -\frac{3}{2}$$

$$\text{Slope of normal at } \left(\theta = \frac{\pi}{4}\right) = m' = \frac{2}{3}$$

When, $\theta = \frac{\pi}{4}$

$$x = 2 \sin^3\left(\frac{\pi}{4}\right) = 2 \left(\frac{1}{\sqrt{2}}\right)^3 = \frac{1}{\sqrt{2}}$$

$$y = 3 \cos^3\left(\frac{\pi}{4}\right) = 3 \left(\frac{1}{\sqrt{2}}\right)^3 = \frac{3}{2\sqrt{2}}$$

$$\therefore \text{ The point is } P = \left(\frac{1}{\sqrt{2}}, \frac{3}{2\sqrt{2}}\right)$$

Equation of tangent at P is given by

$$y - \frac{3}{2\sqrt{2}} = -\frac{3}{2} \left(x - \frac{1}{\sqrt{2}}\right) \Rightarrow y - \frac{3}{2\sqrt{2}} = -\frac{3x}{2} + \frac{3}{2\sqrt{2}}$$

$$\frac{3x}{2} + y - \frac{3}{\sqrt{2}} = 0 \quad \text{i.e. } 3x + 2y - 3\sqrt{2} = 0$$

Equation of normal is given by

$$y - \frac{3}{2\sqrt{2}} = \frac{2}{3} \left(x - \frac{1}{\sqrt{2}}\right) \Rightarrow y - \frac{3}{2\sqrt{2}} = \frac{2x}{3} - \frac{2}{3\sqrt{2}}$$

$$\frac{2x}{3} - y - \frac{2}{3\sqrt{2}} + \frac{3}{2\sqrt{2}} = 0$$

$$\text{i.e. } 4\sqrt{2}x - 6\sqrt{2}y + 5 = 0 \quad \dots \text{ [Multiply by } 6\sqrt{2} \text{]}$$

Ex. 2 : Find points on the curve given by $y = x^3 - 6x^2 + x + 3$ where the tangents are parallel to the line $y = x + 5$.

Solution : Equation of curve is $y = x^3 - 6x^2 + x + 3$

Differentiate w. r. t. x

$$\frac{dy}{dx} = \frac{d}{dx} (x^3 - 6x^2 + x + 3) = 3x^2 - 12x + 1$$

Given that the tangent is parallel to $y = x + 5$ whose slope is 1.

$$\therefore \text{Slope of tangent} = \frac{dy}{dx} = 1 \Rightarrow 3x^2 - 12x + 1 = 1$$

$$3x(x - 4) = 0 \quad \text{so, } x = 0 \text{ or } x = 4$$

$$\text{When } x = 0, y = (0)^3 - 6(0)^2 + (0) + 3 = 3$$

$$\text{When } x = 4, y = (4)^3 - 6(4)^2 + (4) + 3 = -25$$

So the required points on the curve are $(0, 3)$ and $(4, -25)$.

2.1.3 Derivative as a Rate measure :

If $y = f(x)$ is the given function then a change in x from x_1 to x_2 is generally denoted by $\delta x = x_2 - x_1$ and the corresponding change in y is denoted by $\delta y = f(x_2) - f(x_1)$. The difference quotient $\frac{\delta y}{\delta x} = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$ is called the **average rate of change** with respect to x . This can also be interpreted geometrically as the slope of the secant line joining the points $P(x_1, f(x_1))$ and $Q(x_2, f(x_2))$ on the graph of function $y = f(x)$.

Consider the average rate of change over smaller and smaller intervals by letting x_2 to approach x_1 and therefore letting δx to approach 0. The limit of these average rates of change is called the **instantaneous rate of change** of y with respect to x at $x = x_1$, which is interpreted as the slope of the tangent to the curve $y = f(x)$ at $P(x_1, f(x_1))$. Therefore instantaneous rate of change is given by

$$\lim_{\delta x \rightarrow 0} \left(\frac{\delta y}{\delta x} \right) = \lim_{x_2 \rightarrow x_1} \left(\frac{f(x_2) - f(x_1)}{x_2 - x_1} \right)$$

We recognize this limit as being the derivative of $f(x)$ at $x = x_1$, i.e. $f'(x_1)$. We know that one interpretation of the derivative $f'(a)$ is the instantaneous rate of change of $y = f(x)$ with respect x when $x = a$. The other interpretation is $f'(a)$ is the slope of the tangent to $y = f(x)$ at $(a, f(a))$.



SOLVED EXAMPLES

Ex. 1 : A stone is dropped in to a quiet lake and waves in the form of circles are generated, radius of the circular wave increases at the rate of 5 cm/ sec. At the instant when the radius of the circular wave is 8 cm, how fast the area enclosed is increasing ?

Solution : Let R be the radius and A be the area of the circular wave.

$$\therefore A = \pi \cdot R^2$$

Differentiate w. r. t. t

$$\frac{dA}{dt} = \pi \frac{d}{dt} (R^2)$$

$$\frac{dA}{dt} = 2\pi R \frac{dR}{dt} \quad \dots (I)$$

Given that $\frac{dR}{dt} = 5$ cm/sec.

Thus when $R = 8$ cm, from (I) we get,

$$\frac{dA}{dt} = 2\pi(8)(5) = 80\pi$$

Hence when the radius of the circular wave is 8 cm, the area of the circular wave is increasing at the rate of 80π cm²/ sec.

Ex. 2 : The volume of the spherical ball is increasing at the rate of 4π cc/sec. Find the rate at which the radius and the surface area are changing when the volume is 288π cc.

Solution : Let R be the radius, S be the surface area and V be the volume of the spherical ball.

$$V = \frac{4}{3} \pi R^3 \quad \dots (I)$$

Differentiate w. r. t. t

$$\frac{dV}{dt} = \frac{4\pi}{3} \cdot \frac{d}{dt} (R^3)$$

$$4\pi = \frac{4\pi}{3} \cdot 3R^2 \frac{dR}{dt} \quad \dots \text{ [Given } \frac{dV}{dt} = 4\pi \text{ cc/sec]}$$

$$\frac{dR}{dt} = \frac{1}{R^2} \quad \dots (II)$$

When volume is 288π cc.

i.e. $\frac{4}{3} \pi \cdot R^3 = 288\pi$ we get, $R^3 = 216 \Rightarrow R = 6$... [From (I)]

From (II) we get, $\frac{dR}{dt} = \frac{1}{36}$

So, the radius of the spherical ball is increasing at the rate of $\frac{1}{36}$ cc/sec.

Now, $S = 4\pi R^2$

Differentiate w. r. t. t .

$$\frac{dS}{dt} = 4\pi \frac{d}{dt} (R^2) = 8\pi R \frac{dR}{dt}$$

So, when $R = 6$ cm

$$\left[\frac{dS}{dt} \right]_{R=6} = 8\pi(6) \frac{1}{36} = \frac{4\pi}{3}$$

\therefore Surface area is increasing at the rate of $\frac{4\pi}{3}$ cm²/ sec.



Ex. 3 : Water is being poured at the rate of $36 \text{ m}^3/\text{sec}$ in to a cylindrical vessel of base radius 3 meters. Find the rate at which water level is rising.

Solution : Let R be the radius of the base, H be the height and V be the volume of the cylindrical vessel at any time t . R , V and H are functions of t .

$$V = \pi R^2 H$$

$$V = \pi(3)^2 H = 9\pi H \quad \dots \text{ [Given : } R = 3 \text{]}$$

Differentiate w. r. t. t

$$\frac{dV}{dt} = 9\pi \frac{dH}{dt}$$

$$\frac{dH}{dt} = \frac{1}{9\pi} \cdot \frac{dV}{dt} \quad \dots \text{ (I)}$$

Given that,

$$\frac{dV}{dt} = 36 \text{ m}^3/\text{sec} \quad \dots \text{ (II)}$$

From (I) we get,
$$\frac{dH}{dt} = \frac{1}{9\pi} \cdot (36) = \frac{4}{\pi}$$

\therefore Water level is rising at the rate of $\frac{4}{\pi}$ meter/sec.

Ex. 4 : A man of height 180 cm is moving away from a lamp post at the rate of 1.2 meters per second. If the height of the lamp post is 4.5 meters, find the rate at which (i) his shadow is lengthening. (ii) the tip of the shadow is moving.

Solution : Let OA be the lamp post, MN be the man, $MB = x$ be the length of shadow and $OM = y$ be the distance of the man from the lamp post at time t . Given that man is moving away from the lamp post at the rate of 1.2 meter/sec. x and y are functions of t .

Hence $\frac{dy}{dt} = 1.2$. The rate at which shadow is lengthening = $\frac{dx}{dt}$.

B is the tip of the shadow and it is at a distance of $(x + y)$ from the post.

$$\frac{x}{1.8} = \frac{x + y}{4.5} \quad \text{i.e. } 45x = 18x + 18y \quad \text{i.e. } 27x = 18y$$

$$\therefore x = \frac{2y}{3}$$


Differentiate w. r. t. t

$$\frac{dx}{dt} = \frac{2}{3} \times \frac{dy}{dt} = \frac{2}{3} \times 1.2 = 0.8 \text{ meter/sec.}$$

rate at which the tip of the shadow is moving is given by

$$\frac{d}{dt}(x + y) = \frac{dx}{dt} + \frac{dy}{dt}$$

$$\therefore \frac{d}{dt}(x + y) = 0.8 + 1.2 = 2 \text{ meter/sec.}$$



Shadow is lengthening at the rate of 0.8 meter/ sec. and its tip is moving at the rate of 2 meters/sec.

2.1.4 Velocity, Acceleration and Jerk :

If $s = f(t)$ is the displacement function of a particle that moves along a straight line, then $f'(t)$ is the rate of change of the displacement s with respect to the time t . In other words, $f'(t)$ is the **velocity** of the particle. The **speed** of the particle is the absolute value of the velocity, that is, $|f'(t)|$.

The rate of change of velocity with respect to time is called the **acceleration** of the particle denoted by $a(t)$. Thus the acceleration function is the derivative of the velocity function and is therefore the second derivative of the position function $s = f(t)$.

$$\text{Thus, } a = \frac{dv}{dt} = \frac{d^2s}{dt^2} \quad \text{i.e. } a(t) = v'(t) = s''(t).$$

Let us consider the third derivative of the position function $s = f(t)$ of an object that moves along a straight line. $s'''(t) = v''(t) = a'(t)$ is derivative of the acceleration function and is called the **Jerk** (j).

Thus, $j = \frac{da}{dt} = \frac{d^3s}{dt^3}$. Hence the jerk j is the rate of change of acceleration. It is aptly named because a jerk means a sudden change in acceleration, which causes an abrupt movement in a vehicle.



SOLVED EXAMPLES

Ex. 1 : A car is moving in such a way that the distance it covers, is given by the equation $s = 4t^2 + 3t$ where s is in meters and t is in seconds. What would be the velocity and the acceleration of the car at time $t = 20$ second ?

Solution : Let v be the velocity and a be the acceleration of the car.

Distance traveled by the car is given by

$$s = 4t^2 + 3t$$

Differentiate w. r. t. t .

\therefore Velocity of the car is given by

$$v = \frac{ds}{dt} = \frac{d}{dt}(4t^2 + 3t) = 8t + 3 \quad \dots \text{(I)}$$

and Acceleration of the car is given by

$$a = \frac{d}{dt} \left(\frac{dv}{dt} \right) = \frac{d}{dt}(8t + 3) = 8 \quad \dots \text{(II)}$$

Put $t = 20$ in (I),

\therefore Velocity of the car, $v_{t=20} = 8(20) + 3 = 163$ m/sec.

Put $t = 20$ in (II),

\therefore Acceleration of the car, $a_{t=20} = 8$ m/sec².

Note : In this problem, the acceleration is independent of time. Such a motion is said to be uniformly accelerated motion.

Ex. 2 : The displacement of a particle at time t is given by $s = 2t^3 - 5t^2 + 4t - 3$. Find the time when the acceleration is 14 ft/ sec², the velocity and the displacement at that time.

Solution : Displacement of a particle is given by

$$s = 2t^3 - 5t^2 + 4t - 3 \quad \dots \text{(I)}$$

Differentiate w. r. t. t .

$$\text{Velocity, } v = \frac{ds}{dt} = \frac{d}{dt}(2t^3 - 5t^2 + 4t - 3)$$

$$\therefore v = 6t^2 - 10t + 4 \quad \dots \text{(II)}$$

$$\text{Acceleration, } a = \frac{dv}{dt} = \frac{d}{dt}(6t^2 - 10t + 4)$$

$$\therefore a = 12t - 10 \quad \dots \text{(III)}$$

Given : Acceleration = 14 ft/ sec².

$$\therefore 12t - 10 = 14 \Rightarrow 12t = 24 \Rightarrow t = 2$$

So, the particle reaches an acceleration of 14 ft/ sec² in 2 seconds.

Velocity, when $t = 2$ is

$$\therefore v_{t=2} = 6(2)^2 - 10(2) + 4 = 8 \text{ ft/ sec.}$$

Displacement when $t = 2$ is

$$\therefore s_{t=2} = 2(2)^3 - 5(2)^2 + 4(2) - 3 = 1 \text{ foot.}$$

Hence the velocity is 8 ft/ sec and the displacement is 1 foot after 2 seconds.

EXERCISE 2.1

- (1) Find the equations of tangents and normals to the curve at the point on it.
- (i) $y = x^2 + 2e^x + 2$ at $(0, 4)$
(ii) $x^3 + y^3 - 9xy = 0$ at $(2, 4)$
(iii) $x^2 - \sqrt{3}xy + 2y^2 = 5$ at $(\sqrt{3}, 2)$
(iv) $2xy + \pi \sin y = 2\pi$ at $\left(1, \frac{\pi}{2}\right)$
(v) $x \sin 2y = y \cos 2x$ at $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$
(vi) $x = \sin \theta$ and $y = \cos 2\theta$ at $\theta = \frac{\pi}{6}$
(vii) $x = \sqrt{t}, y = t - \frac{1}{\sqrt{t}}$ at $t = 4$.
- (2) Find the point on the curve $y = \sqrt{x-3}$ where the tangent is perpendicular to the line $6x + 3y - 5 = 0$.
- (3) Find the points on the curve $y = x^3 - 2x^2 - x$ where the tangents are parallel to $3x - y + 1 = 0$.
- (4) Find the equations of the tangents to the curve $x^2 + y^2 - 2x - 4y + 1 = 0$ which are parallel to the X-axis.
- (5) Find the equations of the normals to the curve $3x^2 - y^2 = 8$, which are parallel to the line $x + 3y = 4$.
- (6) If the line $y = 4x - 5$ touches the curve $y^2 = ax^3 + b$ at the point $(2, 3)$ find a and b .
- (7) A particle moves along the curve $6y = x^2 + 2$. Find the points on the curve at which y-coordinate is changing 8 times as fast as the X-coordinate.
- (8) A spherical soap bubble is expanding so that its radius is increasing at the rate of 0.02 cm/sec. At what rate is the surface area is increasing, when its radius is 5 cm?
- (9) The surface area of a spherical balloon is increasing at the rate of 2 cm²/ sec. At what rate the volume of the balloon is increasing when radius of the balloon is 6 cm?
- (10) If each side of an equilateral triangle increases at the rate of $\sqrt{2}$ cm/ sec, find the rate of increase of its area when its side of length 3 cm.
- (11) The volume of a sphere increase at the rate of 20 cm³/ sec. Find the rate of change of its surface area when its radius is 5 cm.
- (12) The edge of a cube is decreasing at the rate of 0.6 cm/sec. Find the rate at which its volume is decreasing when the edge of the cube is 2 cm.
- (13) A man of height 2 meters walks at a uniform speed of 6 km/hr away from a lamp post of 6 meters high. Find the rate at which the length of the shadow is increasing.
- (14) A man of height 1.5 meters walks toward a lamp post of height 4.5 meters, at the rate of $\left(\frac{3}{4}\right)$ meter/sec. Find the rate at which (i) his shadow is shortening. (ii) the tip of the shadow is moving.
- (15) A ladder 10 meter long is leaning against a vertical wall. If the bottom of the ladder is pulled horizontally away from the wall at the rate of 1.2 meters per second, find how fast the top of the ladder is sliding down the wall when the bottom is 6 meters away from the wall.
- (16) If water is poured into an inverted hollow cone whose semi-vertical angel is 30° , so that its depth (measured along the axis) increases at the rate of 1 cm/ sec. Find the rate at which the volume of water increasing when the depth is 2 cm.

2.2.1 Approximations

If $f(x)$ is a differentiable function of x , then its derivative at $x = a$ is given by

$$f'(a) = \lim_{h \rightarrow 0} \left[\frac{f(a+h) - f(a)}{h} \right]$$

Here we use \doteq sign for approximation.

For a sufficiently small h we have,

$$f'(a) \doteq \left[\frac{f(a+h) - f(a)}{h} \right]$$

i.e. $hf'(a) \doteq f(a+h) - f(a)$

$\therefore f(a+h) \doteq f(a) + hf'(a)$

This is the formula to find the approximate value of the function at $x = a + h$, when $f'(a)$ exists. Let us solve some problems by using this formula.

SOLVED EXAMPLES

Ex. 1 : Find the approximate value of $\sqrt{64.1}$.

Solution :

Let $f(x) = \sqrt{x}$... (I)

Differentiate *w. r. t. x*.

$$f'(x) = \frac{1}{2\sqrt{x}}$$
 ... (II)

Let $a = 64, h = 0.1$

For $x = a = 64$, from (I) we get

$$f(a) = f(64) = \sqrt{64} = 8$$
 ... (III)

For $x = a = 64$, from (II) we get

$$f'(a) = f'(64) = \frac{1}{2\sqrt{64}} = \frac{1}{16}$$

$\therefore f'(a) = 0.0625$... (IV)

We have, $f(a+h) \doteq f(a) + hf'(a)$

$$f(64 + 0.1) \doteq f(64) + (0.1) \cdot f'(64)$$

$$f(64.1) \doteq 8 + (0.1) \cdot (0.0625) \dots$$

[From (III) and (IV)]

$$\doteq 8 + 0.00625$$

$$\therefore f(64.1) = \sqrt{64.1} \doteq 8.00625$$

Ex. 2 : Find the approximate value of $(3.98)^3$.

Solution :

Let $f(x) = x^3$... (I)

Differentiate *w. r. t. x*.

$$f'(x) = 3x^2$$
 ... (II)

Let $a = 4, h = -0.02$

For $x = a = 4$, from (I) we get

$$f(a) = f(4) = (4)^3 = 64$$
 ... (III)

For $x = a = 4$, from (II) we get

$$f'(a) = f'(4) = 3(4)^2 = 48$$
 ... (IV)

We have, $f(a+h) \doteq f(a) + hf'(a)$

$$f[4 + (-0.02)] \doteq f(4) + (-0.02) \cdot f'(4)$$

$$f(3.98) \doteq 64 + (-0.02) \cdot (48) \dots$$

[From (III) and (IV)]

$$f(3.98) \doteq 64 - 0.96$$

$$\therefore f(3.98) = (3.98)^3 \doteq 63.04$$

Ex. 3 : Find the approximate value of $\sin(30^\circ 30')$. Given that $1^\circ = 0.0175^c$ and $\cos 30^\circ = 0.866$.

Solution : Let $f(x) = \sin x$... (I)

Differentiate w. r. t. x .

$$f'(x) = \cos x$$

$$\begin{aligned} \text{Now, } 30^\circ 30' &= 30^\circ + 30' = 30^\circ + \left(\frac{1}{2}\right)^\circ \\ &= \frac{\pi}{6} + \frac{0.1750^c}{2} \end{aligned}$$

$$30^\circ 30' = \frac{\pi}{6} + 0.00875 \quad \dots \text{ (II)}$$

$$\text{Let } a = \frac{\pi}{6}, h = 0.00875$$

For $x = a = \frac{\pi}{6}$, from (I) we get

$$f(a) = f\left(\frac{\pi}{6}\right) = \sin\left(\frac{\pi}{6}\right) = \frac{1}{2} = 0.5 \quad \dots \text{ (III)}$$

For $x = a = \frac{\pi}{6}$, from (II) we get

$$f'(a) = f'\left(\frac{\pi}{6}\right) = \cos\left(\frac{\pi}{6}\right) = 0.866 \quad \dots \text{ (IV)}$$

We have, $f(a + h) \doteq f(a) + hf'(a)$

$$f\left(\frac{\pi}{6} + 0.00875^c\right) \doteq f\left(\frac{\pi}{6}\right) + (0.00875) \cdot f'\left(\frac{\pi}{6}\right)$$

$$\begin{aligned} f(30^\circ 30') &\doteq 0.5 + (0.00875) \cdot (0.866) \dots \\ &\dots \text{ [From (III) and (IV)]} \end{aligned}$$

$$\doteq 0.5 + 0.075775$$

$$\therefore f(30^\circ 30') = \sin(30^\circ 30') \doteq 0.575775$$

Ex. 4 : Find the approximate value of $\tan^{-1}(0.99)$, Given that $\pi \doteq 3.1416$.

Solution : Let $f(x) = \tan^{-1} x$... (I)

Differentiate w. r. t. x .

$$f'(x) = \frac{1}{1+x^2} \quad \dots \text{ (II)}$$

$$\text{Let } a = 1, h = -0.01$$

For $x = a = 1$, from (I) we get

$$f(a) = f(1) = \tan^{-1}(1) = \frac{\pi}{4} \quad \dots \text{ (III)}$$

For $x = a = 1$, from (II) we get

$$f'(a) = f'(1) = \frac{1}{1+1^2} = 0.5 \quad \dots \text{ (IV)}$$

We have, $f(a + h) \doteq f(a) + hf'(a)$

$$f[(1) + (-0.01)] \doteq f(1) + (-0.01) \cdot f'(1)$$

$$\begin{aligned} f(0.99) &\doteq \frac{\pi}{4} - (0.01) \cdot (0.5) \dots \text{ [From} \\ &\text{(III) and (IV)]} \end{aligned}$$

$$\doteq \frac{\pi}{4} - 0.005$$

$$\doteq \frac{3.1416}{4} - 0.005$$

$$\doteq 0.7854 - 0.005 = 0.7804$$

$$\therefore f(0.99) = \tan^{-1}(0.99) \doteq 0.7804$$

Ex. 5 : Find the approximate value of $e^{1.005}$. Given that $e = 2.7183$.

Solution : Let $f(x) = e^x$... (I)

Differentiate w. r. t. x .

$$f'(x) = e^x \quad \dots \text{ (II)}$$

$$\text{Let } a = 1, h = 0.005$$

For $x = a = 1$, from (I) we get

$$f(a) = f(1) = e^1 = 2.7183 \quad \dots \text{ (III)}$$

For $x = a = 1$, from (II) we get

$$f'(a) = f'(1) = e^1 = 2.7183 \quad \dots \text{ (IV)}$$

We have, $f(a + h) \doteq f(a) + hf'(a)$

$$f(1 + 0.005) \doteq f(1) + (0.005) \cdot f'(1)$$

$$\begin{aligned} f(1.005) &\doteq 2.7183 + (0.005) (2.7183) \dots \\ &\dots \text{ [From (III) and (IV)]} \end{aligned}$$

$$f(1.005) \doteq 2.7183 + 0.0135915$$

$$\doteq 2.7318915$$

$$f(1.005) = e^{1.005} \doteq 2.73189$$

Ex. 6 : Find the approximate value of $\log_{10} (998)$. Given that $\log_{10} e = 0.4343$.

Solution : Let $f(x) = \log_{10} x = \frac{\log x}{\log 10}$
 $\therefore f(x) = (\log_{10} e) \cdot \log x \quad \dots (I)$
 Differentiate *w. r. t. x.*
 $f'(x) = \frac{\log_{10} e}{x} = \frac{0.4343}{x} \quad \dots (II)$
 Let $a = 1000, h = -2$
 For $x = a = 1000$, from (I) we get
 $f(a) = f(1000) = \log_{10} 1000$
 $\therefore f(a) = 3 \log_{10} 10 = 3 \quad \dots (III)$
 For $x = a = 1000$, from (II) we get
 $f'(a) = f'(1000) = \frac{0.4343}{1000}$
 $\therefore f'(a) = 0.0004343 \quad \dots (IV)$
 We have, $f(a + h) \doteq f(a) + hf'(a)$
 $f[1000 + (-2)] \doteq f(1000) + (-2)f'(1000)$
 $f(998) \doteq 3 - (2)(0.0004343) \dots$
 [From (III) and (IV)]
 $\doteq 3 - 0.0008686$
 $f(998) = \log (998) \doteq 2.9991314$

Ex. 7 : Find the approximate value of $f(x) = x^3 + 5x^2 - 2x + 3$ at $x = 1.98$.

Solution : Let $f(x) = x^3 + 5x^2 - 2x + 3 \quad \dots (I)$
 Differentiate *w. r. t. x.*
 $f'(x) = 3x^2 + 10x - 2 \quad \dots (II)$
 Let $a = 2, h = -0.02$
 For $x = a = 2$, from (I) we get
 $f(a) = f(2) = (2)^3 + 5(2)^2 - 2(2) + 3$
 $\therefore f(a) = 27 \quad \dots (III)$
 For $x = a = 2$, from (II) we get
 $f'(a) = f'(2) = 3(2)^2 + 10(2) - 2$
 $\therefore f'(a) = 30 \quad \dots (IV)$
 We have, $f(a + h) \doteq f(a) + hf'(a)$
 $f[(2) + (-0.02)] \doteq f(2) + (-0.02) \cdot f'(2)$
 $f(1.98) \doteq 27 - (0.02) \cdot (30) \dots$ [From (III) and (IV)]
 $\doteq 27 - 0.6$
 $f(1.98) \doteq 26.4$

EXERCISE 2.2

- (1) Find the approximate value of given functions, at required points.
 (i) $\sqrt{8.95}$ (ii) $\sqrt[3]{28}$ (iii) $\sqrt[5]{31.98}$
 (iv) $(3.97)^4$ (v) $(4.01)^3$
- (2) Find the approximate value of
 (i) $\sin (61^\circ)$ given that $1^\circ = 0.0174^c$, $\sqrt{3} = 1.732$
 (ii) $\sin (29^\circ 30')$ given that $1^\circ = 0.0175^c$, $\sqrt{3} = 1.732$
 (iii) $\cos (60^\circ 30')$ given that $1^\circ = 0.0175^c$, $\sqrt{3} = 1.732$
 (iv) $\tan (45^\circ 40')$ given that $1^\circ = 0.0175^c$.
- (3) Find the approximate value of
 (i) $\tan^{-1} (0.999)$ (ii) $\cot^{-1} (0.999)$
 (iii) $\tan^{-1} (1.001)$
- (4) Find the approximate value of
 (i) $e^{0.995}$ (ii) $e^{2.1}$ given that $e^2 = 7.389$
 (iii) $3^{2.01}$ given that $\log 3 = 1.0986$
- (5) Find the approximate value of
 (i) $\log_e (101)$ given that $\log_e 10 = 2.3026$
 (ii) $\log_e (9.01)$ given that $\log 3 = 1.0986$
 (iii) $\log_{10} (1016)$ given that $\log_{10} e = 0.4343$
- (6) Find the approximate value of
 (i) $f(x) = x^3 - 3x + 5$ at $x = 1.99$
 (ii) $f(x) = x^3 + 5x^2 - 7x + 10$ at $x = 1.12$

2.3.1 Rolle's Theorem or Rolle's Lemma :

If a real-valued function f is continuous on $[a, b]$, differentiable on the open interval (a, b) and $f(a) = f(b)$, then there exists at least one c in the open interval (a, b) such that $f'(c) = 0$.

Rolle's Theorem essentially states that any real-valued differentiable function that attains equal values at two distinct points on it, must have at least one stationary point somewhere in between them, that is, a point where the first derivative (the slope of the tangent line to the graph of the function) is zero.

Geometrical Significance :

Let $f(x)$ be a real valued function defined on $[a, b]$ and it is continuous on $[a, b]$. This means that we can draw the graph $f(x)$ between the values $x = a$ and $x = b$. Also $f(x)$ is differentiable on (a, b) which means the graph of $f(x)$ has a tangent




Fig. 2.3.1

at each point of (a, b) . Now the existence of real number $c \in (a, b)$ such that $f'(c) = 0$ shows that the tangent to the curve at $x = c$ has slope zero, that is, tangent is parallel to X-axis since $f(a) = f(b)$.

SOLVED EXAMPLES

Ex. 1 : Check whether conditions of Rolle's theorem are satisfied by the following functions.

- (i) $f(x) = 2x^3 - 5x^2 + 3x + 2, x \in \left[0, \frac{3}{2}\right]$
- (ii) $f(x) = x^2 - 2x + 3, x \in [1, 4]$

Solution :

(i) Given that $f(x) = 2x^3 - 5x^2 + 3x + 2 \dots (I)$

$f(x)$ is a polynomial which is continuous on $\left[0, \frac{3}{2}\right]$ and it is differentiable on $\left(0, \frac{3}{2}\right)$.

Let $a = 0$, and $b = \frac{3}{2}$,

For $x = a = 0$ from (I) we get,

$$f(a) = f(0) = 2(0)^3 - 5(0)^2 + 3(0) + 2 = 2$$

For $x = b = \left(\frac{3}{2}\right)$ from (I) we get,

$$f(b) = f\left(\frac{3}{2}\right) = 2\left(\frac{3}{2}\right)^3 - 5\left(\frac{3}{2}\right)^2 + 3\left(\frac{3}{2}\right) + 2 = \frac{54}{8} - \frac{45}{4} + \frac{9}{2} + 2$$

$$f(b) = f\left(\frac{3}{2}\right) = \frac{54 - 90 + 36}{8} + 2 = 2$$

So, here $f(a) = f(b)$ i.e. $f(0) = f\left(\frac{3}{2}\right) = 2$

Hence conditions of Rolle's Theorem are satisfied.



(ii) Given that $f(x) = x^2 - 2x + 3 \dots (I)$
 $f(x)$ is a polynomial which is continuous on $[1, 4]$ and it is differentiable on $(1, 4)$.

Let $a = 1$, and $b = 4$

For $x = a = 1$ from (I) we get,

$$f(a) = f(1) = (1)^2 - 2(1) + 3 = 2$$

For $x = b = 4$ from (I) we get,

$$f(b) = f(4) = (4)^2 - 2(4) + 3 = 11$$

So, here $f(a) \neq f(b)$ i.e. $f(1) \neq f(4)$

Hence conditions of Rolle's theorem are not satisfied.

Ex. 2 : Verify Rolle's theorem for the function
 $f(x) = x^2 - 4x + 10$ on $[0, 4]$.

Solution :

Given that $f(x) = x^2 - 4x + 10 \dots (I)$

$f(x)$ is a polynomial which is continuous on $[0, 4]$ and it is differentiable on $(0, 4)$.

Let $a = 0$, and $b = 4$

For $x = a = 0$ from (I) we get,

$$f(a) = f(0) = (0)^2 - 4(0) + 10 = 10$$

For $x = b = 4$ from (I) we get,

$$f(b) = f(4) = (4)^2 - 4(4) + 10 = 10$$

So, here $f(a) = f(b)$ i.e. $f(0) = f(4) = 10$

All the conditions of Rolle's theorem are satisfied.

To get the value of c , we should have

$$f'(c) = 0 \text{ for some } c \in (0, 4)$$

Differentiate (I) w. r. t. x .

$$f'(x) = 2x - 4 = 2(x - 4)$$

Now, for $x = c$,

$$f'(c) = 0 \Rightarrow 2(c - 2) = 0 \Rightarrow c = 2$$

Also $c = 2 \in (0, 4)$

Thus Rolle's theorem is verified.

Ex. 3 : Given an interval $[a, b]$ that satisfies hypothesis of Rolle's theorem for the function $f(x) = x^3 - 2x^2 + 3$. It is known that $a = 0$. Find the value of b .

Solution :

Given that $f(x) = x^3 - 2x^2 + 3 \dots (I)$

Let $g(x) = x^3 - 2x^2 = x^2(x - 2)$

From (I), $f(x) = g(x) + 3$

We see that $g(x)$ becomes zero for $x = 0$ and $x = 2$.

We observe that for $x = 0$,

$$f(0) = g(0) + 3 = 3$$

and for $x = 2$,

$$f(2) = g(2) + 3 = 3$$

\therefore We can write that $f(0) = f(2) = 3$

It is obvious that the function $f(x)$ is everywhere continuous and differentiable as a cubic polynomial. Consequently, it satisfies all the conditions of Rolle's theorem on the interval $[0, 2]$.

So $b = 2$.

Ex. 4 : Verify Rolle's theorem for the function $f(x) = e^x (\sin x - \cos x)$ on $\left[\frac{\pi}{4}, \frac{5\pi}{4}\right]$.

Solution : Given that, $f(x) = e^x (\sin x - \cos x) \dots (I)$

We know that e^x , $\sin x$ and $\cos x$ are continuous and differentiable on their domains. Therefore $f(x)$ is continuous and differentiable on $\left[\frac{\pi}{4}, \frac{5\pi}{4}\right]$ and $\left(\frac{\pi}{4}, \frac{5\pi}{4}\right)$ respectively.

Let $a = \frac{\pi}{4}$, and $b = \frac{5\pi}{4}$

For $x = a = \frac{\pi}{4}$ from (I) we get,

$$f(a) = f\left(\frac{\pi}{4}\right) = e^{\frac{\pi}{4}} \left[\sin\left(\frac{\pi}{4}\right) - \cos\left(\frac{\pi}{4}\right) \right] = e^{\frac{\pi}{4}} \left(\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}} \right) = 0$$

For $x = b = \left(\frac{5\pi}{4}\right)$ from (I) we get,

$$f(a) = f\left(\frac{5\pi}{4}\right) = e^{\frac{5\pi}{4}} \left[\sin\left(\frac{5\pi}{4}\right) - \cos\left(\frac{5\pi}{4}\right) \right] = e^{\frac{5\pi}{4}} \left(-\frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \right) = 0$$

$\therefore f(a) = f(b)$ i.e. $f\left(\frac{\pi}{4}\right) = f\left(\frac{5\pi}{4}\right)$.

All the conditions of Rolle's theorem are satisfied.

To get the value of c , we should have $f'(c) = 0$ for some $c \in \left[\frac{\pi}{4}, \frac{5\pi}{4}\right]$

Differentiate (I) w. r. t. x .

$$f'(x) = e^x (\cos x + \sin x) + (\sin x - \cos x) e^x = 2e^x \sin x$$

Now, for $x = c$, $f'(c) = 0 \Rightarrow 2e^c \sin c = 0$. As $e^c \neq 0$ for any $c \in R$

$$\sin c = 0 \Rightarrow c = 0, \pm \pi, \pm 2\pi, \pm 3\pi, \dots$$

It is clearly seen that $\pi \in \left[\frac{\pi}{4}, \frac{5\pi}{4}\right] \therefore c = \pi$


Thus Rolle's theorem is verified.

2.3.2 Lagrange's Mean Value Theorem (LMVT) :

If a real-valued function f is continuous on a closed $[a, b]$ and differentiable on the open interval (a, b) then there exists at least one c in the open interval (a, b) such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

Lagrange's mean value theorem states, that for any real-valued differentiable function which is continuous at the two end points, there is at least one point at which the tangent is parallel to the secant through its end points.



Geometrical Significance :

Draw the curve $y = f(x)$ (see Figure 2.3.2) and take the end points $A(a, f(a))$ and $B(b, f(b))$ on the curve, then

$$\text{Slope of the chord } AB = \frac{f(b) - f(a)}{b - a}$$

Since by statement of Lagrange's Mean Value.

Theorem
$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f'(c) = \text{Slope of the chord } AB.$$




Fig. 2.3.2

This shows that the tangent to the curve $y = f(x)$ at the point $x = c$ is parallel to the chord AB.

SOLVED EXAMPLES

Ex. 1 : Verify Lagrange's mean value theorem for the function $f(x) = \sqrt{x + 4}$ on the interval $[0, 5]$.

Solution : Given that $f(x) = \sqrt{x + 4}$... (I)

The function $f(x)$ is continuous on the closed interval $[0, 5]$ and differentiable on the open interval $(0, 5)$, so the LMVT is applicable to the function.

Differentiate (I) w. r. t. x .

$$f'(x) = \frac{1}{2\sqrt{x+4}} \quad \dots \text{(II)}$$

Let $a = 0$ and $b = 5$

$$\text{From (I), } f(a) = f(0) = \sqrt{0 + 4} = 2$$

$$f(b) = f(5) = \sqrt{5 + 4} = 3$$

Let $c \in (0, 5)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$\frac{1}{2\sqrt{c+4}} = \frac{3 - 2}{5 - 0} = \frac{1}{5}$$

$$\therefore \sqrt{c+4} = \frac{5}{2} \Rightarrow c+4 = \frac{25}{4} \therefore c = \frac{9}{4} \in (0, 5)$$

Thus Lagrange's Mean Value Theorem is verified.

Ex. 2 : Verify Lagrange's mean value theorem for the function $f(x) = x + \frac{1}{x}$ on the interval $[1, 3]$.

Solution : Given that $f(x) = x + \frac{1}{x}$... (I)

The function $f(x)$ is continuous on the closed interval $[1, 3]$ and differentiable on the open interval $(1, 3)$, so the LMVT is applicable to the function.

Differentiate (I) w. r. t. x .

$$f'(x) = 1 - \frac{1}{x^2} \quad \dots \text{(II)}$$

Let $a = 1$ and $b = 3$

$$\text{From (I), } f(a) = f(1) = 1 + \frac{1}{1} = 2$$

$$f(b) = f(3) = 3 + \frac{1}{3} = \frac{10}{3}$$

Let $c \in (1, 3)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$1 - \frac{1}{c^2} = \frac{\frac{10}{3} - 2}{3 - 1}$$

$$1 - \frac{1}{c^2} = \frac{\frac{4}{3}}{2} = \frac{2}{3}$$

$$\therefore c^2 = 3 \Rightarrow c = \pm \sqrt{3}$$

$$\therefore c = \sqrt{3} \in (1, 3) \text{ and } c = -\sqrt{3} \notin (1, 3)$$

EXERCISE 2.3

- (1) Check the validity of the Rolle's theorem for the following functions.
 - (i) $f(x) = x^2 - 4x + 3, x \in [1, 3]$
 - (ii) $f(x) = e^{-x} \sin x, x \in [0, \pi]$
 - (iii) $f(x) = 2x^2 - 5x + 3, x \in [1, 3]$
 - (iv) $f(x) = \sin x - \cos x + 3, x \in [0, 2\pi]$
 - (v) $f(x) = x^2$ if $0 \leq x \leq 2$
 $= 6 - x$ if $2 \leq x \leq 6$
 - (vi) $f(x) = x^{\frac{2}{3}}, x \in [-1, 1]$
- (2) Given an interval $[a, b]$ that satisfies hypothesis of Rolle's thorem for the function $f(x) = x^4 + x^2 - 2$. It is known that $a = -1$. Find the value of b .
- (3) Verify Rolle's theorem for the following functions.
 - (i) $f(x) = \sin x + \cos x + 7, x \in [0, 2\pi]$
 - (ii) $f(x) = \sin\left(\frac{x}{2}\right), x \in [0, 2\pi]$
 - (iii) $f(x) = x^2 - 5x + 9, x \in [1, 4]$
- (4) If Rolle's theorem holds for the function $f(x) = x^3 + px^2 + qx + 5, x \in [1, 3]$ with $c = 2 + \frac{1}{\sqrt{3}}$, find the values of p and q .
- (5) Rolle's theorem holds for the function $f(x) = (x - 2) \log x, x \in [1, 2]$, show that the equation $x \log x = 2 - x$ is satisfied by at least one value of x in $(1, 2)$.
- (6) The function $f(x) = x(x + 3)e^{-\frac{x}{2}}$ satisfies all the conditions of Rolle's theorem on $[-3, 0]$. Find the value of c such that $f'(c) = 0$.
- (7) Verify Lagrange's mean value theorem for the following functions.
 - (i) $f(x) = \log x$, on $[1, e]$
 - (ii) $f(x) = (x - 1)(x - 2)(x - 3)$ on $[0, 4]$
 - (iii) $f(x) = x^2 - 3x - 1, x \in \left[-\frac{11}{7}, \frac{13}{7}\right]$
 - (iv) $f(x) = 2x - x^2, x \in [0, 1]$
 - (v) $f(x) = \frac{x - 1}{x - 3}$ on $[4, 5]$

2.4.1 Increasing and decreasing functions :

Increasing functions :

Definition : A function f is said to be a monotonically (or strictly) increasing function on an interval (a, b) if for any $x_1, x_2 \in (a, b)$ with if $x_1 < x_2$, we have $f(x_1) < f(x_2)$.

Consider an increasing function $y = f(x)$ in (a, b) . Let $h > 0$ be a small increment in x then,

$$x < x + h \quad [x = x_1, x + h = x_2]$$

$$f(x) < f(x + h) \quad [f(x_1) < f(x_2)]$$



$$\therefore f(x + h) > f(x)$$

$$\therefore f(x + h) - f(x) > 0$$

$$\therefore \frac{f(x + h) - f(x)}{h} > 0$$

$$\therefore \lim_{h \rightarrow 0} \left[\frac{f(x + h) - f(x)}{h} \right] \geq 0$$

$$\therefore f'(x) \geq 0$$



If $f'(a) > 0$, then in a small δ -neighborhood of a i.e. $(a - \delta, a + \delta)$, we have f strictly increasing if

$$\frac{f(a+h) - f(a)}{h} > 0 \quad \text{for } |h| < \delta$$

Hence if $0 < h < \delta$, $f(a+h) - f(a) > 0$ and $f(a-h) - f(a) < 0$


Thus for $0 < h < \delta$, $f(a-h) < f(a) < f(a+h)$

Decreasing functions :

Definition : A function f is said to be a monotonically (strictly) decreasing function on an interval (a, b) if for any $x_1, x_2 \in (a, b)$ with $x_1 < x_2$, we have $f(x_1) > f(x_2)$.

Consider a decreasing function $y = f(x)$ in (a, b) . Let $h > 0$ be a small increment in x then,

$$\begin{aligned} x+h &> x && [x = x_1, x+h = x_2] \\ f(x) &< f(x+h) && [f(x_1) < f(x_2)] \\ \therefore f(x+h) &< f(x) \\ \therefore f(x+h) - f(x) &< 0 \\ \therefore \frac{f(x+h) - f(x)}{h} &< 0 \\ \therefore \lim_{h \rightarrow 0} \left[\frac{f(x+h) - f(x)}{h} \right] &\leq 0 \\ \therefore f'(x) &\leq 0 \end{aligned}$$



If $f'(a) < 0$, then in a small δ -neighborhood of a i.e. $(a - \delta, a + \delta)$, we have f strictly decreasing

because
$$\frac{f(a+h) - f(a)}{h} < 0 \quad \text{for } |h| < \delta$$

Hence for $0 < h < \delta$, $f(a-h) > f(a) > f(a+h)$

Note : Whenever $f'(x) = 0$, at that point the tangent is parallel to X-axis, we cannot deduce that whether $f(x)$ is increasing or decreasing at that point.



SOLVED EXAMPLES

Ex. 1: Show that the function $f(x) = x^3 + 10x + 7$ for $x \in \mathbb{R}$ is strictly increasing.

Solution : Given that $f(x) = x^3 + 10x + 7$

Differentiate w. r. t. x .

$$f'(x) = 3x^2 + 10$$

Here, $3x^2 \geq 0$ for all $x \in \mathbb{R}$ and $10 > 0$.

$$\therefore 3x^2 + 10 > 0 \Rightarrow f'(x) > 0$$

Thus $f(x)$ is a strictly increasing function.

Ex. 2: Test whether the function

$f(x) = x^3 + 6x^2 + 12x - 5$ is increasing or decreasing for all $x \in \mathbb{R}$.

Solution : Given that $f(x) = x^3 + 6x^2 + 12x - 5$

Differentiate w. r. t. x .


$$f'(x) = 3x^2 + 12x + 12 = 3(x^2 + 4x + 4)$$

$$f'(x) = 3(x+2)^2$$

$3(x+2)^2$ is always positive for $x \neq -2$

$$\therefore f'(x) \geq 0 \text{ for all } x \in \mathbb{R}$$

Hence $f(x)$ is an increasing function for all $x \in \mathbb{R}$.



Ex. 3: Find the values of x , for which the function $f(x) = x^3 + 12x^2 + 36x + 6$ is (i) monotonically increasing. (ii) monotonically decreasing.

Solution : Given that $f(x) = x^3 + 12x^2 + 36x + 6$

Differentiate w. r. t. x .

$$f'(x) = 3x^2 + 24x + 36$$

$$= 3(x^2 + 8x + 12)$$

$$f'(x) = 3(x + 2)(x + 6)$$

(i) $f(x)$ is monotonically increasing if $f'(x) > 0$

i.e. $3(x + 2)(x + 6) > 0$, $(x + 2)(x + 6) > 0$

then either $(x + 2) < 0$ and $(x + 6) < 0$ or $(x + 2) > 0$ and $(x + 6) > 0$

Case (I) : $x + 2 < 0$ and $x + 6 < 0$

$$x < -2 \text{ and } x < -6$$

Thus for every $x < -6$, $(x + 2)(x + 6) > 0$, hence f is monotonically increasing.

Case (II) : $x + 2 > 0$ and $x + 6 > 0$

$$x > -2 \text{ and } x > -6$$

Thus for every $x > -2$, $(x + 2)(x + 6) > 0$ and f is monotonically increasing.

\therefore From Case (I) and Case (II), $f(x)$ is monotonically increasing if and only if $x < -6$ or $x > -2$.

Hence, $x \in (\infty, -6)$ or $x \in (-2, \infty) \Rightarrow f$ is monotonically increasing.

(ii) $f(x)$ is said to be monotonically decreasing if $f'(x) < 0$

i.e. $3(x + 2)(x + 6) < 0$, $(x + 2)(x + 6) < 0$

then either $(x + 2) < 0$ and $(x + 6) > 0$ or $(x + 2) > 0$ and $(x + 6) < 0$

Case (I) : $x + 2 < 0$ and $x + 6 > 0$


$$x < -2 \text{ and } x > -6$$

Thus for $x \in (-6, -2)$, f is monotonically decreasing.

Case (II) : $x + 2 > 0$ and $x + 6 < 0$

$$x > -2 \text{ and } x < -6$$

\therefore This case does not arise. . . . [check. why ?]



2.4.2 Maxima and Minima :

Maxima of a function $f(x)$: A function $f(x)$ is said to have a maxima at $x = c$ if the value of the function at $x = c$ is greater than any other value of $f(x)$ in a δ -neighborhood of c . That is for a small $\delta > 0$ and for $x \in (c - \delta, c + \delta)$ we have $f(c) > f(x)$. The value $f(c)$ is called a Maxima of $f(x)$. Thus the function $f(x)$ will have maxima at $x = c$ if $f(x)$ is increasing in $c - \delta < x < c$ and decreasing in $c < x < c + \delta$.

Minima of a function $f(x)$: A function $f(x)$ is said to have a minima at $x = c$ if the value of the function at $x = c$ is less than any other value of $f(x)$ in a δ -neighborhood of c . That is for a small $\delta > 0$ and for $x \in (c - \delta, c + \delta)$ we have $f(c) < f(x)$. The value $f(c)$ is called a Minima of $f(x)$. Thus the function $f(x)$ will have minima at $x = c$ if $f(x)$ is decreasing in $c - \delta < x < c$ and increasing in $c < x < c + \delta$.

If $f'(c) = 0$ then at $x = c$ the function is neither increasing nor decreasing, such a point on the curve is called **turning point** or **stationary point** of the function. Any point at which the tangent to the graph is horizontal is a turning point. We can locate the turn points by looking for points at which $\frac{dy}{dx} = 0$. At these points if the function has Maxima or Minima then these are called extreme values of the function.

Note : The maxima and the minima of a function are not necessarily the greatest and the least values of the function in the whole domain. Actually these are the greatest and the least values of the function in a small interval. Hence the maxima or the minima defined above are known as **local (or relative) maximum** and **the local (or relative) minimum** of the function $f(x)$.

To find the extreme values of the function let us use following tests.

2.4.3 First derivative test :

A function $f(x)$ has a maxima at $x = c$ if

- (i) $f'(c) = 0$
- (ii) $f'(c - h) > 0$ [$f(x)$ is increasing for values of $x < c$]
- (iii) $f'(c + h) < 0$ [$f(x)$ is decreasing for values of $x > c$]

where h is a small positive number.


A function $f(x)$ has a minima at $x = c$ if

- (i) $f'(c) = 0$
- (ii) $f'(c - h) < 0$ [$f(x)$ is decreasing for values of $x < c$]
- (iii) $f'(c + h) > 0$ [$f(x)$ is increasing for values of $x > c$]

where h is a small positive number.

Note : If $f'(c) = 0$ and $f'(c - h) > 0, f'(c + h) > 0$ or $f'(c - h) < 0, f'(c + h) < 0$ then $f(c)$ is neither maxima nor minima. In such a case $x = c$ is called a **point of inflexion**. e.g. $f(x) = x^3, f(x) = x^5$ in $[-2, 2]$.





SOLVED EXAMPLES

Ex. 1: Find the local maxima or local minima of $f(x) = x^3 - 3x$.

Solution : Given that $f(x) = x^3 - 3x$. . . (I)

Differentiate (I) w. r. t. x .

$$f'(x) = 3x^2 - 3 = 3(x^2 - 1) \quad \dots \text{(II)}$$

For extreme values, $f'(x) = 0$

$$3x^2 - 3 = 0 \quad \text{i.e. } 3(x^2 - 1) = 0$$

$$\text{i.e. } x^2 - 1 = 0 \Rightarrow 0 \Rightarrow x^2 = 1 \Rightarrow x = \pm 1$$

The turning points are $x = 1$ and $x = -1$

Let's consider the turning point, $x = 1$

Let $x = 1 - h$ for a small, $h > 0$, from (II) we get,

$$f'(1 - h) = 3[(1 - h)^2 - 1] = 3(1 - 2h + h^2 - 1) = 3h(h - 2)$$

$$\therefore f'(1 - h) < 0 \dots [\text{since, } h > 0, h - 2 < 0]$$

$$\therefore f'(x) \text{ for } x = 1 - h \Rightarrow f(x) \text{ is decreasing for, } x > 1.$$

Now for $x = 1 + h$ for a small, $h > 0$, from (II) we get,

$$f'(1 + h) = 3[(1 + h)^2 - 1] = 3(1 + 2h + h^2 - 1) = 3(h^2 + 2h)$$

$$\therefore f'(1 + h) > 0 \dots [\text{since, } h > 0, h^2 + 2h > 0]$$

$$\therefore f'(x) < 0 \text{ for } x = 1 + h \Rightarrow f(x) \text{ is increasing for, } x < 1.$$

$$\therefore f'(x) < 0 \text{ for } 1 - h < x < 1$$

$$\therefore f'(x) > 0 \text{ for } 1 < x < 1 + h.$$

$$\therefore x = 1 \text{ is the point of local minima.}$$

Minima of $f(x)$, is $f(1) = 1^3 - 3(1) = -2$

Now, let's consider the turning point, $x = -1$

Let $x = -1 - h$ for a small, $h > 0$, from (II) we get,

$$\therefore f'(-1 - h) = 3[(-1 - h)^2 - 1] = 3(1 + 2h + h^2 - 1) = 3(h^2 + 2h)$$

$$\therefore f'(-1 - h) > 0 \dots [\text{since, } h > 0, h^2 + 2h > 0]$$

$$\therefore f'(x) > 0 \text{ for } x = -1 - h \Rightarrow f(x) \text{ is increasing for, } x < -1.$$

Now for $x = -1 + h$ for a small, $h > 0$, from (II) we get,

$$\therefore f'(-1 + h) = 3[(-1 + h)^2 - 1] = 3(1 - 2h + h^2 - 1) = -3h(2 - h)$$

$$\therefore f'(-1 + h) < 0 \dots [\text{since, } h > 0, 2 - h > 0]$$

$$\therefore f'(x) < 0 \text{ for } x = -1 + h \Rightarrow f(x) \text{ is decreasing for, } x > -1.$$

$$\therefore f'(x) > 0 \text{ for } -1 - h < x < -1$$

$$\therefore f'(x) > 0 \text{ for } -1 < x < -1 + h.$$

$$\therefore x = -1 \text{ is the point of local maxima.}$$

Maxima of $f(x)$, is $f(-1) = (-1)^3 - 3(-1) = -1 + 3 = 2$

Hence, Maxima of $f(x) = 2$ and Minima of $f(x) = -2$





2.4.4 Second derivative test :

A function $f(x)$ has a maxima at $x = c$ if $f'(c) = 0$ and $f''(c) < 0$

A function $f(x)$ has a minima at $x = c$ if $f'(c) = 0$ and $f''(c) > 0$

Note : If $f''(c) = 0$ then second derivative test fails so, you may try using first derivative test.



Maxima at A : Consider the slopes of the tangents (See Fig 2.4.4a) Slope of L_1 is +ve, slope of $L_2 = 0$ and slope of L_3 is -ve. Thus the slope is seen to be decreasing if there is a maximum at A.

Minima at A : Consider the slopes of the tangents (See Fig 2.4.4b) slope of L_1 is -ve, slope of $L_2 = 0$ and slope of L_3 is +ve. Thus the slope is seen to be increasing if there is a minima at A.

SOLVED EXAMPLES

Ex. 1 : Find the local maximum and local minimum value of $f(x) = x^3 - 3x^2 - 24x + 5$.

Solution : Given that $f(x) = x^3 - 3x^2 - 24x + 5 \dots$ (I)

Differentiate (I) w. r. t. x .

$$f'(x) = 3x^2 - 6x - 24 \dots$$
 (II)

For extreme values, $f'(x) = 0$

$$3x^2 - 6x - 24 \text{ i.e. } 3(x^2 - 2x - 8) = 0$$

$$\text{i.e. } x^2 - 2x - 8 = 0 \text{ i.e. } (x + 2)(x - 4) = 0$$

$$\Rightarrow x + 2 = 0 \text{ or } x - 4 = 0 \Rightarrow x = -2 \text{ or } x = 4$$

The stationary points are $x = -2$ and $x = 4$.

Differentiate (II) w. r. t. x .

$$f''(x) = 6x - 6 \dots$$
 (III)

For $x = -2$, from (III) we get,

$$f''(-2) = 6(-2) - 6 = -18 < 0$$

\therefore At $x = -2$, $f(x)$ has a maximum value.

For maximum of $f(x)$, put $x = -2$ in (I)

$$f(-2) = (-2)^3 - 3(-2)^2 - 24(-2) + 5 = 33.$$

For $x = 4$, from (III) we get

$$f''(4) = 6(4) - 6 = 18 > 0$$

\therefore At $x = 4$, $f(x)$ has a minimum value.

For minima of $f(x)$, put $x = 4$ in (I)

$$f(4) = (4)^3 - 3(4)^2 - 24(4) + 5 = -75$$

\therefore Local maximum of $f(x)$ is 33 when $x = -2$ and

Local minimum of $f(x)$ is -75 when $x = 4$.



Ex. 2: A wire of length 120 cm is bent in the form of a rectangle. Find its dimensions if the area of the rectangle is maximum.

Solution : Let x cm and y cm be the length and the breadth of the rectangle. Perimeter of rectangle = 120 cm.

$$\therefore 2(x + y) = 120 \text{ so, } x + y = 60$$

$$\therefore y = 60 - x \quad \dots \text{ (I)}$$

Let A be the area of the rectangle

$$\therefore A = xy = x(60 - x) = 60x - x^2 \dots \text{ [From (I)]}$$

Differentiate *w. r. t. x.*

$$\frac{dA}{dx} = 60 - 2x \quad \dots \text{ (II)}$$

$$\text{For maximum area } \frac{dA}{dx} = 0$$

$$\text{i.e. } 60 - 2x = 0 \Rightarrow x = 30$$

Differentiate (II) *w. r. t. x.*

$$\frac{d^2A}{dx^2} = -2 \quad \dots \text{ (III)}$$

For, $x = 30$ from (III) we get,

$$\left(\frac{d^2A}{dx^2}\right)_{x=30} = -2 < 0$$

When, $x = 30$, Area of the rectangle is maximum.


$$\text{Put } x = 30 \text{ in (I) we get } y = 60 - 30 = 30$$

\therefore Area of the rectangle is maximum if length = breadth = 30 cm.

Ex. 3: A Rectangular sheet of paper has it area 24 sq. meters. The margin at the top and the bottom are 75 cm each and at the sides 50 cm each. What are the dimensions of the paper, if the area of the printed space is maximum?

Solution : Let x m and y m be the width and the length of the rectangular sheet of paper respectively. Area of the paper = 24 sq. m.

$$\therefore xy = 24 \Rightarrow y = \frac{24}{x} \quad \dots \text{ (I)}$$



After leaving the margins, length of the printing space is $(x - 1)$ m and breadth of the printing space is $(y - 1.5)$ m.

Let A be the area of the printing space

$$A = (x - 1)(y - 1.5) = (x - 1)\left(\frac{24}{x} - 1.5\right)$$

$$= 24 - 1.5x - \frac{24}{x} + 1.5 \dots \text{ [From (I)]}$$

$$A = 25.5 - 1.5x - \frac{24}{x} \quad \dots \text{ (II)}$$

Differentiate *w. r. t. x.*

$$\frac{dA}{dx} = -1.5 + \frac{24}{x^2} \quad \dots \text{ (III)}$$

$$\text{For maximum printing space } \frac{dA}{dx} = 0$$

$$\text{i.e. } -1.5x + \frac{24}{x^2} = 0 \Rightarrow 1.5x^2 = 24 \Rightarrow x = \pm 4, x \neq -4$$

$$\therefore x = 4$$

Differentiate (III) *w. r. t. x.*

$$\frac{d^2A}{dx^2} = -\frac{48}{x^3} \dots \text{ (IV)}$$

For, $x = 4$, from (IV) we get,

$$\left(\frac{d^2A}{dx^2}\right)_{x=4} = -\frac{48}{(4)^3} < 0$$

When, $x = 4$ Area of the rectangular printing space is maximum.

$$\text{Put } x = 4 \text{ in (I) we get } y = \frac{24}{4} = 6$$

\therefore Area of the printing space is maximum when width printing space = 4 m. and length of the printing space = 6 m.



Ex. 4: An open box is to be cut out of piece of square card board of side 18 cm by cutting of equal squares from the corners and turning up the sides. Find the maximum volume of the box.

Solution : Let the side of each of the small squares cut be x cm, so that each side of the box to be made is $(18 - 2x)$ cm. and height x cm.




Fig. 2.4.6

Let V be the volume of the box.

$$V = \text{Area of the base} \times \text{Height}$$

$$= (18 - 2x)^2 x = (324 - 72x + 4x^2) x$$

$$V = 4x^3 - 72x^2 + 324x \quad \dots \text{(I)}$$

Differentiate w. r. t. x

$$\frac{dV}{dx} = 12x^2 - 144x + 324 \quad \dots \text{(II)}$$

For maximum volume $\frac{dV}{dx} = 0$

i.e. $12x^2 - 144x + 324 = 0 \Rightarrow x^2 - 12x + 27 = 0$
 $(x - 3)(x - 9) = 0 \Rightarrow x - 3 = 0$ or $x - 9 = 0$

$\therefore x = 3$ or $x = 9$, but $x \neq 9 \therefore x = 3$

Differentiate (II) w. r. t. x

$$\frac{d^2V}{dx^2} = 24x - 144 \quad \dots \text{(III)}$$

For, $x = 3$ from (III) we get,

$$\left(\frac{d^2V}{dx^2}\right)_{x=3} = 24(3) - 144 = -72 < 0$$

Volume of the box is maximum when $x = 3$.

Maximum volume of the box

$$= (18 - 6)^2 (3) = 432 \text{ c.c.}$$

Ex. 5: Two sides of a triangle are given, find the angle between them such that the area of the triangle is maximum.

Solution : Let ABC be a triangle. Let the given sides be $AB = c$ and $AC = b$.




Fig. 2.4.7

Let Δ be the area of the triangle.

$$\Delta = \frac{1}{2} bc \sin A \quad \dots \text{(I)}$$

Differentiate w. r. t. A .

$$\frac{d\Delta}{dA} = \frac{bc}{2} \cos A \quad \dots \text{(II)}$$

For maximum area $\frac{d\Delta}{dA} = 0$

i.e. $\frac{bc}{2} \cos A = 0 \Rightarrow \cos A = 0 \Rightarrow A = \frac{\pi}{2}$

Differentiate (II) w. r. t. A .

$$\frac{d^2\Delta}{dA^2} = -\frac{bc}{2} \sin A \quad \dots \text{(III)}$$

For, $A = \frac{\pi}{2}$ from (III) we get,

$$\left(\frac{d^2\Delta}{dA^2}\right)_{A=\frac{\pi}{2}} = -\frac{bc}{2} \sin\left(\frac{\pi}{2}\right) = \frac{bc}{a} < 0$$

When, $A = \frac{\pi}{2}$ Area of the triangle is maximum.

Hence, the area of the triangle is maximum when the angle between the given sides $\frac{\pi}{2}$.

Note : $\sin A$ is maximum (=1), when $A = \frac{\pi}{2}$



Ex. 6: The slant side of a right circular cone is l . Show that the semi-vertical angle of the cone of maximum volume is $\tan^{-1}(\sqrt{2})$.

Solution : Let x be the height of the cone and r be the radius of the base.

$$\text{So, } r^2 = l^2 - x^2 \quad \dots \text{ (I)}$$

Let V be the volume of the cone.

$$V = \frac{1}{3} \pi r^2 x = \frac{\pi}{3} (l^2 - x^2) x$$

$$\therefore V = \frac{\pi}{3} (l^2 x - x^3)$$

Differentiate w. r. t. x

$$\frac{dV}{dx} = \frac{\pi}{3} (l^2 - 3x^2) \quad \dots \text{ (II)}$$

$$\text{For maximum volume } \frac{dV}{dx} = 0$$

$$\text{i.e. } \frac{\pi}{3} (l^2 - 3x^2) = 0 \Rightarrow x^2 = \frac{l^2}{3}$$

$$x = \pm \frac{l}{\sqrt{3}} \Rightarrow x = \frac{l}{\sqrt{3}} \text{ or } x = -\frac{l}{\sqrt{3}} \text{ is the stationary point but, } x \neq -\frac{l}{\sqrt{3}} \therefore x = \frac{l}{\sqrt{3}}$$

Differentiate (II) w. r. t. x

$$\left(\frac{d^2V}{dx^2}\right) = -2\pi x \quad \dots \text{ (III)}$$

For, $x = \frac{l}{\sqrt{3}}$ from (III) we get,

$$\left(\frac{d^2V}{dx^2}\right)_{x=\frac{l}{\sqrt{3}}} = -\frac{2\pi l}{\sqrt{3}} < 0$$

Volume of the cone is maximum when height of the cone is $x = \frac{l}{\sqrt{3}}$.

$$\text{Put } x = \frac{l}{\sqrt{3}} \text{ in (I) we get, } r = \sqrt{l^2 - \left(\frac{l}{\sqrt{3}}\right)^2} = \frac{l\sqrt{2}}{\sqrt{3}}$$

Let α be the semi-vertical angle.

$$\text{Then } \tan \alpha = \frac{r}{x} = \frac{\frac{l\sqrt{2}}{\sqrt{3}}}{\frac{l}{\sqrt{3}}} = \sqrt{2}$$


$$\therefore \alpha = \tan^{-1}(\sqrt{2})$$

Ex. 7: Find the height of a covered box of fixed volume so that the total surface area of the box is minimum whose base is a rectangle with one side three times as long as the other.

Solution : Given that, box has a rectangular base with one side three times as long as other.

Let x and $3x$ be the sides of the rectangular base.

Let h be the height of the box and V be its volume.



$V = (x)(3x)(h) = 3x^2h \dots$ [Observe that V is constant]

Differentiate *w. r. t. x.*

$$\frac{dV}{dx} = 3x^2 \frac{dh}{dx} + h \frac{d}{dx}(3x^2)$$

$$\therefore 3x^2 \frac{dh}{dx} + 6xh = 0 \Rightarrow \frac{dh}{dx} = -\frac{2h}{x} \dots \text{(I)}$$

Let S be the surface area of the box.

$$\therefore S = (2 \times 3x^2) + (2 \times 3xh) + (2 \times xh) = 6x^2 + 8xh$$

Differentiate *w. r. t. x.*

$$\frac{dS}{dx} = 12x + 8 \left(x \frac{dh}{dx} + h \frac{d}{dx}(x) \right)$$

$$\frac{dS}{dx} = 12x + 8 \left[x \left(-\frac{2h}{x} \right) + h \right] \dots \text{[from (I)]}$$

$$= 12x + 8(-2h + h)$$

$$\therefore \frac{dS}{dx} = 12x - 8h \dots \text{(II)}$$

For minimum surface area

$$\frac{dS}{dx} = 0 \Rightarrow 12x - 8h = 0 \Rightarrow h = \frac{3x}{2}$$

Differentiate (II) *w. r. t. x.*

$$\left(\frac{d^2S}{dx^2} \right) = 12 - 8 \frac{dh}{dx} = 12 - 8 \left(-\frac{2h}{x} \right) = 12 + \frac{16h}{x} \dots \text{(III)} \dots \text{[from (I)]}$$

Both x and h are positive, from (III) we get,

$$\left(\frac{d^2S}{dx^2} \right) = 12 + \frac{16h}{x} > 0$$

Surface area of the box is minimum if height = $\frac{3}{2} \times$ shorter side of base.

EXERCISE 2.4

(1) Test whether the following functions are increasing or decreasing.

(i) $f(x) = x^3 - 6x^2 + 12x - 16, x \in R$

(ii) $f(x) = 2 - 3x + 3x^2 - x^3, x \in R$

(iii) $f(x) = x - \frac{1}{x}, x \in R$ and $x \neq 0$

(2) Find the values of x for which the following functions are strictly increasing -

(i) $f(x) = 2x^3 - 3x^2 - 12x + 6$

(ii) $f(x) = 3 + 3x - 3x^2 + x^3$

(iii) $f(x) = x^3 - 6x^2 - 36x + 7$

(3) Find the values of x for which the following functions are strictly decreasing -

(i) $f(x) = 2x^3 - 3x^2 - 12x + 6$

(ii) $f(x) = x + \frac{25}{x}$

(iii) $f(x) = x^3 - 9x^2 + 24x + 12$



- (4) Find the values of x for which the function $f(x) = x^3 - 12x^2 - 144x + 13$
 (a) Increasing (b) Decreasing
- (5) Find the values of x for which $f(x) = 2x^3 - 15x^2 - 144x - 7$ is
 (a) strictly increasing
 (b) strictly decreasing
- (6) Find the values of x for which $f(x) = \frac{x}{x^2 + 1}$ is
 (a) strictly increasing
 (b) strictly decreasing
- (7) Show that $f(x) = 3x + \frac{1}{3x}$ increasing in $\left(\frac{1}{3}, 1\right)$ and decreasing in $\left(\frac{1}{9}, \frac{1}{3}\right)$.
- (8) Show that $f(x) = x - \cos x$ is increasing for all x .
- (9) Find the maximum and minimum of the following functions -
 (i) $y = 5x^3 + 2x^2 - 3x$
 (ii) $f(x) = 2x^3 - 21x^2 + 36x - 20$
 (iii) $f(x) = x^3 - 9x^2 + 24x$
 (iv) $f(x) = x^2 + \frac{16}{x^2}$
 (v) $f(x) = x \log x$ (vi) $f(x) = \frac{\log x}{x}$
- (10) Divide the number 30 in to two parts such that their product is maximum.
- (11) Divide that number 20 in to two parts such that sum of their squares is minimum.
- (12) A wire of length 36 meters is bent in the form of a rectangle. Find its dimensions if the area of the rectangle is maximum.
- (13) A ball is thrown in the air. Its height at any time t is given by $h = 3 + 14t - 5t^2$. Find the maximum height it can reach.
- (14) Find the largest size of a rectangle that can be inscribed in a semi circle of radius 1 unit, So that two vertices lie on the diameter.
- (15) An open cylindrical tank whose base is a circle is to be constructed of metal sheet so as to contain a volume of πa^3 cu. cm of water. Find the dimensions so that sheet required is minimum.
- (16) The perimeter of a triangle is 10 cm. If one of the side is 4 cm. What are the other two sides of the triangle for its maximum area ?
- (17) A box with a square base is to have an open top. The surface area of the box is 192 sq.cm. What should be its dimensions in order that the volume is largest ?
- (18) The profit function $P(x)$ of a firm, selling x items per day is given by $P(x) = (150 - x)x - 1625$. Find the number of items the firm should manufacture to get maximum profit. Find the maximum profit.
- (19) Find two numbers whose sum is 15 and when the square of one multiplied by the cube of the other is maximum.
- (20) Show that among rectangles of given area, the square has the least perimeter.
- (21) Show that the height of a closed right circular cylinder of a given volume and least surface area is equal to its diameter.
- (22) Find the volume of the largest cylinder that can be inscribed in a sphere of radius ' r ' cm.
- (23) Show that $y = \log(1+x) - \frac{2x}{2+x}$, $x > -1$ is an increasing function on its domain.
- (24) Prove that $y = \frac{4 \sin \theta}{2 + \cos \theta} - \theta$ is an increasing function of $\theta \in \left[0, \frac{\pi}{2}\right]$.



Let us Remember

- Equations of tangent and Normal at $P(x_1, y_1)$ respectively are given by

$$y - y_1 = m(x - x_1) \text{ where } m = \left[\frac{dy}{dx} \right]_{(x_1, y_1)}$$

$$y - y_1 = m'(x - x_1) \text{ where } m' = -\frac{1}{\left[\frac{dy}{dx} \right]_{(x_1, y_1)}}, \text{ if } \left[\frac{dy}{dx} \right]_{(x_1, y_1)} \neq 0$$

- Approximate value of the function $f(x)$ at $x = a + h$ is given by $f(a + h) \doteq f(a) + hf'(a)$
- Rolle's theorem :** If real-valued function f is continuous on a closed $[a, b]$, differentiable on the open interval (a, b) and $f(a) = f(b)$, then there exists at least one c in the open interval (a, b) such that $f'(c) = 0$.
- Lagrange's Mean Value Theorem (LMVT) :** If a real-valued function f is continuous on a closed $[a, b]$ and differentiable on the open interval (a, b) then there exists at least one c in the open interval (a, b) such that $f'(c) = \frac{f(b) - f(a)}{b - a}$
- Increasing and decreasing functions :**
 - A function f is monotonically increasing if $f'(x) > 0$.
 - A function f is monotonically decreasing if $f'(x) < 0$.
 - A function f is increasing if $f'(x) \geq 0$.
 - A function f is decreasing if $f'(x) \leq 0$.
- (i) First Derivative test :**

A function $f(x)$ has a maxima at $x = c$ if

 - $f'(c) = 0$
 - $f'(c - h) > 0$ [$f(x)$ is increasing for values of $x < c$]
 - $f'(c + h) < 0$ [$f(x)$ is decreasing for values of $x > c$]

where h is a small positive number.

A function $f(x)$ has a minima at $x = c$ if

 - $f'(c) = 0$
 - $f'(c - h) < 0$ [$f(x)$ is decreasing for values of $x < c$]
 - $f'(c + h) > 0$ [$f(x)$ is increasing for values of $x > c$]

where h is a small positive number.
- (ii) Second Derivative test :**

A function $f(x)$ has a maxima at $x = c$ if $f'(c) = 0$ and $f''(c) < 0$.

A function $f(x)$ has a minimum at $x = c$ if $f'(c) = 0$ and $f''(c) > 0$.

MISCELLANEOUS EXERCISE 2

(I) Choose the correct option from the given alternatives :


- (1) If the function $f(x) = ax^3 + bx^2 + 11x - 6$ satisfies conditions of Rolle's theorem in $[1, 3]$ and $f'\left(2 + \frac{1}{\sqrt{3}}\right) = 0$, then values of a and b are respectively.
 (A) 1, -6 (B) -2, 1 (C) -1, -6 (D) -1, 6
- (2) If $f(x) = \frac{x^2 - 1}{x^2 + 1}$, for every real x , then the minimum value of f is -
 (A) 1 (B) 0 (C) -1 (D) 2
- (3) A ladder 5 m in length is resting against vertical wall. The bottom of the ladder is pulled along the ground away from the wall at the rate of 1.5 m/ sec. The length of the higher point of ladder when the foot of the ladder is 4.0 m away from the wall decreases at the rate of
 (A) 1 (B) 2 (C) 2.5 (D) 3
- (4) Let $f(x)$ and $g(x)$ be differentiable for $0 < x < 1$ such $f(0) = 0, g(0) = 0, f(1) = 6$. Let there exist a real number c in $(0, 1)$ such that $f'(c) = 2g'(c)$, then the value of $g(1)$ must be
 (A) 1 (B) 3 (C) 2.5 (D) -1
- (5) Let $f(x) = x^3 - 6x^2 + 9x + 18$, then $f(x)$ is strictly decreasing in -
 (A) $(-\infty, 1)$ (B) $[3, \infty)$ (C) $(-\infty, 1] \cup [3, \infty)$ (D) $(1, 3)$
- (6) If $x = -1$ and $x = 2$ are the extreme points of $y = \alpha \log x + \beta x^2 + x$ then
 (A) $\alpha = -6, \beta = \frac{1}{2}$ (B) $\alpha = -6, \beta = -\frac{1}{2}$ (C) $\alpha = 2, \beta = -\frac{1}{2}$ (D) $\alpha = 2, \beta = \frac{1}{2}$
- (7) The normal to the curve $x^2 + 2xy - 3y^2 = 0$ at $(1, 1)$
 (A) Meets the curve again in second quadrant. (B) Does not meet the curve again.
 (C) Meets the curve again in third quadrant. (D) Meets the curve again in fourth quadrant.
- (8) The equation of the tangent to the curve $y = 1 - e^{\frac{x}{2}}$ at the point of intersection with Y-axis is
 (A) $x + 2y = 0$ (B) $2x + y = 0$ (C) $x - y = 2$ (D) $x + y = 2$
- (9) If the tangent at $(1, 1)$ on $y^2 = x(2 - x)^2$ meets the curve again at P then P is
 (A) $(4, 4)$ (B) $(-1, 2)$ (C) $(3, 6)$ (D) $\left(\frac{9}{4}, \frac{3}{8}\right)$
- (10) The approximate value of $\tan(44^\circ 30')$ given that $1^\circ = 0.0175$.
 (A) 0.8952 (B) 0.9528 (C) 0.9285 (D) 0.9825

- (II) (1) If the curves $ax^2 + by^2 = 1$ and $a'x^2 + b'y^2 = 1$ intersect orthogonally, then prove that $\frac{1}{a} - \frac{1}{b} = \frac{1}{a'} - \frac{1}{b'}$.
- (2) Determine the area of the triangle formed by the tangent to the graph of the function $y = 3 - x^2$ drawn at the point $(1, 2)$ and the coordinate axes.
- (3) Find the equation of the tangent and normal drawn to the curve $y^4 - 4x^4 - 6xy = 0$ at the point $M(1, 2)$.
- (4) A water tank in the form of an inverted cone is being emptied at the rate of 2 cubic feet per second. The height of the cone is 8 feet and the radius is 4 feet. Find the rate of change of the water level when the depth is 6 feet.
- (5) Find all points on the ellipse $9x^2 + 16y^2 = 400$, at which the y-coordinate is decreasing and the x-coordinate is increasing at the same rate.
- (6) Verify Rolle's theorem for the function $f(x) = \frac{2}{e^x + e^{-x}}$ on $[-1, 1]$.
- (7) The position of a particle is given by the function $s(t) = 2t^2 + 3t - 4$. Find the time $t = c$ in the interval $0 \leq t \leq 4$ when the instantaneous velocity of the particle equals to its average velocity in this interval.
- (8) Find the approximate value of the function $f(x) = \sqrt{x^2 + 3x}$ at $x = 1.02$.
- (9) Find the approximate value of $\cos^{-1}(0.51)$ given $\pi = 3.1416$, $\frac{2}{\sqrt{3}} = 1.1547$.
- (10) Find the intervals on which the function $y = x^x$, ($x > 0$) is increasing and decreasing.
- (11) Find the intervals on the which the function $f(x) = \frac{x}{\log x}$, is increasing and decreasing.
- (12) An open box with a square base is to be made out of a given quantity of sheet of area a^2 , Show the maximum volume of the box is $\frac{a^3}{6\sqrt{3}}$.
- (13) Show that of all rectangles inscribed in a given circle, the square has the maximum area.
- (14) Show that a closed right circular cylinder of given surface area has maximum volume if its height equals the diameter of its base.
- (15) A window is in the form of a rectangle surmounted by a semi-circle. If the perimeter be 30 m, find the dimensions so that the greatest possible amount of light may be admitted.
- (16) Show that the height of a right circular cylinder of greatest volume that can be inscribed in a right circular cone is one-third of that of the cone.
- (17) A wire of length l is cut in to two parts. One part is bent into a circle and the other into a square. Show that the sum of the areas of the circle and the square is least, if the radius of the circle is half the side of the square.

- (18) A rectangular Sheet of paper of fixed perimeter with the sides having their length in the ratio 8 : 15 converted in to an open rectangular box by folding after removing the squares of equal area from all corners. If the total area of the removed squares is 100, the resulting box has maximum valume. Find the lengths of the rectangular sheet of paper.
- (19) Show that the altitude of the right circular cone of maximum volume that can be inscribed in a shpere of radius r is $\frac{4r}{3}$.
- (20) Show that the height of the cylinder of maximum volume that can be inscribed in a sphere of radius R is $\frac{2R}{\sqrt{3}}$. Also find the maximum volume.
- (21) Find the maximum and minimum values of the function $f(x) = \cos^2 x + \sin x$.



3. INDEFINITE INTEGRATION



Let us Study

- Definition and Properties
- Different Techniques : 1. by substitution 2. by parts 3. by partial fraction

Introduction :

In differential calculus, we studied differentiation or derivatives of some functions. We saw that derivatives are used for finding the slopes of tangents, maximum or minimum values of the function.

Now we will try to find the function whose derivative is known, or given $f(x)$. We will find $g(x)$ such that $g'(x) = f(x)$. Here the integration of $f(x)$ with respect to x is $g(x)$ or $g(x)$ is called the primitive of $f(x)$. For example, we know that the derivative of x^3 w. r. t. x is $3x^2$. So $\frac{d}{dx} x^3 = 3x^2$; and integral of $3x^2$ w. r. t. x is x^3 . This is shown with the sign of integration namely ' \int '. We write $\int 3x^2 \cdot dx = x^3$.

In this chapter we restrict ourselves only to study the methods of integration. The theory of integration is developed by Sir Isaac Newton and Gottfried Leibnitz.

$\int f(x) \cdot dx = g(x)$, read as an integral of $f(x)$ with respect to x , is $g(x)$. Since the derivative of constant function with respect to x is zero (0), we can also write

$\int f(x) \cdot dx = g(x) + c$, where c is an arbitrary constant and c can take infinitely many values.

For example :

$f(x) = x^2 + c$ represents family of curves for different values of c .

$f'(x) = 2x$ gives the slope of the tangent to $f(x) = x^2 + c$.

In the figure we have shown the curves

$$y = x^2, y = x^2 + 4, y = x^2 - 5.$$

Note that at the points (2, 4), (2, 8) (2, -1) respectively on those curves, the slopes of tangents are $2(2) = 4$.




Fig. 3.1.1

3.1.1 Elementary Integration Formulae

$$(i) \quad \frac{d}{dx} \left(\frac{x^{n+1}}{n+1} \right) = \frac{(n+1)x^n}{(n+1)}, n \neq -1$$

$$= x^n \quad \Rightarrow \quad \therefore \int x^n \cdot dx = \frac{x^{n+1}}{n+1} + c$$

$$\frac{d}{dx} \left(\frac{(ax+b)^{n+1}}{(n+1) \cdot a} \right) = \frac{(n+1)(ax+b)^n}{(n+1)}$$

$$= (ax+b)^n \quad \Rightarrow \quad \therefore \int (ax+b)^n \cdot dx = \frac{(ax+b)^{n+1}}{n+1} \cdot \frac{1}{a} + c$$

This result can be extended for n replaced by any rational $\frac{p}{q}$.

$$(ii) \quad \frac{d}{dx} \left(\frac{a^x}{\log a} \right) = a^x, a > 0 \quad \Rightarrow \quad \therefore \int a^x \cdot dx = \frac{a^x}{\log a} + c$$

$$\int A^{ax+b} \cdot dx = \frac{A^{ax+b}}{\log A} \cdot \frac{1}{a} + c, A > 0$$

$$(iii) \quad \frac{d}{dx} e^x = e^x \quad \Rightarrow \quad \int e^x \cdot dx = e^x + c$$

$$\int e^{ax+b} \cdot dx = e^{ax+b} \cdot \frac{1}{a} + c$$

$$(iv) \quad \frac{d}{dx} \sin x = \cos x \quad \Rightarrow \quad \int \cos x \cdot dx = \sin x + c$$

$$\int \cos(ax+b) \cdot dx = \sin(ax+b) \cdot \frac{1}{a} + c$$

$$(v) \quad \frac{d}{dx} \cos x = -\sin x \quad \Rightarrow \quad \int \sin x \cdot dx = -\cos x + c$$

$$\int \sin(ax+b) \cdot dx = -\cos(ax+b) \cdot \frac{1}{a} + c$$

$$(vi) \quad \frac{d}{dx} \tan x = \sec^2 x \quad \Rightarrow \quad \int \sec^2 x \cdot dx = \tan x + c$$

$$\int \sec^2(ax+b) \cdot dx = \tan(ax+b) \cdot \frac{1}{a} + c$$

$$(vii) \quad \frac{d}{dx} \sec x = \sec x \cdot \tan x \quad \Rightarrow \quad \int \sec x \cdot \tan x \cdot dx = \sec x + c$$

$$\int \sec(ax+b) \cdot \tan(ax+b) \cdot dx = \sec(ax+b) \cdot \frac{1}{a} + c$$

$$(viii) \quad \frac{d}{dx} \operatorname{cosec} x = -\operatorname{cosec} x \cdot \cot x \quad \Rightarrow \quad \int \operatorname{cosec} x \cdot \cot x \cdot dx = -\operatorname{cosec} x + c$$

$$\int \operatorname{cosec}(ax+b) \cdot \cot(ax+b) \cdot dx = -\operatorname{cosec}(ax+b) \cdot \frac{1}{a} + c$$

$$(ix) \quad \frac{d}{dx} \cot x = -\operatorname{cosec}^2 x \quad \Rightarrow \quad \int \operatorname{cosec}^2 x \cdot dx = -\cot x + c$$

$$\int \operatorname{cosec}^2(ax+b) \cdot dx = -\cot(ax+b) \cdot \frac{1}{a} + c$$

$$(x) \quad \frac{d}{dx} \log x = \frac{1}{x}, x > 0 \quad \Rightarrow \quad \int \frac{1}{x} dx = \log x + c, x \neq 0.$$

$$\therefore \text{also } \int \frac{1}{(ax+b)} \cdot dx = \log(ax+b) \cdot \frac{1}{a} + c$$

We assume that the trigonometric functions and logarithmic functions are defined on the respective domains.

3.1.2

Theorem 1 : If f and g are real valued integrable functions of x , then

$$\int [f(x) + g(x)] \cdot dx = \int f(x) \cdot dx + \int g(x) \cdot dx$$

Theorem 2 : If f and g are real valued integrable functions of x , then

$$\int [f(x) - g(x)] \cdot dx = \int f(x) \cdot dx - \int g(x) \cdot dx$$

Theorem 3 : If f and g are real valued integrable functions of x , and k is constant, then

$$\int k [f(x)] \cdot dx = k \int f(x) \cdot dx$$

Proof : 1. Let $\int f(x) \cdot dx = g_1(x) + c_1$ and $\int g(x) \cdot dx = g_2(x) + c_2$ then

$$\frac{d}{dx} [(g_1(x) + c_1)] = f(x) \quad \text{and} \quad \frac{d}{dx} [(g_2(x) + c_2)] = g(x)$$

$$\begin{aligned} \therefore \frac{d}{dx} [(g_1(x) + c_1) + (g_2(x) + c_2)] \\ &= \frac{d}{dx} [(g_1(x) + c_1)] + \frac{d}{dx} [(g_2(x) + c_2)] \\ &= f(x) + g(x) \end{aligned}$$

By definition of integration.

$$\begin{aligned} \int f(x) + g(x) &= (g_1(x) + c_1) + (g_2(x) + c_2) \\ &= \int f(x) \cdot dx + \int g(x) \cdot dx \end{aligned}$$

Note : Students can construct the proofs of the other two theorems (Theorem 2 and Theorem 3).



SOLVED EXAMPLES

Ex. : Evaluate the following :

1. $\int (x^3 + 3^x) \cdot dx$

Solution :

$$\begin{aligned} &\int (x^3 + 3^x) \cdot dx \\ &= \int x^3 \cdot dx + \int 3^x \cdot dx \\ &= \frac{x^4}{4} + \frac{3^x}{\log 3} + c \end{aligned}$$

2.
$$\int \left(\sin x + \frac{1}{x} + \frac{1}{\sqrt[3]{x}} \right) \cdot dx$$

Solution :
$$\int \left(\sin x + \frac{1}{x} + \frac{1}{\sqrt[3]{x}} \right) \cdot dx$$

$$= \int \sin x \cdot dx + \int \frac{1}{x} \cdot dx + \int \frac{1}{\sqrt[3]{x}} \cdot dx$$

$$= \int \sin x \cdot dx + \int \frac{1}{x} \cdot dx + \int x^{-\frac{1}{3}} \cdot dx$$

$$= -\cos x + \log x + \frac{x^{-\frac{1}{3}+1}}{-\frac{1}{3}+1} + c$$

$$= -\cos x + \log x + \frac{x^{\frac{2}{3}}}{\frac{2}{3}} + c$$

4.
$$\int \frac{\sqrt{x} + 1}{x + \sqrt{x}} \cdot dx$$

Solution :
$$\int \frac{\sqrt{x} + 1}{x + \sqrt{x}} \cdot dx$$

$$= \int \frac{\sqrt{x} + 1}{\sqrt{x}(\sqrt{x} + 1)} \cdot dx$$

$$= \int \frac{1}{\sqrt{x}} \cdot dx$$

$$= 2 \cdot \int \frac{1}{2\sqrt{x}} \cdot dx$$

$$= 2\sqrt{x} + c$$

3.
$$\int (\tan x + \cot x)^2 \cdot dx$$

Solution :
$$\int (\tan x + \cot x)^2 \cdot dx$$

$$= \int (\tan^2 x + 2 \tan x \cdot \cot x + \cot^2 x) \cdot dx$$

$$= \int (\tan^2 x + 2 + \cot^2 x) \cdot dx$$

$$= \int (\sec^2 x - 1 + 2 + \operatorname{cosec}^2 x - 1) \cdot dx$$

$$= \int (\sec^2 x + \operatorname{cosec}^2 x) \cdot dx$$

$$= \int \sec^2 x \cdot dx + \int \operatorname{cosec}^2 x \cdot dx$$

$$= \tan x + (-\cot x) + c$$

$$= \tan x - \cot x + c$$

5.
$$\int \frac{e^{4 \log x} - e^{5 \log x}}{x^5} \cdot dx$$

Solution :
$$\int \frac{e^{4 \log x} - e^{5 \log x}}{x^5} \cdot dx$$

$$= \int \frac{e^{\log x^4} - e^{\log x^5}}{x^5} \cdot dx, \quad \because a^{\log a f(x)} = f(x)$$

$$= \int \left(\frac{x^4 - x^5}{x^5} \right) \cdot dx$$

$$= \int \left(\frac{1}{x} - 1 \right) \cdot dx$$

$$= \log(x) - x + c$$

6.
$$\int \frac{2x + 3}{5x - 1} \cdot dx$$

Solution :
$$\frac{N}{D} = Q + \frac{R}{D}$$

$$(5x - 1) \overline{\begin{array}{r} \frac{2}{5} \\ 2x + 3 \\ - 2x - \frac{2}{5} \\ \hline 3 + \frac{2}{5} = \frac{17}{5} \end{array}}$$

$$\therefore 2x + 3 = \frac{2}{5}(5x - 1) + 3 + \frac{2}{5}$$

$$I = \int \left[\frac{2}{5} + \frac{\frac{17}{5}}{5x - 1} \right] \cdot dx$$

$$= \frac{2}{5}x + \frac{17}{5} \log(5x - 1) \cdot \frac{1}{5} + c$$

$$= \frac{2}{5}x + \frac{17}{25} \log(5x - 1) + c$$

7.
$$\int \frac{1}{\sqrt{3x+1} - \sqrt{3x-5}} \cdot dx$$

Solution :

$$\int \frac{1}{\sqrt{3x+1} - \sqrt{3x-5}} \cdot dx$$

$$= \int \left(\frac{1}{\sqrt{3x+1} - \sqrt{3x-5}} \right) \cdot \left(\frac{\sqrt{3x+1} + \sqrt{3x-5}}{\sqrt{3x+1} + \sqrt{3x-5}} \right) \cdot dx$$

$$= \int \frac{\sqrt{3x+1} + \sqrt{3x-5}}{3x+1 - 3x+5} \cdot dx$$

$$= \int \frac{\sqrt{3x+1} + \sqrt{3x-5}}{6} \cdot dx$$

$$= \frac{1}{6} \cdot \int \left((3x+1)^{\frac{1}{2}} + (3x-5)^{\frac{1}{2}} \right) \cdot dx$$

$$= \frac{1}{6} \cdot \left\{ \int (3x+1)^{\frac{1}{2}} \cdot dx + \int (3x-5)^{\frac{1}{2}} \cdot dx \right\}$$

$$= \frac{1}{6} \cdot \left\{ \frac{(3x+1)^{\frac{1}{2}+1}}{\left(\frac{1}{2}+1\right) \cdot 3} + \frac{(3x-5)^{\frac{1}{2}+1}}{\left(\frac{1}{2}+1\right) \cdot 3} \right\} + c$$

$$= \frac{1}{18} \cdot \left\{ \frac{2}{3} (3x+1)^{\frac{3}{2}} + \frac{2}{3} (3x-5)^{\frac{3}{2}} \right\} + c$$

$$= \frac{1}{27} \cdot \left\{ (3x+1)^{\frac{3}{2}} + (3x-5)^{\frac{3}{2}} \right\} + c$$

9.
$$\int \frac{x^3}{x-1} \cdot dx$$

Solution :

$$I = \int \frac{x^3 - 1 + 1}{x-1} \cdot dx$$

$$= \int \left(\frac{x^3 - 1}{x-1} + \frac{1}{x-1} \right) \cdot dx$$

$$= \int \left(\frac{(x-1)(x^2+x+1)}{(x-1)} + \frac{1}{x-1} \right) \cdot dx$$

$$= \int \left(x^2 + x + 1 + \frac{1}{x-1} \right) \cdot dx$$

$$= \frac{x^3}{3} + \frac{x^2}{2} + x + \log(x-1) + c$$

8.
$$\int \frac{2x-7}{\sqrt{3x-2}} \cdot dx$$

Solution : Express $(2x-7)$ in terms of $(3x-2)$

$$2x-7 = \frac{2}{3}(3x-2) + \frac{4}{3} - 7$$

$$= \frac{2}{3}(3x-2) - \frac{17}{3}$$

$$I = \int \left[\frac{\frac{2}{3}(3x-2) - \frac{17}{3}}{\sqrt{3x-2}} \right] \cdot dx$$

$$= \int \left[\frac{\frac{2}{3}(3x-2)}{\sqrt{3x-2}} - \frac{\frac{17}{3}}{\sqrt{3x-2}} \right] \cdot dx$$

$$= \frac{2}{3} \int \sqrt{3x-2} \cdot dx - \frac{17}{3} \int \frac{1}{\sqrt{3x-2}} \cdot dx$$

$$= \frac{2}{3} \int (3x-2)^{\frac{1}{2}} \cdot dx - \frac{17}{3} \int \frac{1}{\sqrt{3x-2}} \cdot dx$$

$$= \frac{2}{3} \cdot \frac{(3x-2)^{\frac{3}{2}}}{\left(\frac{3}{2}\right)} \cdot \frac{1}{3} - \frac{17}{3} \cdot 2 \cdot (\sqrt{3x-2}) \cdot \frac{1}{3} + c$$

$$= \frac{4}{27} \cdot (3x-2)^{\frac{3}{2}} - \frac{34}{9} \cdot (3x-2)^{\frac{1}{2}} + c$$

10.
$$\int \frac{3^x - 4^x}{5^x} \cdot dx$$

Solution :

$$I = \int \left(\frac{3^x}{5^x} - \frac{4^x}{5^x} \right) \cdot dx$$

$$= \int \left[\left(\frac{3}{5} \right)^x - \left(\frac{4}{5} \right)^x \right] \cdot dx$$

$$= \frac{\left(\frac{3}{5}\right)^x}{\log \frac{3}{5}} - \frac{\left(\frac{4}{5}\right)^x}{\log \frac{4}{5}} + c$$

11. $\int \cos^3 x \cdot dx$

Solution : $\cos 3A = 4 \cos^3 A - 3 \cos A$

$$\begin{aligned} I &= \int \frac{1}{4} (\cos 3x + 3 \cos x) \cdot dx \\ &= \frac{1}{4} \left(\sin 3x \cdot \frac{1}{3} + 3 \cdot \sin x \right) + c \\ &= \frac{1}{12} \sin 3x + \frac{3}{4} \sin x + c \end{aligned}$$

13. $\int \sin^4 x \cdot dx$

Solution :

$$\begin{aligned} I &= \int (\sin^2 x)^2 \cdot dx \\ &= \int \left(\frac{1}{2} (1 - \cos 2x) \right)^2 \cdot dx \\ &= \frac{1}{4} \cdot \int (1 - 2 \cos 2x + \cos^2 2x) \cdot dx \\ &= \frac{1}{4} \cdot \int \left[1 - 2 \cos 2x + \frac{1}{2} (1 + \cos 4x) \right] \cdot dx \\ &= \frac{1}{4} \cdot \int \left(1 - 2 \cos 2x + \frac{1}{2} + \frac{1}{2} \cos 4x \right) \cdot dx \\ &= \frac{1}{4} \cdot \int \left(\frac{3}{2} - 2 \cos 2x + \frac{1}{2} \cos 4x \right) \cdot dx \\ &= \frac{1}{4} \cdot \left[\frac{3}{2}x - 2 \sin 2x \cdot \frac{1}{2} + \frac{1}{2} \sin 4x \cdot \frac{1}{4} \right] + c \\ &= \frac{1}{4} \cdot \left[\frac{3}{2}x - \sin 2x + \frac{1}{8} \sin 4x \right] + c \end{aligned}$$

12. $\int \sqrt{1 + \sin 3x} \cdot dx$

Solution :

$$\begin{aligned} I &= \int \sqrt{\cos^2 \frac{3x}{2} + \sin^2 \frac{3x}{2} + 2 \sin \frac{3x}{2} \cdot \cos \frac{3x}{2}} \cdot dx \\ &= \int \sqrt{\left(\cos \frac{3x}{2} + \sin \frac{3x}{2} \right)^2} \cdot dx \\ &= \int \left(\cos \frac{3x}{2} + \sin \frac{3x}{2} \right) \cdot dx \\ &= \sin \frac{3x}{2} \cdot \frac{1}{\frac{3}{2}} - \cos \frac{3x}{2} \cdot \frac{1}{\frac{3}{2}} + c \\ &= \frac{2}{3} \left(\sin \frac{3x}{2} - \cos \frac{3x}{2} \right) + c \end{aligned}$$

14. $\int \sin 5x \cdot \cos 7x \cdot dx$

Solution : We know that

$$2 \sin A \cdot \cos B = \sin (A + B) + \sin (A - B)$$

$$\begin{aligned} I &= \frac{1}{2} \int 2 \sin 5x \cdot \cos 7x \cdot dx \\ &= \frac{1}{2} \int [\sin (5x + 7x) + \sin (5x - 7x)] \cdot dx \\ &= \frac{1}{2} \int [\sin (12x) + \sin (-2x)] \cdot dx \\ &= \frac{1}{2} \int (\sin 12x - \sin 2x) \cdot dx \\ &= \frac{1}{2} \cdot \left[-\cos 12x \cdot \frac{1}{12} + \cos 2x \cdot \frac{1}{2} \right] + c \\ I &= -\frac{1}{24} \cos 12x + \frac{1}{4} \cos 2x + c \end{aligned}$$

15. $\int \frac{\sin^3 x - \cos^3 x}{\sin^2 x \cdot \cos^2 x} \cdot dx$

Solution : $I = \int \left(\frac{\sin^3 x}{\sin^2 x \cdot \cos^2 x} - \frac{\cos^3 x}{\sin^2 x \cdot \cos^2 x} \right) \cdot dx$

$$= \int \left(\frac{\sin x}{\cos^2 x} - \frac{\cos x}{\sin^2 x} \right) \cdot dx$$

$$\begin{aligned} &= \int \left(\frac{1}{\cos x} \cdot \frac{\sin x}{\cos x} - \frac{1}{\sin x} \cdot \frac{\cos x}{\sin x} \right) \cdot dx \\ &= \int (\sec x \cdot \tan x - \operatorname{cosec} x \cdot \cot x) \cdot dx \\ &= \sec x - (-\operatorname{cosec} x) + c \\ I &= \sec x + \operatorname{cosec} x + c \end{aligned}$$

16. $\int \frac{1}{1 - \sin x} \cdot dx$

Solution :

$$\begin{aligned} I &= \int \left(\frac{1}{1 - \sin x} \right) \left(\frac{1 + \sin x}{1 + \sin x} \right) \cdot dx \\ &= \int \frac{1 + \sin x}{1 - \sin^2 x} \cdot dx \\ &= \int \frac{1 + \sin x}{\cos^2 x} \cdot dx \\ &= \int \left(\frac{1}{\cos^2 x} + \frac{\sin x}{\cos^2 x} \right) \cdot dx \\ &= \int (\sec^2 x + \sec x \cdot \tan x) \cdot dx \\ &= \tan x + \sec x + c \end{aligned}$$

17. $\int \left(\frac{\cos x}{1 - \cos x} \right) \cdot dx$

Solution :

$$\begin{aligned} I &= \int \left(\frac{\cos x}{1 - \cos x} \right) \left(\frac{1 + \cos x}{1 + \cos x} \right) \cdot dx \\ &= \int \frac{\cos x (1 + \cos x)}{1 - \cos^2 x} \cdot dx \\ &= \int \left(\frac{\cos x + \cos^2 x}{\sin^2 x} \right) \cdot dx \\ &= \int \left(\frac{\cos x}{\sin^2 x} + \frac{\cos^2 x}{\sin^2 x} \right) \cdot dx \\ &= \int (\operatorname{cosec} x \cdot \cot x + \cot^2 x) \cdot dx \\ &= \int (\operatorname{cosec} x \cdot \cot x + \operatorname{cosec}^2 x - 1) \cdot dx \\ &= (-\operatorname{cosec} x) + (-\cot x) - x + c \\ &= -\operatorname{cosec} x - \cot x - x + c \end{aligned}$$

Activity :

18. $\int \frac{\cos x - \cos 2x}{1 - \cos x} \cdot dx$

Solution :

$$\begin{aligned} &\int \frac{\cos x - \cos 2x}{1 - \cos x} \cdot dx \\ &= \int \frac{\cos x - (\dots\dots\dots)}{1 - \cos x} \cdot dx \\ &= \int \frac{\cos x - \dots\dots\dots}{1 - \cos x} \cdot dx \\ &= \int \frac{\cos x (1 - \cos x) + \dots\dots\dots}{1 - \cos x} \cdot dx \\ &= \int \left[\cos x + \frac{\dots\dots\dots}{1 - \cos x} \right] \cdot dx \\ &= \int [\cos x + (1 + \cos x)] \cdot dx \\ &= \int (1 + 2 \cos x) \cdot dx \\ &= x + 2 \sin x + c \end{aligned}$$



19. $\int \sin^{-1}(\cos 3x) \cdot dx$

Solution :

$$\begin{aligned} I &= \int \sin^{-1}\left(\sin \frac{\pi}{2} - 3x\right) \cdot dx \\ &= \int \left(\frac{\pi}{2} - 3x\right) \cdot dx \\ &= \frac{\pi}{2}x - 3 \frac{x^2}{2} + c \end{aligned}$$

20. $\int \tan^{-1}\left(\frac{\sin 2x}{1 + \cos 2x}\right) \cdot dx$

Solution :

$$\begin{aligned} I &= \int \cot^{-1}\left(\frac{1 + \cos 2x}{\sin 2x}\right) \cdot dx \\ &= \int \cot^{-1}\left(\frac{2 \cos^2 x}{2 \sin x \cdot \cos x}\right) \cdot dx \\ &= \int \cot^{-1}(\cot x) \cdot dx \\ &= \int x \cdot dx = \frac{x^2}{2} + c \end{aligned}$$

21. $\int \tan^{-1} \sqrt{\frac{1 - \sin x}{1 + \sin x}} \cdot dx$

Solution :

$$\begin{aligned} I &= \int \tan^{-1} \sqrt{\frac{1 - \cos\left(\frac{\pi}{2} - x\right)}{1 + \cos\left(\frac{\pi}{2} - x\right)}} \cdot dx \\ &= \int \tan^{-1} \sqrt{\frac{2 \sin^2\left(\frac{\pi}{4} - \frac{x}{2}\right)}{2 \cos^2\left(\frac{\pi}{4} - \frac{x}{2}\right)}} \cdot dx \\ &= \int \tan^{-1} \sqrt{\tan^2\left(\frac{\pi}{4} - \frac{x}{2}\right)} \cdot dx \\ &= \int \tan^{-1} \left[\tan\left(\frac{\pi}{4} - \frac{x}{2}\right) \right] \cdot dx \\ &= \int \left(\frac{\pi}{4} - \frac{x}{2}\right) \cdot dx \\ &= \frac{\pi}{4}x - \frac{1}{2} \cdot \frac{x^2}{2} + c \\ &= \frac{\pi}{4}x - \frac{x^2}{4} + c \end{aligned}$$

EXERCISE 3.1

I. Integrate the following functions w. r. t. x :

- (i) $x^3 + x^2 - x + 1$
- (ii) $x^2 \left(1 - \frac{2}{x}\right)^2$
- (iii) $3 \sec^2 x - \frac{4}{x} + \frac{1}{x\sqrt{x}} - 7$
- (iv) $2x^3 - 5x + \frac{3}{x} + \frac{4}{x^5}$
- (v) $\frac{3x^3 - 2x + 5}{x\sqrt{x}}$

II. Evaluate :

- (i) $\int \tan^2 x \cdot dx$
- (ii) $\int \frac{\sin 2x}{\cos x} \cdot dx$
- (iii) $\int \frac{\sin x}{\cos^2 x} \cdot dx$
- (iv) $\int \frac{\cos 2x}{\sin^2 x} \cdot dx$
- (v) $\int \frac{\cos 2x}{\sin^2 x \cdot \cos^2 x} \cdot dx$
- (vi) $\int \frac{\sin x}{1 + \sin x} \cdot dx$
- (vii) $\int \frac{\tan x}{\sec x + \tan x} \cdot dx$
- (viii) $\int \sqrt{1 + \sin 2x} \cdot dx$
- (ix) $\int \sqrt{1 - \cos 2x} \cdot dx$
- (x) $\int \sin 4x \cdot \cos 3x \cdot dx$

III. Evaluate :

- (i) $\int \frac{x}{x+2} \cdot dx$
- (ii) $\int \frac{4x+3}{2x+1} \cdot dx$
- (iii) $\int \frac{5x+2}{3x-4} \cdot dx$
- (iv) $\int \frac{x-2}{\sqrt{x+5}} \cdot dx$
- (v) $\int \frac{2x-7}{\sqrt{4x-1}} \cdot dx$
- (vi) $\int \frac{\sin 4x}{\cos 2x} \cdot dx$
- (vii) $\int \sqrt{1 + \sin 5x} \cdot dx$
- (viii) $\int \cos^2 x \cdot dx$
- (ix) $\int \frac{2}{\sqrt{x} - \sqrt{x+3}} \cdot dx$
- (x) $\int \frac{3}{\sqrt{7x-2} - \sqrt{7x-5}} \cdot dx$

IV. $f'(x) = x - \frac{3}{x^3}, f(1) = \frac{11}{2}$ then find $f(x)$.

3.2 Methods of integration :

We have evaluated the integrals which can be reduced to standard forms by algebraic or trigonometric simplifications. This year we are going to study three special methods of reducing an integral to a standard form, namely –

1. Integration by substitution
2. Integration by parts
3. Integration by partial fraction

3.2.1 Integration by substitution :

Theorem 1 : If $x = \phi(t)$ is a differentiable function of t , then $\int f(x) \cdot dx = \int f[\phi(t)] \cdot \phi'(t) dt$.

Proof : $x = \phi(t)$ is a differentiable function of t .

$$\therefore \frac{dx}{dt} = \phi'(t)$$

$$\text{Let } \int f(x) dx = g(x) \Rightarrow \frac{d}{dx} [g(x)] = f(x)$$

By Chain rule,

$$\begin{aligned} \frac{d}{dt} [g(x)] &= \frac{d}{dx} [g(x)] \cdot \frac{dx}{dt} \\ &= f(x) \cdot \frac{dx}{dt} \\ &= f[\phi(t)] \cdot \phi'(t) \end{aligned}$$

By definition of integration,

$$\begin{aligned} g(x) &= \int f[\phi(t)] \cdot \phi'(t) \cdot dt \\ \therefore \int f(x) \cdot dx &= \int f[\phi(t)] \cdot \phi'(t) \cdot dt \end{aligned}$$

For example 1 : $\int 3x^2 \sin(x^3) \cdot dx$

$$\text{Let } x^3 = t$$

$$\begin{aligned} \therefore 3x^2 dx &= dt \\ &= \int \sin t \cdot dt \\ &= -\cos t + c \\ &= -\cos(x^3) + c \end{aligned}$$

Corollary I :

If $\int f(x) \cdot dx = g(x) + c$

then $\int f(ax + b) \cdot dx = g(ax + b) \frac{1}{a} + c$

Proof : Let $I = \int f(ax + b) \cdot dx$

put $ax + b = t$

Differentiating both the sides

$a \cdot dx = 1 \cdot dt \Rightarrow dx = \frac{1}{a} dt$

$I = \int f(t) \cdot \frac{1}{a} \cdot dt$

$= \frac{1}{a} \cdot \int f(t) \cdot dt$

$= \frac{1}{a} \cdot g(t) + c$

$= \frac{1}{a} \cdot g(ax + b) + c$

$\therefore \int f(ax + b) \cdot dx = g(ax + b) \frac{1}{a} + c$

For example : $\int \sec^2(5x - 4) \cdot dx$

$= \frac{1}{5} \tan(5x - 4) + c$

Corollary III :

$\int \frac{f'(x)}{f(x)} \cdot dx = \log(f(x)) + c$

Proof : Consider $\int \frac{f'(x)}{f(x)} \cdot dx$

put $f(x) = t$

Differentiating both the sides

$f'(x) \cdot dx = dt$

$I = \int \frac{1}{t} \cdot dt$

$= \log(t) + c$

$= \log(f(x)) + c$

$\therefore \int \frac{f'(x)}{f(x)} \cdot dx = \log(f(x)) + c$

Corollary II :

$\int [f(x)]^n \cdot f'(x) \cdot dx = \frac{[f(x)]^{n+1}}{n+1} + c, n \neq -1$

Proof : Let $I = \int [f(x)]^{n+1} \cdot f'(x) \cdot dx$

put $f(x) = t$

Differentiating both the sides

$f'(x) \cdot dx = dt$

$I = \int [t]^n \cdot dt$

$= \frac{t^{n+1}}{n+1} + c, n \neq -1$

$= \frac{[f(x)]^{n+1}}{n+1} + c$

$\therefore \int [f(x)]^n \cdot f'(x) \cdot dx = \frac{[f(x)]^{n+1}}{n+1} + c$

For example : $\int \frac{(\sin^{-1}x)^3}{\sqrt{1-x^2}} \cdot dx$

$= \int [(\sin^{-1}x)^3] \cdot \left(\frac{1}{\sqrt{1-x^2}}\right) \cdot dx$

$= \frac{(\sin^{-1}x)^4}{4} + c$

For example : $\int \cot x \cdot dx$

$= \int \frac{\cos x}{\sin x} \cdot dx$

$\frac{d}{dx} \sin x = \cos x$

$= \int \left(\frac{\frac{d}{dx} \sin x}{\sin x} \cdot dx\right)$

$= \log(\sin x) + c$

Corollary IV :

$$\int \frac{f'(x)}{\sqrt{f(x)}} \cdot dx = 2 \sqrt{f(x)} + c$$

Proof : Consider $\int \frac{f'(x)}{\sqrt{f(x)}} \cdot dx$

put $f(x) = t$

Differentiating both the sides

$$f'(x) \cdot dx = dt$$

$$\begin{aligned} I &= \int \frac{1}{\sqrt{t}} \cdot dt \\ &= 2 \cdot \int \frac{1}{2\sqrt{t}} \cdot dt \\ &= 2 \sqrt{t} + c \\ &= 2 \sqrt{f(x)} + c \end{aligned}$$

$\therefore \int \frac{f'(x)}{\sqrt{f(x)}} \cdot dx = 2 \sqrt{f(x)} + c$

Using corollary III, $\int \frac{f'(x)}{f(x)} \cdot dx = \log (f(x)) + c$ we find the integrals of some trigonometric functions.

For example :
$$\begin{aligned} &\int \frac{1}{x\sqrt{\log x}} \cdot dx \\ &= \int \left(\frac{\frac{1}{x}}{\sqrt{\log x}} \right) \cdot dx \\ &= \int \left(\frac{\frac{d}{dx} \log x}{\sqrt{\log x}} \cdot dx \right) \\ &= 2 \sqrt{\log x} + c \end{aligned}$$

3.2.2 Integrals of trigonometric functions :

1. $\int \tan x \cdot dx$

Solution :

$$\begin{aligned} I &= \int \tan x \cdot dx \\ &= \int \frac{\sin x}{\cos x} \cdot dx \\ &= - \int \frac{-\sin x}{\cos x} \cdot dx \\ &= -\log (\cos x) + c \\ &= \log (\sec x) + c \end{aligned}$$

Activity :

2. $\int \cot (5x - 4) \cdot dx$

Solution :

$$\begin{aligned} I &= \int \frac{\dots\dots\dots}{\sin (5x - 4)} \cdot dx \\ &= \frac{1}{\dots\dots\dots} \int \frac{5 \cos (5x - 4)}{\dots\dots\dots} \cdot dx \\ \frac{d}{dx} (\dots\dots\dots) &= \dots\dots\dots \\ &= \frac{1}{5} \log [\sec (5x - 4)] + c \end{aligned}$$

3. $\int \sec x \cdot dx = \log (\sec x + \tan x) + c$

Solution : Let $I = \int \sec x \cdot dx$

$$= \int \frac{(\sec x)(\sec x + \tan x)}{\sec x + \tan x} \cdot dx$$

$$= \int \frac{\sec^2 x + \sec x \cdot \tan x}{\sec x + \tan x} \cdot dx$$

$$= \int \frac{\sec x \cdot \tan x + \sec^2 x}{\sec x + \tan x} \cdot dx$$

$$\therefore \frac{d}{dx} (\sec x + \tan x) = \sec x \cdot \tan x + \sec^2 x$$

$$\therefore \int \sec x \cdot dx = \log (\sec x + \tan x) + c$$

Also,

$$\int \sec x \cdot dx = \log \left[\tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right] + c$$

Activity :

4. $\int \operatorname{cosec} x \cdot dx = \log (\operatorname{cosec} x - \cot x) + c$

Solution : Let $I = \int \operatorname{cosec} x \cdot dx$

$$= \int \frac{(\operatorname{cosec} x)(\dots\dots\dots)}{(\dots\dots\dots)} \cdot dx$$

$$= \int \frac{\dots\dots\dots}{\dots\dots\dots} \cdot dx$$

$$= \int \frac{-\operatorname{cosec} x \cdot \cot x + \operatorname{cosec}^2 x}{\dots\dots\dots} \cdot dx$$

$$\frac{d}{dx} (\operatorname{cosec} x - \cot x)$$

$$= \dots\dots\dots$$

$$= \log (\operatorname{cosec} x - \cot x) + c$$

$$\therefore \int \operatorname{cosec} x \cdot dx = \log (\operatorname{cosec} x - \cot x) + c$$

Also,

$$\int \operatorname{cosec} x \cdot dx = \log \left(\tan \frac{x}{2} \right) + c$$

SOLVED EXAMPLES

Ex. : Evaluate the following functions :

1. $\int \frac{\cot (\log x)}{x} \cdot dx$

Solution : Let $I = \int \frac{\cot (\log x)}{x} \cdot dx$

put $\log x = t$

$$\therefore \frac{1}{x} \cdot dx = 1 \cdot dt$$

$$= \int \cot t \cdot dt$$

$$= \log (\sin t) + c$$

$$= \log (\sin \log x) + c$$

2. $\int \frac{\cos \sqrt{x}}{\sqrt{x}} \cdot dx$

Solution : Let $I = \int \frac{\cos \sqrt{x}}{\sqrt{x}} \cdot dx$

put $\sqrt{x} = t$

$$\therefore \frac{1}{2\sqrt{x}} \cdot dx = 1 \cdot dt$$

$$\therefore \frac{1}{\sqrt{x}} \cdot dx = 2 \cdot dt$$

$$= 2 \cdot \int \cos t \cdot dt$$

$$= 2 \cdot \sin t + c$$

$$= 2 \cdot \sin \sqrt{x} + c$$

3. $\int \frac{\sec^8 x}{\operatorname{cosec} x} \cdot dx$

Solution : $I = \int \sec^7 x \cdot \sec x \cdot \frac{1}{\operatorname{cosec} x} \cdot dx$
 $= \int \sec^7 x \cdot \frac{1}{\cos x} \cdot \sin x \cdot dx$
 $= \int \sec^7 x \cdot \tan x \cdot dx$
 $= \int \sec^6 x \cdot \sec x \cdot \tan x \cdot dx$
 put $\sec x = t$
 $\therefore \sec x \cdot \tan x \cdot dx = dt$
 $= \int t^6 \cdot dt$
 $= \frac{t^7}{7} + c$
 $= \frac{\sec^7 x}{7} + c$

5. $\int 5^{5^x} \cdot 5^x \cdot dx$

Solution : $I = \int 5^{5^x} \cdot 5^x \cdot dx$
 put $5^x = t$
 $\therefore 5^x \cdot \log 5 \cdot dx = 1 dt$
 $5^x \cdot dx = \frac{1}{\log 5} \cdot dt$
 $= \int 5^t \cdot \frac{1}{\log 5} \cdot dt$
 $I = \frac{1}{\log 5} \cdot \int 5^t \cdot dt$
 $= \frac{1}{\log 5} \cdot 5^t \cdot \frac{1}{\log 5} + c$
 $= \left(\frac{1}{\log 5} \right)^2 \cdot 5^{5^x} + c$

7. $\int \frac{e^x (1+x)}{\cos (x \cdot e^x)} \cdot dx$

Solution : put $x \cdot e^x = t$
 Differentiating both sides
 $(x \cdot e^x + e^x \cdot 1) \cdot dx = 1 dt$
 $e^x (1+x) \cdot dx = 1 dt$

4. $\int \frac{1}{x + \sqrt{x}} \cdot dx$

Solution : $I = \int \frac{1}{x + \sqrt{x}} \cdot dx$
 $= \int \frac{1}{\sqrt{x} (\sqrt{x} + 1)} \cdot dx$
 put $\sqrt{x} + 1 = t$
 $\therefore \frac{1}{2\sqrt{x}} \cdot dx = 1 \cdot dt$
 $\therefore \frac{1}{\sqrt{x}} \cdot dx = 2 \cdot dt$
 $= \int \frac{1}{t} \cdot 2 \cdot dt$
 $= 2 \cdot \int \frac{1}{t} \cdot dt$
 $= 2 \cdot \log (t) + c$
 $= 2 \cdot \log (\sqrt{x} + 1) + c$

6. $\int \frac{1}{1 + e^{-x}} \cdot dx$

Solution : $I = \int \frac{1}{1 + e^{-x}} \cdot dx$
 $= \int \frac{1}{1 + \frac{1}{e^x}} \cdot dx$
 $= \int \frac{1}{\frac{e^x + 1}{e^x}} \cdot dx$
 $= \int \frac{e^x}{e^x + 1} \cdot dx$
 $\therefore \frac{d}{dx} (e^x + 1) \cdot dx = e^x$
 $= \log [e^x + 1] + c$

$I = \int \frac{1}{\cos t} \cdot dt$

$= \int \sec t \cdot dt$
 $= \log (\sec t + \tan t) + c$
 $= \log (\sec (xe^x) + \tan (xe^x)) + c$

8. $\int \frac{1}{3x + 7x^{-n}} \cdot dx$

Solution : Consider $\int \frac{1}{3x + 7x^{-n}} \cdot dx$

$$= \int \frac{1}{3x + \frac{7}{x^n}} \cdot dx = \int \frac{1}{\frac{3x^{n+1} + 7}{x^n}} \cdot dx$$

$$= \int \frac{x^n}{3x^{n+1} + 7} \cdot dx$$

put $3x^{n+1} + 7 = t$
Differentiate w. r. t. x
 $3(n+1)x^n \cdot dx = dt$

$$\therefore x^n \cdot dx = \frac{1}{3(n+1)} dt$$

$$= \int \frac{1}{\frac{3(n+1)}{t}} \cdot dt$$

$$= \frac{1}{3(n+1)} \cdot \log(t) + c$$

$$= \frac{1}{3(n+1)} \cdot \log(3x^{n+1} + 7) + c$$

9. $\int (3x + 2) \sqrt{x-4} \cdot dx$

Solution : put $x - 4 = t$
 $\therefore x = 4 + t$
Differentiate
 $1 \cdot dx = 1 \cdot dt$

$$= \int [3(4 + t) + 2] \cdot \sqrt{t} \cdot dt$$

$$= \int (14 + 3t) \cdot t^{\frac{1}{2}} \cdot dt$$

$$= \int \left(14t^{\frac{1}{2}} + 3t^{\frac{3}{2}} \right) \cdot dt$$

$$= 14 \frac{t^{\frac{3}{2}}}{\frac{3}{2}} + 3 \frac{t^{\frac{5}{2}}}{\frac{5}{2}} \cdot dx$$

$$= \frac{28}{3} (x-4)^{\frac{3}{2}} + \frac{6}{5} (x-4)^{\frac{5}{2}} + c$$

10. $\int \frac{\sin(x+a)}{\cos(x-b)} \cdot dx$

Solution :

$$= \int \frac{\sin[(x-b) + (a+b)]}{\cos(x-b)} \cdot dx$$

$$= \int \frac{\sin(x-b) \cdot \cos(a+b) + \cos(x-b) \cdot \sin(a+b)}{\cos(x-b)} \cdot dx$$

$$= \int \left[\frac{\sin(x-b) \cdot \cos(a+b)}{\cos(x-b)} + \frac{\cos(x-b) \cdot \sin(a+b)}{\cos(x-b)} \right]$$

$$= \int [\cos(a+b) \cdot \tan(x-b) + \sin(a+b)] \cdot dx$$

$$= \cos(a+b) \cdot \log(\sec(x-b)) + x \cdot \sin(a+b) + c$$

EXERCISE 3.2 (A)

I. Integrate the following functions w. r. t. x :

1. $\frac{(\log x)^n}{x}$
2. $\frac{(\sin^{-1} x)^{\frac{3}{2}}}{\sqrt{1-x^2}}$
3. $\frac{1+x}{x \cdot \sin(x+\log x)}$
4. $\frac{x \cdot \sec^2(x^2)}{\sqrt{\tan^3(x^2)}}$
5. $\frac{e^{3x}}{e^{3x}+1}$
6. $\frac{(x^2+2)}{(x^2+1)} \cdot a^{x+\tan^{-1}x}$
7. $\frac{e^x \cdot \log(\sin e^x)}{\tan(e^x)}$
8. $\frac{e^{2x}+1}{e^{2x}-1}$
9. $\sin^4 x \cdot \cos^3 x$
10. $\frac{1}{4x+5x^{-11}}$
11. $x^9 \cdot \sec^2(x^{10})$
12. $e^{3 \log x} \cdot (x^4+1)^{-1}$
13. $\frac{\sqrt{\tan x}}{\sin x \cdot \cos x}$
14. $\frac{(x-1)^2}{(x^2+1)^2}$
15. $\frac{2 \sin x \cdot \cos x}{3 \cos^2 x + 4 \sin^2 x}$
16. $\frac{1}{\sqrt{x} + \sqrt{x^3}}$
17. $\frac{10x^9 + 10^x \cdot \log 10}{10^x + x^{10}}$
18. $\frac{x^{n-1}}{\sqrt{1+4x^n}}$
19. $(2x+1)\sqrt{x+2}$
20. $x^5 \cdot \sqrt{a^2+x^2}$
21. $(5-3x)(2-3x)^{-\frac{1}{2}}$
22. $\frac{7+4x+5x^2}{(2x+3)^{\frac{3}{2}}}$

23. $\frac{x^2}{\sqrt{9-x^6}}$
24. $\frac{1}{x(x^3-1)}$
25. $\frac{1}{x \cdot \log x \cdot \log(\log x)}$

II. Integrate the following functions w. r. t. x :

1. $\frac{\cos 3x - \cos 4x}{\sin 3x + \sin 4x}$
2. $\frac{\cos x}{\sin(x-a)}$
3. $\frac{\sin(x-a)}{\cos(x+b)}$
4. $\frac{1}{\sin x \cdot \cos x + 2 \cos^2 x}$
5. $\frac{\sin x + 2 \cos x}{3 \sin x + 4 \cos x}$
6. $\frac{1}{2+3 \tan x}$
7. $\frac{4e^x - 25}{2e^x - 5}$
8. $\frac{20+12e^x}{3e^x+4}$
9. $\frac{3e^{2x}+5}{4e^{2x}-5}$
10. $\cos^8 x \cdot \cot x$
11. $\tan^5 x$
12. $\cos^7 x$
13. $\tan 3x \cdot \tan 2x \cdot \tan x$
14. $\sin^5 x \cdot \cos^8 x$
15. $3^{\cos^2 x} \cdot \sin 2x$
16. $\frac{\sin 6x}{\sin 10x \cdot \sin 4x}$
17. $\frac{\sin x \cdot \cos^3 x}{1+\cos^2 x}$

3.2.3 Some Special Integrals

1. $\int \frac{1}{x^2+a^2} \cdot dx = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + c$
2. $\int \frac{1}{x^2-a^2} \cdot dx = \frac{1}{2a} \log\left(\frac{x-a}{x+a}\right) + c$
3. $\int \frac{1}{a^2-x^2} \cdot dx = \frac{1}{2a} \log\left(\frac{a+x}{a-x}\right) + c$
4. $\int \frac{1}{\sqrt{a^2-x^2}} \cdot dx = \sin^{-1}\left(\frac{x}{a}\right) + c$
5. $\int \frac{1}{\sqrt{x^2-a^2}} \cdot dx = \log(x + \sqrt{x^2-a^2}) + c$
6. $\int \frac{1}{\sqrt{x^2+a^2}} \cdot dx = \log(x + \sqrt{x^2+a^2}) + c$
7. $\int \frac{1}{x\sqrt{x^2-a^2}} \cdot dx = \frac{1}{a} \sec^{-1}\left(\frac{x}{a}\right) + c$

While evaluating an integral there is no unique substitution, we can use some standard substitutions and try.

No.	Function	Substitution
1.	$\sqrt{a^2 - x^2}$	$x = a \cdot \sin \theta$ ($x = a \cdot \cos \theta$ can also be used.)
2.	$\sqrt{a^2 + x^2}$	$x = a \cdot \tan \theta$
3.	$\sqrt{x^2 - a^2}$	$x = a \cdot \sec \theta$
4.	$\sqrt{\frac{a-x}{a+x}}$	$x = a \cdot \cos 2\theta$

1.
$$\int \frac{1}{x^2 + a^2} \cdot dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$$

Proof :

Let $I = \int \frac{1}{x^2 + a^2} \cdot dx$

put $x = a \cdot \tan \theta \Rightarrow \tan \theta = \frac{x}{a}$

i.e. $\theta = \tan^{-1} \left(\frac{x}{a} \right)$

$\therefore dx = a \cdot \sec^2 \theta \cdot d\theta$

$$I = \int \frac{1}{a^2 \cdot \tan^2 \theta + a^2} \cdot a \cdot \sec^2 \theta \cdot d\theta$$

$$= \int \frac{a \cdot \sec^2 \theta}{a^2 (\tan^2 \theta + 1)} \cdot d\theta$$

$$= \int \frac{\sec^2 \theta}{a \cdot \sec^2 \theta} \cdot d\theta$$

$$= \frac{1}{a} \int d\theta$$

$$= \frac{1}{a} \theta + c$$

$$= \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$$

$\therefore \int \frac{1}{x^2 + a^2} \cdot dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$

e.g.
$$\int \frac{1}{x^2 + 5^2} \cdot dx = \frac{1}{5} \tan^{-1} \left(\frac{x}{5} \right) + c$$

Alternatively

Consider,

$$\frac{d}{dx} \left[\frac{1}{a} \cdot \tan^{-1} \left(\frac{x}{a} \right) + c \right]$$

$$= \frac{d}{dx} \left[\frac{1}{a} \cdot \tan^{-1} \left(\frac{x}{a} \right) \right] + \frac{d}{dx} c$$

$$= \frac{1}{a} \cdot \frac{1}{1 + \left(\frac{x}{a} \right)^2} \cdot \frac{d}{dx} \left(\frac{x}{a} \right) + 0$$

$$= \frac{1}{a} \cdot \frac{1}{1 + \frac{x^2}{a^2}} \cdot \frac{1}{a}$$

$$= \frac{1}{a^2} \cdot \frac{1}{\frac{a^2 + x^2}{a^2}}$$

$$= \frac{1}{x^2 + a^2}$$

Therefore,

by definition of integration

$$\therefore \int \frac{1}{x^2 + a^2} \cdot dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$$

$$2. \int \frac{1}{x^2 - a^2} \cdot dx = \frac{1}{2a} \log \left(\frac{x - a}{x + a} \right) + c$$

Proof :

$$\begin{aligned} \text{Let } I &= \int \frac{1}{x^2 - a^2} \cdot dx \\ &= \int \frac{1}{(x + a)(x - a)} \cdot dx \\ &= \int \frac{1}{2a} \cdot \left[\frac{1}{x - a} - \frac{1}{x + a} \right] \cdot dx \\ &= \frac{1}{2a} \cdot \int \left[\frac{1}{x - a} - \frac{1}{x + a} \right] \cdot dx \\ &= \frac{1}{2a} \cdot [\log (x - a) - \log (x + a)] + c \\ &= \frac{1}{2a} \cdot \log \left(\frac{x - a}{x + a} \right) + c \end{aligned}$$

$$\therefore \int \frac{1}{x^2 - a^2} \cdot dx = \frac{1}{2a} \log \left(\frac{x - a}{x + a} \right) + c$$

e.g. $\int \frac{1}{x^2 - 9} \cdot dx = \frac{1}{2(3)} \log \left(\frac{x - 3}{x + 3} \right) + c$

Activity :

$$3. \int \frac{1}{a^2 - x^2} \cdot dx = \frac{1}{2a} \log \left(\frac{a + x}{a - x} \right) + c$$

Proof : Consider,

$$\begin{aligned} I &= \int \frac{1}{a^2 - x^2} \cdot dx \\ &= \int \frac{1}{(\dots)(\dots)} \cdot dx \\ &= \int \frac{1}{2a} \cdot \left[\frac{1}{\dots} - \frac{1}{\dots} \right] \cdot dx \\ &= \frac{1}{2a} \cdot \int \left[\frac{\dots}{\dots} - \frac{1}{a + x} \right] \cdot dx \\ &= \frac{1}{2a} \cdot [\log (a + x) - \log (a - x)] + c \\ &= \frac{1}{2a} \cdot \log \left(\frac{\dots}{\dots} \right) + c \end{aligned}$$

$$\therefore \int \frac{1}{a^2 - x^2} \cdot dx = \frac{1}{2a} \log \left(\frac{a + x}{a - x} \right) + c$$

e.g. $\int \frac{1}{16 - x^2} \cdot dx = \frac{1}{2(4)} \log \left(\frac{4 + x}{4 - x} \right) + c$

$$4. \int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx = \sin^{-1} \left(\frac{x}{a} \right) + c$$

Proof :

$$\text{Let } I = \int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx$$

$$\text{put } x = a \sin \theta \Rightarrow \sin \theta = \frac{x}{a}$$

$$\therefore \theta = \sin^{-1} \left(\frac{x}{a} \right)$$

$$dx = a \cdot \cos \theta \, d\theta$$

$$I = \int \frac{1}{\sqrt{a^2 - a^2 \sin^2 \theta}} \cdot a \cdot \cos \theta \, d\theta$$

$$I = \int \frac{a \cdot \cos \theta}{a \sqrt{1 - a^2 \sin^2 \theta}} \cdot d\theta$$

$$= \int \frac{\cos \theta}{\cos \theta} \cdot d\theta$$

$$= \int 1 \cdot d\theta$$

$$= \theta + c$$

$$\therefore \int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx = \sin^{-1} \left(\frac{x}{a} \right) + c$$

e.g. $\int \frac{1}{\sqrt{81 - x^2}} \cdot dx = \sin^{-1} \left(\frac{x}{9} \right) + c$

$$5. \int \frac{1}{\sqrt{x^2 - a^2}} \cdot dx = \log(x + \sqrt{x^2 - a^2}) + c$$

Proof: Let $I = \int \frac{1}{\sqrt{x^2 - a^2}} \cdot dx$

put $x = a \sec \theta \Rightarrow \theta = \sec^{-1}\left(\frac{x}{a}\right)$

$$\therefore dx = a \cdot \sec \theta \cdot \tan \theta \cdot d\theta$$

$$I = \int \frac{1}{\sqrt{a^2 \sec^2 \theta - a^2}} \cdot a \cdot \sec \theta \cdot \tan \theta \cdot d\theta$$

$$= \int \frac{a \cdot \sec \theta \cdot \tan \theta}{\sqrt{a^2(\sec^2 \theta - 1)}} \cdot d\theta$$

$$= \int \frac{a \cdot \sec \theta \cdot \tan \theta}{\sqrt{a^2 \cdot \tan^2 \theta}} \cdot d\theta$$

$$= \int \frac{a \cdot \sec \theta \cdot \tan \theta}{a \cdot \tan \theta} \cdot d\theta$$

$$= \int \sec \theta \cdot d\theta$$

$$= \log(\sec \theta + \tan \theta) + c$$

$$= \log(\sec \theta + \sqrt{\sec^2 \theta - 1}) + c$$

$$= \log\left(\frac{x}{a} + \sqrt{\frac{x^2}{a^2} - 1}\right) + c_1$$

$$= \log\left(\frac{x}{a} + \sqrt{\frac{x^2}{a^2} - 1}\right) + c_1$$

$$= \log\left(\frac{x + \sqrt{x^2 - a^2}}{a}\right) + c_1$$

$$= \log(x + \sqrt{x^2 - a^2}) - \log a + c_1$$

$$= \log(x + \sqrt{x^2 - a^2}) + c$$

where $c = c_1 - \log a$

$$\therefore \int \frac{1}{\sqrt{x^2 - a^2}} \cdot dx = \log(x + \sqrt{x^2 - a^2}) + c$$

e.g. $\int \frac{1}{\sqrt{x^2 - 16}} \cdot dx = \log(x + \sqrt{x^2 - 16}) + c$

Activity :

$$6. \int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx = \log(x + \sqrt{a^2 - x^2}) + c$$

Proof: use substitution $x = a \cdot \tan \theta$

.....

.....

.....

.....

.....

Activity :

$$7. \int \frac{1}{x\sqrt{x^2 - a^2}} \cdot dx = \frac{1}{a} \sec^{-1}\left(\frac{x}{a}\right) + c$$

Proof: Let $I = \int \frac{1}{x\sqrt{x^2 - a^2}} \cdot dx$

put $x = a \sec \theta \Rightarrow \theta = \sec^{-1}\left(\frac{x}{a}\right)$

$$\therefore dx = a \cdot \sec \theta \cdot \tan \theta \cdot d\theta$$

$$I = \int \frac{1}{a \sec \theta \sqrt{\dots - a^2}} \cdot \dots$$

$$= \int \frac{\tan \theta}{\sqrt{a^2(\dots)}} \cdot d\theta$$

$$= \frac{1}{a} \int 1 \cdot d\theta$$

$$= \frac{1}{a} \cdot \theta + c$$

$$= \frac{1}{a} \cdot \sec^{-1}\left(\frac{x}{a}\right) + c$$

$$\therefore \int \frac{1}{x\sqrt{x^2 - a^2}} \cdot dx = \frac{1}{a} \sec^{-1}\left(\frac{x}{a}\right) + c$$

e.g. $\int \frac{1}{x\sqrt{x^2 - 64}} \cdot dx = \frac{1}{8} \sec^{-1}\left(\frac{x}{8}\right) + c$

3.2.4

In order to evaluate the integrals of type $\int \frac{1}{ax^2 + bx + c} \cdot dx$ and $\int \frac{1}{\sqrt{ax^2 + bx + c}} \cdot dx$ we can use the following steps.

- (1) Write $ax^2 + bx + c$ as, $a\left(x^2 + \frac{b}{a}x + \frac{c}{a}\right)$, $a > 0$ and take a or \sqrt{a} out of the integral sign.
- (2) $\left(x^2 + \frac{b}{a}x\right)$ or $\left(\frac{b}{a}x - x^2\right)$ is expressed by the method of completing square by adding and subtracting $\left(\frac{1}{2} \text{ coefficient of } x\right)^2$.
- (3) Express the quadratic expression as a sum or difference of two squares
i.e. $((x + \beta)^2 \pm \alpha^2)$ or $(\alpha^2 - (x + \beta)^2)$
- (4) We know that $\int f(x) dx = g(x) + c \Rightarrow \int f(x + \beta) dx = g(x + \beta) + c$
 $\int f(ax + \beta) dx = \frac{1}{\alpha} g(ax + \beta) + c$
- (5) Use the standard integral formula and express the result in terms of x .

3.2.5

In order to evaluate the integral of type $\int \frac{1}{a \sin^2 x + b \cos^2 x + c} \cdot dx$ we can use the following steps.

- (1) Divide the numerator and denominator by $\cos^2 x$ or $\sin^2 x$.
- (2) In denominator replace $\sec^2 x$ by $1 + \tan^2 x$ and /or $\operatorname{cosec}^2 x$ by $1 + \cot^2 x$, if exists.
- (3) Put $\tan x = t$ or $\cot x = t$ so that the integral reduces to the form $\int \frac{1}{at^2 + bt + c} \cdot dt$
- (4) Use the standard integral formula and express the result in terms of x .

3.2.6

To evaluate the integral of the form $\int \frac{1}{a \sin x + b \cos x + c} \cdot dx$, we use the standard substitution $\tan \frac{x}{2} = t$.

If $\tan \frac{x}{2} = t$ then (i) $\sec^2 \frac{x}{2} \cdot \frac{1}{2} \cdot dx = 1 \cdot dt$
 i.e. $dx = \frac{2}{\sec^2 \frac{x}{2}} \cdot dt = \frac{2}{1 + \tan^2 \frac{x}{2}} \cdot dt = \frac{2 dt}{1 + t^2}$
 (ii) $\sin x = \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} = \frac{2t}{1 + t^2}$



$$(iii) \cos x = \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} = \frac{1 - t^2}{1 + t^2}$$

We put $\tan x = t$ for the integral of the type $\int \frac{1}{a \sin 2x + b \cos 2x + c} \cdot dx$

therefore $dx = \frac{1}{1 + t^2} \cdot dt$

$$\sin 2x = \frac{2t}{1 + t^2} \cdot dt$$

and $\cos 2x = \frac{1 - t^2}{1 + t^2} \cdot dt$

With this substitution the integral reduces to the form $\int \frac{1}{ax^2 + bx + c} \cdot dx$. Now use the standard integral formula and express the result in terms of x .

 **SOLVED EXAMPLES**

Ex. : Evaluate :

1. $\int \frac{1}{4x^2 + 11} \cdot dx$

Solution : $I = \int \frac{1}{4 \left(x^2 + \frac{11}{4} \right)} \cdot dx$
 $= \frac{1}{4} \cdot \int \frac{1}{x^2 + \left(\frac{\sqrt{11}}{2} \right)^2} \cdot dx$

$\therefore \int \frac{1}{x^2 + a^2} \cdot dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$

$$I = \frac{1}{4} \cdot \left[\frac{1}{\left(\frac{\sqrt{11}}{2} \right)} \right] \cdot \tan^{-1} \left[\frac{x}{\left(\frac{\sqrt{11}}{2} \right)} \right] + c$$

$$= \frac{1}{2\sqrt{11}} \tan^{-1} \left(\frac{2x}{\sqrt{11}} \right) + c$$

2. $\int \frac{1}{a^2 - b^2 x^2} \cdot dx$

Solution : $I = \int \frac{1}{b^2 \left(\frac{a^2}{b^2} - x^2 \right)} \cdot dx$
 $= \frac{1}{b^2} \cdot \int \frac{1}{\left(\frac{a}{b} \right)^2 - x^2} \cdot dx$

$\therefore \int \frac{1}{a^2 - x^2} \cdot dx = \frac{1}{2a} \log \left(\frac{a+x}{a-x} \right) + c$

$$I = \frac{1}{b^2} \cdot \frac{1}{2 \left(\frac{a}{b} \right)} \cdot \log \left(\frac{\frac{a}{b} + x}{\frac{a}{b} - x} \right) + c$$

$$= \frac{1}{b^2} \cdot \frac{1}{2 \left(\frac{a}{b} \right)} \cdot \log \left(\frac{\frac{a}{b} + x}{\frac{a}{b} - x} \right) + c$$

$$= \frac{1}{2ab} \cdot \log \left(\frac{a + bx}{a - bx} \right) + c$$

3. $\int \frac{1}{\sqrt{3x^2 - 7}} \cdot dx$

Solution : $I = \int \frac{1}{\sqrt{3\left(x^2 - \frac{7}{3}\right)}} \cdot dx$
 $= \int \frac{1}{\sqrt{3} \cdot \sqrt{x^2 - \left(\frac{\sqrt{7}}{\sqrt{3}}\right)^2}} \cdot dx$
 $= \frac{1}{\sqrt{3}} \cdot \int \frac{1}{\sqrt{x^2 - \left(\frac{\sqrt{7}}{\sqrt{3}}\right)^2}} \cdot dx$

$\therefore \int \frac{1}{\sqrt{x^2 - a^2}} \cdot dx = \log |x + \sqrt{x^2 - a^2}| + c$
 $I = \frac{1}{\sqrt{3}} \cdot \log \left(x + \sqrt{x^2 - \left(\frac{\sqrt{7}}{\sqrt{3}}\right)^2} \right) + c$
 $= \frac{1}{\sqrt{3}} \cdot \log \left(x + \sqrt{x^2 - \frac{7}{3}} \right) + c$

4. $\int \frac{1}{x^2 + 8x + 12} \cdot dx$

Solution : $I = \int \frac{1}{x^2 + 8x + 16 - 4} \cdot dx$
 $= \int \frac{1}{(x + 4)^2 - (2)^2} \cdot dx$
 $\therefore \int \frac{1}{x^2 - a^2} \cdot dx = \frac{1}{2a} \log \left(\frac{x - a}{x + a} \right) + c$
 $I = \frac{1}{2(2)} \cdot \log \left(\frac{(x + 4) - 2}{(x + 4) + 2} \right) + c$
 $= \frac{1}{4} \cdot \log \left(\frac{x + 2}{x + 6} \right) + c$

$\therefore \int \frac{1}{x^2 + 8x + 12} \cdot dx = \frac{1}{4} \cdot \log \left(\frac{x + 2}{x + 6} \right) + c$

5. $\int \frac{1}{\sqrt{3x^2 - 4x + 2}} \cdot dx$

Solution : $= \int \frac{1}{\sqrt{3\left(x^2 - \frac{4}{3}x + \frac{2}{3}\right)}} \cdot dx$

$\therefore \left\{ \left(\frac{1}{2} \text{ coefficient of } x \right)^2 = \left(\frac{1}{2} \left(-\frac{4}{3} \right) \right)^2 = \left(-\frac{2}{3} \right)^2 = \frac{4}{9} \right\}$

$= \int \frac{1}{\sqrt{3} \cdot \sqrt{x^2 - \frac{4}{3}x + \frac{4}{9} - \frac{4}{9} + \frac{2}{3}}} \cdot dx$
 $= \frac{1}{\sqrt{3}} \cdot \int \frac{1}{\sqrt{\left(x^2 - \frac{4}{3}x + \frac{4}{9}\right) + \left(\frac{2}{3} - \frac{4}{9}\right)}} \cdot dx$
 $= \frac{1}{\sqrt{3}} \cdot \int \frac{1}{\sqrt{\left(x - \frac{2}{3}\right)^2 + \left(\frac{\sqrt{2}}{3}\right)^2}} \cdot dx$

$\therefore \int \frac{1}{\sqrt{x^2 + a^2}} \cdot dx = \log |x + \sqrt{x^2 + a^2}| + c$
 $= \frac{1}{\sqrt{3}} \cdot \log \left(\left(x - \frac{2}{3}\right) + \sqrt{\left(x - \frac{2}{3}\right)^2 + \left(\frac{\sqrt{2}}{3}\right)^2} \right) + c$
 $= \frac{1}{\sqrt{3}} \cdot \log \left(\left(x - \frac{2}{3}\right) + \sqrt{x^2 - \frac{4}{3}x + \frac{2}{3}} \right) + c$

$$6. \int \frac{1}{3-10x-25x^2} \cdot dx$$

Solution :

$$\begin{aligned}
 I &= \int \frac{1}{25 \left(\frac{3}{25} - \frac{10}{25} x - x^2 \right)} \cdot dx \\
 &= \int \frac{1}{25 \left[\frac{3}{25} - \left(x^2 + \frac{2}{5} x \right) \right]} \cdot dx \\
 \therefore &\left\{ \left(\frac{1}{2} \text{ coefficient of } x \right)^2 \right. \\
 &= \left. \left(\frac{1}{2} \left(\frac{2}{5} \right) \right)^2 = \left(\frac{1}{5} \right)^2 = \frac{1}{25} \right\} \\
 &= \frac{1}{25} \cdot \int \frac{1}{\frac{3}{25} - \left(x^2 - \frac{2}{5} x + \frac{1}{25} - \frac{1}{25} \right)} \cdot dx \\
 &= \frac{1}{25} \cdot \int \frac{1}{\frac{3}{25} - \left(x^2 - \frac{2}{5} x + \frac{1}{25} \right) + \frac{1}{25}} \cdot dx \\
 &= \frac{1}{25} \cdot \int \frac{1}{\frac{4}{25} - \left(x^2 - \frac{2}{5} x + \frac{1}{25} \right)} \cdot dx \\
 &= \frac{1}{25} \cdot \int \frac{1}{\left(\frac{2}{5} \right)^2 - \left(x - \frac{1}{5} \right)^2} \cdot dx \\
 \therefore &\int \frac{1}{a^2 - x^2} \cdot dx = \frac{1}{2a} \log \left(\frac{a+x}{a-x} \right) + c \\
 I &= \frac{1}{25} \cdot \frac{1}{2 \left(\frac{2}{5} \right)} \cdot \log \left(\frac{\frac{2}{5} + \left(x - \frac{1}{5} \right)}{\frac{2}{5} - \left(x - \frac{1}{5} \right)} \right) + c \\
 &= \frac{1}{5} \cdot \log \left(\frac{1+5x}{3-5x} \right) + c
 \end{aligned}$$

Activity :

$$7. \int \frac{1}{\sqrt{1+x-x^2}} \cdot dx$$

$$\text{Solution : } I = \int \frac{1}{\sqrt{1 - \left(\dots \dots \dots \right)}} \cdot dx$$

$$\begin{aligned}
 \therefore &\left\{ \left(\frac{1}{2} \text{ coefficient of } x \right)^2 \right. \\
 &= \left. \left(\frac{1}{2} (-1) \right)^2 = \left(-\frac{1}{2} \right)^2 = \frac{1}{4} \right\} \\
 &= \int \frac{1}{\sqrt{1 - \left(x^2 - x + \frac{1}{4} - \frac{1}{4} \right)}} \cdot dx \\
 &= \int \frac{1}{\sqrt{1 - \left(\dots \dots \dots \right)}} \cdot dx
 \end{aligned}$$

$$\begin{aligned}
 I &= \int \frac{1}{\sqrt{\left(\frac{\sqrt{5}}{2} \right)^2 - \left(x - \frac{1}{2} \right)^2}} \cdot dx \\
 \therefore &\int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx = \sin^{-1} \left(\frac{x}{a} \right) + c
 \end{aligned}$$

$$\begin{aligned}
 I &= \dots \dots \dots \\
 &= \dots \dots \dots
 \end{aligned}$$

8.
$$\int \frac{\sin 2x}{3 \sin^4 x - 4 \sin^2 x + 1} \cdot dx$$

Solution :
$$I = \int \frac{\sin 2x}{3 (\sin^2 x)^2 - 4 (\sin^2 x) + 1} \cdot dx$$

put $\sin^2 x = t$ $\therefore 2 \sin x \cdot \cos x \cdot dx = 1 \cdot dt$
 $\therefore \sin 2x \cdot dx = 1 \cdot dt$

$$= \int \frac{1}{3t^2 - 4t + 1} \cdot dt$$

$$= \int \frac{1}{3 \left(t^2 - \frac{4}{3}t + \frac{1}{3} \right)} \cdot dt$$

$$\therefore \left\{ \left(\frac{1}{2} \text{ coefficient of } t \right)^2 \right.$$

$$= \left. \left(\frac{1}{2} \left(-\frac{4}{3} \right) \right)^2 = \left(-\frac{2}{3} \right)^2 = \frac{4}{9} \right\}$$

$$I = \frac{1}{3} \cdot \int \frac{1}{t^2 - \frac{4}{3}t + \frac{4}{9} - \frac{4}{9} + \frac{1}{3}} \cdot dt$$

$$= \frac{1}{3} \cdot \int \frac{1}{\left(t^2 - \frac{4}{3}t + \frac{4}{9} \right) - \frac{1}{9}} \cdot dt$$

$$= \frac{1}{3} \cdot \int \frac{1}{\left(t - \frac{2}{3} \right)^2 - \left(\frac{1}{3} \right)^2} \cdot dt$$

$$= \frac{1}{3} \cdot \frac{1}{2 \left(\frac{1}{3} \right)} \cdot \log \left(\frac{\left(t - \frac{2}{3} \right) - \frac{1}{3}}{\left(t - \frac{2}{3} \right) + \frac{1}{3}} \right) + c$$

$$= \frac{1}{2} \cdot \log \left(\frac{3t - 3}{3t - 1} \right) + c$$

$$= \frac{1}{2} \cdot \log \left(\frac{3 \sin^2 x - 3}{3 \sin^2 x - 1} \right) + c$$

$$\therefore \int \frac{\sin 2x}{3 \sin^4 x - 4 \sin^2 x + 1} \cdot dx$$

$$= \frac{1}{2} \cdot \log \left(\frac{3 \sin^2 x - 3}{3 \sin^2 x - 1} \right) + c$$

9.
$$\int \frac{e^{\frac{x}{2}}}{\sqrt{e^{-x} - e^x}} \cdot dx$$

Solution :

$$I = \int \frac{\sqrt{e^x}}{\sqrt{\frac{1}{e^x} - e^x}} \cdot dx$$

$$= \int \frac{\sqrt{e^x}}{\sqrt{\frac{1 - (e^x)^2}{e^x}}} \cdot dx$$

$$= \int \frac{\sqrt{e^x}}{\frac{\sqrt{1 - (e^x)^2}}{\sqrt{e^x}}} \cdot dx$$

$$= \int \frac{\sqrt{e^x} \cdot \sqrt{e^x}}{\sqrt{1 - (e^x)^2}} \cdot dx$$

$$= \int \frac{e^x}{\sqrt{1 - (e^x)^2}} \cdot dx$$

put $e^x = t$

$$\therefore e^x \cdot dx = 1 \cdot dt$$

$$I = \int \frac{1}{\sqrt{1 - t^2}} \cdot dt$$

$$= \sin^{-1}(t) + c$$

$$= \sin^{-1}(e^x) + c$$

$$\therefore \int \frac{e^{\frac{x}{2}}}{\sqrt{e^{-x} - e^x}} \cdot dx = \sin^{-1}(e^x) + c$$

10. $\int(\sqrt{\tan x} + \sqrt{\cot x}) \cdot dx$

Solution : $I = \int\left(\sqrt{\tan x} + \frac{1}{\sqrt{\tan x}}\right) \cdot dx$

$$= \int \frac{\tan x + 1}{\sqrt{\tan x}} \cdot dx$$

put $\sqrt{\tan x} = t \quad \therefore \tan x = t^2 \quad \therefore x = \tan^{-1} t^2$

$$\therefore 1 \cdot dx = \frac{1}{1 + (t^2)^2} \cdot 2t \cdot dt$$

$$\therefore \sec^2 x \cdot dx = 2t \cdot dt$$

$$\therefore dx = \frac{2t}{\sec^2 x} \cdot dx = \frac{2t}{1 + \tan^2 x} \cdot dx = \frac{2t}{1 + t^4} \cdot dt$$

$$= \int \frac{t^2 + 1}{t} \cdot \frac{2t}{1 + t^4} \cdot dt = 2 \int \frac{t^2 + 1}{t^4 + 1} \cdot dt$$

$$= 2 \int \frac{1 + \frac{1}{t^2}}{t^2 + \frac{1}{t^2}} \cdot dt = 2 \int \frac{\left(1 + \frac{1}{t^2}\right)}{\left(t - \frac{1}{t}\right)^2 + 2} \cdot dt$$

put $t - \frac{1}{t} = u \quad \therefore \left[\frac{d}{dt}\left(t - \frac{1}{t}\right) = 1 + \frac{1}{t^2}\right]$

$$\therefore \left(t - \left(-\frac{1}{t^2}\right)\right) dt = 1 \cdot du$$

$$\therefore \left(1 + \frac{1}{t^2}\right) dt = 1 \cdot du$$

$$I = 2 \int \frac{1}{u^2 + 2} \cdot du$$

$$= 2 \int \frac{1}{u^2 + (\sqrt{2})^2} \cdot du$$

$$= 2 \cdot \frac{1}{\sqrt{2}} \cdot \tan^{-1}\left(\frac{u}{\sqrt{2}}\right) + c$$

$$= \sqrt{2} \cdot \tan^{-1}\left(\frac{t - \frac{1}{t}}{\sqrt{2}}\right) + c$$

$$= \sqrt{2} \cdot \tan^{-1}\left(\frac{t^2 - 1}{\sqrt{2} t}\right) + c$$

$$= \sqrt{2} \cdot \tan^{-1}\left(\frac{\tan x - 1}{\sqrt{2} \cdot \sqrt{\tan x}}\right) + c$$

11. $\int \frac{1}{5 - 4 \cos x} \cdot dx$

Solution : put $\tan \frac{x}{2} = t$

$$\therefore dx = \frac{2}{1 + t^2} \cdot dt \quad \text{and} \quad \cos x = \frac{1 - t^2}{1 + t^2}$$

$$I = \int \frac{1 \left(\frac{2}{1 + t^2}\right)}{5 - 4 \left(\frac{1 - t^2}{1 + t^2}\right)} \cdot dt$$

$$= \int \frac{\frac{2}{1 + t^2}}{\frac{5(1 + t^2) - 4(1 - t^2)}{1 + t^2}} \cdot dt$$

$$= \int \frac{2}{5 - 5t^2 - 4 + 4t^2} \cdot dt$$

$$= \int \frac{2}{9t^2 + 1} \cdot dt$$

$$= \int \frac{1}{9 \left(t^2 + \frac{1}{9}\right)} \cdot dt$$

$$= \frac{2}{9} \cdot \int \frac{1}{t^2 + \left(\frac{1}{3}\right)^2} \cdot dt$$

$$= \frac{2}{9} \cdot \frac{1}{\left(\frac{1}{3}\right)} \cdot \tan^{-1}\left(\frac{t}{\left(\frac{1}{3}\right)}\right) + c$$

$$= \frac{2}{3} \cdot \tan^{-1}(2t) + c$$

$$= \frac{2}{3} \cdot \tan^{-1}\left(2 \tan \frac{x}{2}\right) + c$$

$$\therefore \int \frac{1}{5 - 4 \cos x} \cdot dx = \frac{2}{3} \cdot \tan^{-1}\left(2 \tan \frac{x}{2}\right) + c$$

12. $\int \frac{1}{2 - 3 \sin 2x} \cdot dx$

Solution : put $\tan x = t$

$\therefore dx = \frac{1}{1+t^2} \cdot dt$ and $\sin 2x = \frac{2}{1+t^2}$

$$\begin{aligned} I &= \int \frac{1 \left(\frac{1}{1+t^2} \right)}{2 - 3 \left(\frac{2}{1+t^2} \right)} \cdot dt \\ &= \int \frac{\frac{1}{1+t^2}}{2(1+t^2) - 3(2t)} \cdot dt \\ &= \int \frac{1}{2+2t^2-6t} \cdot dt = \int \frac{1}{2(t^2-3t+1)} \cdot dt \\ \therefore &\left\{ \left(\frac{1}{2} \text{ coefficient of } t \right)^2 \right. \\ &= \left. \left(\frac{1}{2}(-3) \right)^2 = \left(-\frac{3}{2} \right)^2 = \frac{9}{4} \right\} \\ &= \frac{1}{2} \cdot \int \frac{1}{t^2 - 3t + \frac{9}{4} - \frac{9}{4} + 1} \cdot dt \\ &= \frac{1}{2} \cdot \int \frac{1}{\left(t - 3t + \frac{9}{4} \right) - \frac{5}{4}} \cdot dt \\ &= \frac{1}{2} \cdot \int \frac{1}{\left(t - \frac{3}{2} \right)^2 - \left(\frac{\sqrt{5}}{2} \right)^2} \cdot dt \\ &= \frac{1}{2} \cdot \frac{1}{2 \left(\frac{\sqrt{5}}{2} \right)} \cdot \log \left(\frac{\left(t - \frac{3}{2} \right) - \frac{\sqrt{5}}{2}}{\left(t - \frac{3}{2} \right) + \frac{\sqrt{5}}{2}} \right) + c \\ &= \frac{1}{2\sqrt{5}} \cdot \log \left(\frac{2t - 3 - \sqrt{5}}{2t - 3 + \sqrt{5}} \right) + c \\ &= \frac{1}{2\sqrt{5}} \cdot \log \left(\frac{2 \tan x - 3 - \sqrt{5}}{2 \tan x - 3 + \sqrt{5}} \right) + c \end{aligned}$$

13. $\int \frac{1}{3 - 2 \sin x + 5 \cos x} \cdot dx$

Solution : put $\tan \frac{x}{2} = t$

$\therefore dx = \frac{2}{1+t^2}$

$\therefore \sin x = \frac{2}{1+t^2} \cdot dt$ and $\cos x = \frac{1-t^2}{1+t^2}$

$$\begin{aligned} I &= \int \frac{1 \left(\frac{2}{1+t^2} \right)}{3 - 2 \left(\frac{2}{1+t^2} \right) + 5 \left(\frac{1-t^2}{1+t^2} \right)} \cdot dt \\ &= \int \frac{\frac{2}{1+t^2}}{3(1+t^2) - 2(2t) + 5(1-t^2)} \cdot dt \\ &= \int \frac{2}{3+3t^2-4t+5-5t^2} \cdot dt \\ &= \int \frac{2}{8-4t-2t^2} \cdot dt \\ &= \int \frac{1}{4-2t-t^2} \cdot dt \\ &= \int \frac{1}{4-(t^2+2t)} \cdot dt \\ &= \int \frac{1}{4-(t^2+2t+1-1)} \cdot dt \\ &= \int \frac{1}{5-(t^2+2t+1)} \cdot dt \\ &= \int \frac{1}{(\sqrt{5})^2 - (t+1)^2} \cdot dt \\ &= \frac{1}{2(\sqrt{5})} \cdot \log \left(\frac{\sqrt{5} + (t+1)}{\sqrt{5} - (t+1)} \right) + c \\ &= \frac{1}{2\sqrt{5}} \cdot \log \left(\frac{\sqrt{5} + 1 + \tan \frac{x}{2}}{\sqrt{5} - 1 - \tan \frac{x}{2}} \right) + c \end{aligned}$$

Activity : 14. $\int \frac{1}{\sin x - \sqrt{3} \cos x} \cdot dx$

Solution : put $\tan \frac{x}{2} = t \quad \therefore dx = \dots\dots\dots$

$\therefore \sin x = \dots\dots\dots$ and $\cos x = \dots\dots\dots$

$$\begin{aligned}
 I &= \int \frac{1 \left(\frac{2}{1+t^2} \right)}{\dots\dots\dots + \sqrt{3} \dots\dots\dots} \cdot dt \\
 &= \int \frac{\frac{2}{1+t^2}}{\dots\dots\dots \frac{1+t^2}{1+t^2}} \cdot dt \\
 &= \int \frac{2}{\dots\dots\dots} \cdot dt \\
 &= \int \frac{2}{\sqrt{3} \left(1 - \left(t^2 - \frac{2}{\sqrt{3}} t \right) \right)} \cdot dt \\
 &= \frac{2}{\sqrt{3}} \cdot \int \frac{1}{1 - \left(t^2 - \frac{2}{\sqrt{3}} t + \frac{1}{3} - \frac{1}{3} \right)} \cdot dt \\
 &= \frac{2}{\sqrt{3}} \cdot \int \frac{1}{1 - \left(t^2 - \frac{2}{\sqrt{3}} t + \frac{1}{3} \right) + \frac{1}{3}} \cdot dt \\
 &= \frac{2}{\sqrt{3}} \cdot \int \frac{1}{\left(\dots\dots \right) - \left(t - \frac{1}{\sqrt{3}} \right)^2} \cdot dt \\
 &= \frac{2}{\sqrt{3}} \cdot \int \frac{1}{\left(\dots \right)^2 - \left(\dots \right)^2} \cdot dt \\
 &= \frac{2}{\sqrt{3}} \cdot \frac{1}{2(\dots)} \cdot \log \left(\dots\dots\dots \right) + c \\
 &= \dots \cdot \log \left(\dots\dots\dots \right) + c \\
 &= \frac{1}{2} \cdot \log \left(\frac{1 + \sqrt{3} \tan \frac{x}{2}}{3 - \sqrt{3} \tan \frac{x}{2}} \right) + c
 \end{aligned}$$

Alternative method :

14. $\int \frac{1}{\sin x - \sqrt{3} \cos x} \cdot dx$

Solution : For any two positive numbers a and b , we can find an angle θ , such that

$\therefore \sin \theta = \frac{a}{\sqrt{a^2 + b^2}}$ and $\cos \theta = \frac{b}{\sqrt{a^2 + b^2}}$

Using this we express $\sin x - \sqrt{3} \cos x$
 $= \sqrt{1+3} (\cos \theta \cdot \sin x - \sin \theta \cdot \cos x)$
 $= 2 \cdot \sin (x - \theta)$
 $= 2 \cdot \sin \left(x - \frac{\pi}{3} \right)$

$\therefore I = \int \frac{1}{2 \cdot \sin \left(x - \frac{\pi}{3} \right)} \cdot dx$
 $= \frac{1}{2} \cdot \int \operatorname{cosec} \left(x - \frac{\pi}{3} \right) \cdot dx$
 $= \frac{1}{2} \cdot \log \left(\operatorname{cosec} \left(x - \frac{\pi}{3} \right) - \cot \left(x - \frac{\pi}{3} \right) \right) + c$
 $= \frac{1}{2} \cdot \log \left(\tan \left(\frac{x}{2} + \frac{\pi}{6} \right) \right) + c$

15. $\int \frac{1}{3 + 2 \sin^2 x + 5 \cos^2 x} \cdot dx$

Solution : Divide Numerator and Denominator by $\cos^2 x$

$$\begin{aligned} I &= \int \frac{\frac{1}{\cos^2 x}}{\frac{3 + 2 \sin^2 x + 5 \cos^2 x}{\cos^2 x}} \cdot dx \\ &= \int \frac{\sec^2 x}{3 \sec^2 x + 2 \tan^2 x + 5} \cdot dx \\ &= \int \frac{\sec^2 x}{3(1 + \tan^2 x) + 2 \tan^2 x + 5} \cdot dx \\ &= \int \frac{\sec^2 x}{5 \tan^2 x + 8} \cdot dx \\ &= \frac{1}{5} \int \frac{\sec^2 x}{\tan^2 x + \frac{8}{5}} \cdot dx \end{aligned}$$

put $\tan x = t \therefore \sec^2 x \cdot dx = 1 \cdot dt$

$$\begin{aligned} I &= \frac{1}{5} \int \frac{1}{t^2 + \frac{8}{5}} \cdot dt \\ &= \frac{1}{5} \int \frac{1}{t^2 + \left(\frac{\sqrt{8}}{\sqrt{5}}\right)^2} \cdot dt \\ &= \frac{1}{5} \cdot \frac{1}{\frac{\sqrt{8}}{\sqrt{5}}} \cdot \tan^{-1} \left(\frac{t}{\frac{\sqrt{8}}{\sqrt{5}}} \right) + c \\ &= \frac{1}{\sqrt{5}} \cdot \frac{1}{2\sqrt{2}} \cdot \tan^{-1} \left(\frac{\sqrt{5} t}{2\sqrt{2}} \right) + c \\ &= \frac{1}{2\sqrt{10}} \cdot \tan^{-1} \left(\frac{\sqrt{5} \tan x}{2\sqrt{2}} \right) + c \\ \therefore \int \frac{1}{3 + 2 \sin^2 x + 5 \cos^2 x} \cdot dx &= \frac{1}{2\sqrt{10}} \cdot \tan^{-1} \left(\frac{\sqrt{5} \tan x}{2\sqrt{2}} \right) + c \end{aligned}$$

16. $\int \frac{\cos \theta}{\cos 3\theta} \cdot d\theta$

Solution : $I = \int \frac{\cos \theta}{4 \cos^3 \theta - 3 \cos \theta} \cdot d\theta$
 $= \int \frac{1}{4 \cos^2 \theta - 3} \cdot d\theta$

Divide Numerator and Denominator by $\cos^2 \theta$

$$\begin{aligned} I &= \int \frac{\frac{1}{\cos^2 \theta}}{\frac{4 \cos^2 \theta - 3}{\cos^2 \theta}} \cdot d\theta \\ &= \int \frac{\sec^2 \theta}{4 - 3 \sec^2 \theta} \cdot d\theta \\ &= \int \frac{\sec^2 \theta}{4 - 3(1 + \tan^2 \theta)} \cdot d\theta \\ &= \int \frac{\sec^2 \theta}{1 - 3 \tan^2 \theta} \cdot d\theta \end{aligned}$$

put $\tan \theta = t \therefore \sec^2 \theta \cdot d\theta = 1 \cdot dt$

$$\begin{aligned} I &= \int \frac{1}{1 - 3t^2} \cdot dt \\ &= \frac{1}{3} \int \frac{1}{\frac{1}{3} - t^2} \cdot dt \\ &= \frac{1}{3} \int \frac{1}{\left(\frac{1}{\sqrt{3}}\right)^2 - t^2} \cdot dt \\ &= \frac{1}{3} \cdot \frac{1}{2\left(\frac{1}{\sqrt{3}}\right)} \cdot \log \left(\frac{\frac{1}{\sqrt{3}} + t}{\frac{1}{\sqrt{3}} - t} \right) + c \\ &= \frac{1}{2\sqrt{3}} \cdot \log \left(\frac{1 + \sqrt{3} t}{1 - \sqrt{3} t} \right) + c \\ &= \frac{1}{2\sqrt{3}} \cdot \log \left(\frac{1 + \sqrt{3} \tan \theta}{1 - \sqrt{3} \tan \theta} \right) + c \\ \therefore \int \frac{\cos \theta}{\cos 3\theta} \cdot d\theta &= \frac{1}{2\sqrt{3}} \cdot \log \left(\frac{1 + \sqrt{3} \tan \theta}{1 - \sqrt{3} \tan \theta} \right) + c \end{aligned}$$

EXERCISE 3.2 (B)

I. Evaluate the following :

1. $\int \frac{1}{4x^2 - 3} \cdot dx$

2. $\int \frac{1}{25 - 9x^2} \cdot dx$

3. $\int \frac{1}{7 + 2x^2} \cdot dx$

4. $\int \frac{1}{\sqrt{3x^2 + 8}} \cdot dx$

5. $\int \frac{1}{\sqrt{11 - 4x^2}} \cdot dx$

6. $\int \frac{1}{\sqrt{2x^2 - 5}} \cdot dx$

7. $\int \sqrt{\frac{9+x}{9-x}} \cdot dx$

8. $\int \sqrt{\frac{2+x}{2-x}} \cdot dx$

9. $\int \sqrt{\frac{10+x}{10-x}} \cdot dx$

10. $\int \frac{1}{x^2 + 8x + 12} \cdot dx$

11. $\int \frac{1}{1 + x - x^2} \cdot dx$

12. $\int \frac{1}{4x^2 - 20x + 17} \cdot dx$

13. $\int \frac{1}{5 - 4x - 3x^2} \cdot dx$

14. $\int \frac{1}{\sqrt{3x^2 + 5x + 7}} \cdot dx$

15. $\int \frac{1}{\sqrt{x^2 + 8x - 20}} \cdot dx$

16. $\int \frac{1}{\sqrt{8 - 3x + 2x^2}} \cdot dx$

17. $\int \frac{1}{\sqrt{(x-3)(x+2)}} \cdot dx$

18. $\int \frac{1}{4 + 3 \cos^2 x} \cdot dx$

19. $\int \frac{1}{\cos 2x + 3 \sin^2 x} \cdot dx$

20. $\int \frac{\sin x}{\sin 3x} \cdot dx$

II. Integrate the following functions w. r. t. x :

1. $\int \frac{1}{3 + 2 \sin x} \cdot dx$

2. $\int \frac{1}{4 - 5 \cos x} \cdot dx$

3. $\int \frac{1}{2 + \cos x - \sin x} \cdot dx$

4. $\int \frac{1}{3 + 2 \sin x - \cos x} \cdot dx$

5. $\int \frac{1}{3 - 2 \cos 2x} \cdot dx$

6. $\int \frac{1}{2 \sin 2x - 3} \cdot dx$

7. $\int \frac{1}{3 + 2 \sin 2x + 4 \cos 2x} \cdot dx$

8. $\int \frac{1}{\cos x - \sin x} \cdot dx$

9. $\int \frac{1}{\cos x - \sqrt{3} \sin x} \cdot dx$

3.2.6 Integral of the form $\int \frac{px+q}{ax^2+bx+c} \cdot dx$ and $\int \frac{px+q}{\sqrt{ax^2+bx+c}} \cdot dx$

The integral of the form $\int \frac{px+q}{ax^2+bx+c} \cdot dx$ is evaluated by expressing the integral in the form

$$\int \frac{A \cdot \frac{d}{dx}(ax^2+bx+c)}{ax^2+bx+c} \cdot dx + \int \frac{B}{ax^2+bx+c} \cdot dx \text{ for some constants } A \text{ and } B.$$

The numerator, $px+q = A \cdot \frac{d}{dx}(ax^2+bx+c) + B$

$$\text{i.e. } Nr = A \cdot \frac{d}{dx} Dr + B$$

The first integral is evaluated by putting $ax^2+bx+c = t$

The Second integral is evaluated by expressing the integrand in the form either

$$\frac{1}{A^2+t^2} \text{ or } \frac{1}{t^2-A^2} \text{ or } \frac{1}{A^2-t^2} \text{ and applying the methods discussed previously.}$$

The integral of the form $\int \frac{px+q}{\sqrt{ax^2+bx+c}} \cdot dx$ is evaluated by expressing the integral in the form

$$\int \frac{A \cdot \frac{d}{dx}(ax^2+bx+c)}{\sqrt{ax^2+bx+c}} \cdot dx + \int \frac{B}{\sqrt{ax^2+bx+c}} \cdot dx \text{ for constants } A \text{ and } B.$$

The numerator, $px+q = A \cdot \frac{d}{dx}(ax^2+bx+c) + B$

The first integral is evaluated by putting $ax^2+bx+c = t$

The second integral is evaluated by expressing the integrand in the form either

$$\frac{1}{\sqrt{A^2+t^2}} \text{ or } \frac{1}{\sqrt{t^2-A^2}} \text{ or } \frac{1}{\sqrt{A^2-t^2}} \text{ and applying the methods which discussed previously.}$$

SOLVED EXAMPLES

1. $\int \frac{2x - 3}{3x^2 + 4x + 5} \cdot dx$

Solution : $2x - 3 = A \cdot \frac{d}{dx} (3x^2 + 4x + 5) + B$

$2x - 3 = A(6x + 4) + B$

$= (6A)x + (4A + B)$

comparing the sides/ the co-efficients of like variables and constants

$6A = 2$ and $4A + B = -3$

$\Rightarrow A = \frac{1}{3}$ and $B = -\frac{13}{3}$

$= \int \frac{\frac{1}{3} \cdot \frac{d}{dx} (3x^2 + 4x + 5) + \left(-\frac{13}{3}\right)}{3x^2 + 4x + 5} \cdot dx$

$= \frac{1}{3} \int \frac{\frac{d}{dx} (3x^2 + 4x + 5)}{3x^2 + 4x + 5} \cdot dx - \frac{13}{3} \int \frac{1}{3x^2 + 4x + 5} \cdot dx$

$= \frac{1}{3} \int \frac{6x + 4}{3x^2 + 4x + 5} \cdot dx - \frac{13}{3} \int \frac{1}{3x^2 + 4x + 5} \cdot dx$

$= I_1 - I_2 \dots \dots (i)$

$\therefore I_1 = \frac{1}{3} \int \frac{6x + 4}{3x^2 + 4x + 5} \cdot dx$

put $3x^2 + 4x + 5 = t$

$\therefore (6x + 4) \cdot dx = 1 \cdot dt$

$I_1 = \frac{1}{3} \int \frac{1}{t} \cdot dt$

$= \frac{1}{3} \cdot \log(t) + c_1$

$= \frac{1}{3} \cdot \log(3x^2 + 4x + 5) + c_1 \dots \dots (ii)$

$\therefore I_2 = \frac{13}{3} \int \frac{1}{3x^2 + 4x + 5} \cdot dx$
 $= \frac{13}{3} \cdot \frac{1}{3} \int \frac{1}{x^2 + \frac{4}{3}x + \frac{5}{3}} \cdot dx$

$\therefore \left\{ \left(\frac{1}{2} \text{ coefficient of } t \right)^2 \right.$
 $\left. = \left(\frac{1}{2} \left(\frac{4}{3} \right) \right)^2 = \left(\frac{2}{3} \right)^2 = \frac{4}{9} \right\}$

$I_2 = \frac{13}{9} \int \frac{1}{x^2 + \frac{4}{3}x + \frac{4}{9} - \frac{4}{9} + \frac{5}{3}} \cdot dx$

$= \frac{13}{9} \int \frac{1}{x^2 + \frac{4}{3}x + \frac{4}{9} + \frac{11}{9}} \cdot dx$

$= \frac{13}{9} \int \frac{1}{\left(x + \frac{2}{3}\right)^2 + \left(\frac{\sqrt{11}}{3}\right)^2} \cdot dx$

$\therefore \int \frac{1}{X^2 + A^2} \cdot dx = \frac{1}{A} \tan^{-1} \left(\frac{X}{A} \right) + c$

$= \frac{13}{9} \cdot \frac{1}{\frac{\sqrt{11}}{3}} \cdot \tan^{-1} \left(\frac{x + \frac{2}{3}}{\frac{\sqrt{11}}{3}} \right) + c_1$

$I_2 = \frac{13}{3\sqrt{11}} \cdot \tan^{-1} \left(\frac{3x + 2}{\sqrt{11}} \right) + c_2 \dots \dots (iii)$

thus, from (i), (ii) and (iii)

$\therefore \int \frac{2x - 3}{3x^2 + 4x + 5} \cdot dx$

$= \frac{1}{3} \cdot \log(3x^2 + 4x + 5) - \frac{13}{3\sqrt{11}} \cdot \tan^{-1} \left(\frac{3x + 2}{\sqrt{11}} \right) + c$

$(\because c_1 + c_2 = c)$

$$2. \int \sqrt{\frac{x-5}{x-7}} \cdot dx$$

$$\text{Solution : } I = \int \sqrt{\frac{(x-5) \cdot (x-5)}{(x-7) \cdot (x-5)}} \cdot dx = \int \sqrt{\frac{(x-5)^2}{x^2 - 12x + 35}} \cdot dx$$

$$\therefore x-5 = A \cdot \frac{d}{dx}(x^2 - 12x + 35) + B$$

$$x-5 = A(2x-12) + B$$

$$= (2A)x + (-12A + B)$$

comparing, the co-efficients of like variables and constants

$$2A = 1 \quad \text{and} \quad -12A + B = -5$$

$$\Rightarrow A = \frac{1}{2} \quad \text{and} \quad B = 1$$

$$\begin{aligned} I &= \int \frac{\frac{1}{2} \cdot \frac{d}{dx}(x^2 - 12x + 35) + (1)}{\sqrt{x^2 - 12x + 35}} \cdot dx \\ &= \frac{1}{2} \int \frac{\frac{d}{dx}(x^2 - 12x + 35)}{\sqrt{x^2 - 12x + 35}} \cdot dx + \int \frac{1}{\sqrt{x^2 - 12x + 35}} \cdot dx \\ &= I_1 + I_2 \quad \dots \dots \text{(i)} \end{aligned}$$

$$\therefore I_1 = \frac{1}{2} \int \frac{2x-12}{\sqrt{x^2-12x+35}} \cdot dx$$

$$\text{put } x^2 - 12x + 35 = t$$

$$\therefore (2x-12) \cdot dx = 1 \cdot dt$$

$$I_1 = \frac{1}{2} \int \frac{1}{\sqrt{t}} \cdot dt$$

$$= \int \frac{1}{2\sqrt{t}} \cdot dt$$

$$= \sqrt{t} + c_1$$

$$= \sqrt{x^2 - 12x + 35} + c_1 \quad \dots \dots \text{(ii)}$$

$$\begin{aligned} \therefore I_2 &= \int \frac{1}{\sqrt{x^2 - 12x + 35}} \cdot dx \\ &= \int \frac{1}{\sqrt{x^2 - 12x + 36 - 1}} \cdot dx \\ &= \int \frac{1}{\sqrt{(x-6)^2 - (1)^2}} \cdot dx \end{aligned}$$

$$\therefore \int \frac{1}{\sqrt{X^2 - A^2}} \cdot dx = \log(X + \sqrt{X^2 - A^2}) + c$$

$$I_2 = \log((x-6) + \sqrt{(x-6)^2 - 1}) + c_2$$

$$= \log((x-6) + \sqrt{x^2 - 12x + 35}) + c_2$$

$$\dots \dots \text{(iii)}$$

Thus, from (i), (ii) and (iii)

$$\int \sqrt{\frac{x-5}{x-7}} \cdot dx$$

$$= \sqrt{x^2 - 12x + 35} + \log((x-6) + \sqrt{x^2 - 12x + 35}) + c$$

$$(c_1 + c_2 = c)$$

EXERCISE 3.2 (C)

I. Evaluate :

1. $\int \frac{3x + 4}{x^2 + 6x + 5} \cdot dx$
2. $\int \frac{2x + 1}{x^2 + 4x - 5} \cdot dx$
3. $\int \frac{2x + 3}{2x^2 + 3x - 1} \cdot dx$
4. $\int \frac{3x + 4}{\sqrt{2x^2 + 2x + 1}} \cdot dx$
5. $\int \frac{7x + 3}{\sqrt{3 + 2x - x^2}} \cdot dx$
6. $\int \sqrt{\frac{x - 7}{x - 9}} \cdot dx$
7. $\int \sqrt{\frac{9 - x}{x}} \cdot dx$
8. $\int \frac{3 \cos x}{4 \sin^2 x + 4 \sin x - 1} \cdot dx$
9. $\int \sqrt{\frac{e^{3x} - e^{2x}}{e^x + 1}} \cdot dx$

3.3 Integration by parts :

This method is useful when the integrand is expressed as a product of two different types of functions; one of which can be differentiated and the other can be integrated conveniently.

The following theorem gives the rule of integration by parts.

3.3.1 Theorem : If u and v are two differentiable functions of x then

$$\int u \cdot v \cdot dx = u \cdot \int v \cdot dx - \int \left(\frac{d}{dx} \cdot u \right) (\int v \cdot dx) \cdot dx$$

Proof : Let $\int v \cdot dx = w \dots (i) \Rightarrow v = \frac{dw}{dx} \dots (ii)$

Consider, $\frac{d}{dx} (u \cdot w) = u \cdot \frac{d}{dx} w + w \cdot \frac{d}{dx} u$

$$= u \cdot v + w \cdot \frac{du}{dx}$$

By definition of integration

$$\begin{aligned} u \cdot w &= \int \left[u \cdot v + w \cdot \frac{du}{dx} \right] \cdot dx \\ &= \int u \cdot v \cdot dx + \int w \cdot \frac{du}{dx} \cdot dx \\ &= \int u \cdot v \cdot dx + \int \frac{du}{dx} \cdot w \cdot dx \end{aligned}$$

$$\therefore u \cdot \int v \cdot dx = \int u \cdot v \cdot dx + \int \frac{du}{dx} \cdot \int v \cdot dx \cdot dx$$

$$\therefore \int u \cdot v \cdot dx = u \cdot \int v \cdot dx - \int \left(\frac{d}{dx} \cdot u \right) (\int v \cdot dx) \cdot dx$$

In short, $\int u \cdot v = u \cdot \int v - \int (u' \int v)$

For example : $\int x \cdot e^x \cdot dx = x \int e^x \cdot dx - \int \left(\frac{d(x)}{dx} \cdot \int e^x \cdot dx \right) \cdot dx$

$$= x \cdot e^x - \int (1) \cdot e^x \cdot dx$$

$$= x \cdot e^x - \int e^x \cdot dx$$

$$= x \cdot e^x - e^x + c$$

now let us reverse the choice of u and v

$$\therefore \int e^x \cdot x \cdot dx = e^x \cdot \int x^1 \cdot dx - \int \frac{d}{dx} \cdot e^x \int x \cdot dx \cdot dx$$

$$= e^x \cdot \frac{x^2}{2} - \int e^x \cdot \frac{x^2}{2} \cdot dx$$

$$= \frac{1}{2} \cdot e^x \cdot x^2 - \frac{1}{2} \cdot \int e^x \cdot x^2 \cdot dx$$

We arrive at an integral $\int e^x \cdot x^2 \cdot dx$ which is more difficult, but it helps to get $\int e^x \cdot x^2 \cdot dx$

Thus it is essential to make a proper choice of the first function and the second function. The first function to be selected will be the one, which comes first in the order of **LIATE**.

L **Logarithmic function.**
I **Inverse trigonometric function.**
A **Algebraic function.**
T **Trigonometric function.**
E **Exponential function.**

For example : $\int \sin x \cdot x \cdot dx$


$$= \int x \cdot \sin x \cdot dx \quad \dots \dots \text{by LIATE}$$

$$= x \cdot \int \sin x \cdot dx - \int \frac{d}{dx} \cdot x \cdot \int \sin x \cdot dx \cdot dx$$

$$= x \cdot (-\cos x) - \int (1) \cdot (-\cos x) \cdot dx$$

$$= -x \cdot \cos x + \int \cos x \cdot dx$$

$$= -x \cdot \cos x + \sin x + c$$



SOLVED EXAMPLES

1. $\int x^2 \cdot 5^x \cdot dx$

Solution :

$$\begin{aligned}
 I &= x^2 \cdot \int 5^x \cdot dx - \int \frac{d}{dx} \cdot x^2 \cdot \int 5^x \cdot dx \cdot dx \\
 &= x^2 \cdot 5^x \cdot \frac{1}{\log 5} - \int 2x \cdot 5^x \cdot \frac{1}{\log 5} \cdot dx \\
 &= \frac{1}{\log 5} \cdot x^2 \cdot 5^x - \frac{2}{\log 5} \left\{ x \cdot \int 5^x \cdot dx - \int \frac{d}{dx} \cdot x \cdot \int 5^x \cdot dx \cdot dx \right\} \\
 &= \frac{1}{\log 5} \cdot x^2 \cdot 5^x - \frac{2}{\log 5} \left\{ x \cdot 5^x \cdot \frac{1}{\log 5} - \int (1) \left(5^x \cdot \frac{1}{\log 5} \right) \cdot dx \right\} \\
 &= \frac{1}{\log 5} \cdot x^2 \cdot 5^x - \frac{2}{\log 5} \left\{ \frac{1}{\log 5} \cdot x \cdot 5^x - \int \frac{1}{\log 5} \cdot 5^x \cdot dx \right\} \\
 &= \frac{1}{\log 5} \cdot x^2 \cdot 5^x - \frac{2}{\log 5} \left\{ \frac{1}{\log 5} \cdot x \cdot 5^x - \frac{1}{\log 5} \cdot 5^x \cdot \frac{1}{\log 5} \right\} + c \\
 &= \frac{1}{\log 5} \cdot x^2 \cdot 5^x - \frac{2}{(\log 5)^2} \cdot x \cdot 5^x + \frac{2}{(\log 5)^3} \cdot 5^x + c \\
 \therefore \int x^2 \cdot 5^x \cdot dx &= \frac{5^x}{\log 5} \cdot \left\{ x^2 - \frac{2x}{\log 5} + \frac{2}{(\log 5)^2} \right\} + c
 \end{aligned}$$

2. $\int x \cdot \tan^{-1} x \cdot dx$

Solution :

$$\begin{aligned}
 I &= \int (\tan^{-1} x) \cdot x \cdot dx \quad \dots \dots \text{by LIATE} \\
 &= \tan^{-1} x \cdot \int x \cdot dx - \int \frac{d}{dx} \cdot \tan^{-1} x \cdot \int x \cdot dx \cdot dx \\
 &= \tan^{-1} x \cdot \frac{x^2}{2} - \int \frac{1}{1+x^2} \cdot \frac{x^2}{2} \cdot dx \\
 &= \frac{1}{2} x^2 \cdot \tan^{-1} x - \frac{1}{2} \int \frac{x^2}{1+x^2} \cdot dx \\
 &= \frac{1}{2} x^2 \cdot \tan^{-1} x - \frac{1}{2} \int \frac{1+x^2-1}{1+x^2} \cdot dx \\
 &= \frac{1}{2} x^2 \cdot \tan^{-1} x - \frac{1}{2} \int \left[1 - \frac{1}{1+x^2} \right] \cdot dx \\
 &= \frac{1}{2} x^2 \cdot \tan^{-1} x - \frac{1}{2} [x - \tan^{-1} x] + c \\
 \therefore \int x \cdot \tan^{-1} x \cdot dx &= \frac{1}{2} x^2 \cdot \tan^{-1} x - \frac{1}{2} x + \frac{1}{2} \tan^{-1} x + c
 \end{aligned}$$

3. $\int \frac{x}{1 - \sin x} \cdot dx$

Solution : $I = \int \frac{x}{1 - \sin x} \cdot \frac{(1 + \sin x)}{(1 + \sin x)} \cdot dx$

$$= \int \frac{x(1 + \sin x)}{1 - \sin^2 x} \cdot dx = \int \frac{x(1 + \sin x)}{\cos^2 x} \cdot dx = \int x \cdot \left(\frac{1}{\cos^2 x} + \frac{\sin x}{\cos^2 x} \right) \cdot dx$$

$$= \int x \cdot (\sec^2 x + \sec x \cdot \tan x) \cdot dx$$

$$= \int x \cdot \sec^2 x \cdot dx + \int x \cdot \sec x \cdot \tan x \cdot dx$$

$$= \left(x \cdot \int \sec^2 x \cdot dx - \int \frac{d}{dx} x \cdot \int \sec^2 x \cdot dx \cdot dx \right) + \left(x \cdot \int \sec x \cdot \tan x \cdot dx - \int \frac{d}{dx} x \cdot \int \sec x \cdot \tan x \cdot dx \cdot dx \right)$$

$$= x \cdot \tan x - \int (1) \cdot \tan x \cdot dx + x \cdot \sec x - \int (1) \cdot \sec x \cdot dx$$

$$= x \cdot \tan x - \log (\sec x) + x \cdot \sec x - \log (\sec x + \tan x) + c$$

$$= x \cdot (\sec x + \tan x) - \log (\sec x) - \log (\sec x + \tan x) + c$$

$$\therefore \int \frac{x}{1 - \sin x} \cdot dx = x \cdot (\sec x + \tan x) - \log [(\sec x) (\sec x + \tan x)] + c$$

4. $\int e^{2x} \cdot \sin 3x \cdot dx$

Solution : $I = \int e^{2x} \cdot \sin 3x \cdot dx$

Here we use repeated integration by parts.

To evaluate $\int e^{ax} \cdot \sin (bx + c) \cdot dx$; $\int e^{ax} \cdot \cos (bx + c) \cdot dx$ any function can be taken as a first function.

$$I = e^{2x} \cdot \int \sin 3x \cdot dx - \int \frac{d}{dx} \cdot e^{2x} \cdot \int \sin 3x \cdot dx \cdot dx$$

$$= e^{2x} \cdot \left(-\cos 3x \cdot \frac{1}{3} \right) - \int e^{2x} \cdot 2 \cdot \left(-\cos 3x \cdot \frac{1}{3} \right) \cdot dx$$

$$= -\frac{1}{3} \cdot e^{2x} \cdot \cos 3x + \frac{2}{3} \int e^{2x} \cdot \cos 3x \cdot dx$$

$$= -\frac{1}{3} \cdot e^{2x} \cdot \cos 3x + \frac{2}{3} \left(e^{2x} \cdot \int \cos 3x \cdot dx - \int \frac{d}{dx} \cdot e^{2x} \cdot \int \cos 3x \cdot dx \cdot dx \right)$$

$$= -\frac{1}{3} \cdot e^{2x} \cdot \cos 3x + \frac{2}{3} \left[e^{2x} \cdot \left(\sin 3x \cdot \frac{1}{3} \right) - \int e^{2x} \cdot 2 \cdot \left(\sin 3x \cdot \frac{1}{3} \right) \cdot dx \right]$$

$$= -\frac{1}{3} \cdot e^{2x} \cdot \cos 3x + \frac{2}{9} \cdot e^{2x} \cdot \sin 3x - \frac{4}{9} \cdot \int e^{2x} \cdot \sin 3x \cdot dx$$

$$I = -\frac{1}{3} \cdot e^{2x} \cdot \cos 3x + \frac{2}{9} \cdot e^{2x} \cdot \sin 3x - \frac{4}{9} \cdot I$$

$$I + \frac{4}{9} \cdot I = \frac{e^{2x}}{9} [-3 \cos 3x + 2 \sin 3x] + c$$

$$= \frac{e^{2x}}{13} [2 \sin 3x - 3 \cos 3x] + c$$

$$\frac{13}{9} \cdot I = \frac{e^{2x}}{9} [2 \sin 3x - 3 \cos 3x] + c$$

$$\therefore \int e^{2x} \cdot \sin 3x \cdot dx = \frac{e^{2x}}{13} [2 \sin 3x - 3 \cos 3x] + c$$



Activity :

Prove the following results.

$$(i) \int e^{ax} \cdot \sin (bx + c) \cdot dx = \frac{e^{ax}}{a^2 + b^2} \cdot [a \sin (bx + c) + b \cos (bx + c)] + c$$

$$(ii) \int e^{ax} \cdot \cos (bx + c) \cdot dx = \frac{e^{ax}}{a^2 + b^2} \cdot [a \cos (bx + c) - b \sin (bx + c)] + c$$

$$5. \int \left[\log (\log x) + \frac{1}{(\log x)^2} \right] \cdot dx$$

Solution :

$$\begin{aligned} I &= \int \log (\log x) \cdot 1 \cdot dx + \int \frac{1}{(\log x)^2} \cdot dx \\ &= \log (\log x) \cdot \int 1 \cdot dx - \int \frac{d}{dx} \cdot \log (\log x) \int 1 \cdot dx + \int \frac{1}{(\log x)^2} \cdot dx \\ &= \log (\log x) \cdot x - \int \frac{1}{\log x} \cdot \frac{1}{x} \cdot (x) \cdot dx + \int \frac{1}{(\log x)^2} \cdot dx \\ &= \log (\log x) \cdot x - \int \frac{1}{\log x} \cdot dx + \int \frac{1}{(\log x)^2} \cdot dx \\ &= \log (\log x) \cdot x - \int (\log x)^{-1} \cdot 1 \cdot dx + \int \frac{1}{(\log x)^2} \cdot dx \\ &= \log (\log x) \cdot x - \left\{ (\log x)^{-1} \cdot \int 1 \cdot dx + \int \frac{d}{dx} \cdot (\log x)^{-1} \cdot \int 1 \cdot dx \cdot dx \right\} + \int \frac{1}{(\log x)^2} \cdot dx \\ &= \log (\log x) \cdot x - \left\{ (\log x)^{-1} \cdot x - \int -1 (\log x)^{-2} \cdot \frac{1}{x} \cdot x \cdot dx \right\} + \int \frac{1}{(\log x)^2} \cdot dx \\ &= \log (\log x) \cdot x - (\log x)^{-1} \cdot x - \int (\log x)^{-2} \cdot dx + \int \frac{1}{(\log x)^2} \cdot dx \\ &= x \cdot \log (\log x) - \frac{x}{\log x} - \int \frac{1}{(\log x)^2} \cdot dx + \int \frac{1}{(\log x)^2} \cdot dx \end{aligned}$$

$$\therefore \int \left[\log (\log x) + \frac{1}{(\log x)^2} \right] \cdot dx = x \cdot \log (\log x) - \frac{x}{\log x} + c$$

Note that :

To evaluate the integrals of type $\int \sin^{-1} x \cdot dx$; $\int \tan^{-1} x \cdot dx$; $\int \sec^{-1} x \cdot dx$; $\int \log x \cdot dx$, take the second function (v) to be 1 and then apply integration by parts.

$$\int \sqrt{a^2 - x^2} \cdot dx ; \int \sqrt{a^2 + x^2} \cdot dx ; \int \sqrt{x^2 - a^2} \cdot dx$$

6. $\int \sqrt{a^2 - x^2} \cdot dx$

Solution : Let $I = \int \sqrt{a^2 - x^2} \cdot 1 \cdot dx$

$$= \sqrt{a^2 - x^2} \cdot \int 1 \cdot dx - \int \frac{d}{dx} \cdot \sqrt{a^2 - x^2} \cdot \int 1 \cdot dx \cdot dx$$

$$= \sqrt{a^2 - x^2} \cdot x - \int \frac{1}{2\sqrt{a^2 - x^2}} (-2x) \cdot (x) \cdot dx$$

$$= \sqrt{a^2 - x^2} \cdot x + \int \frac{x^2}{\sqrt{a^2 - x^2}} \cdot dx$$

$$= \sqrt{a^2 - x^2} \cdot x + \int \frac{a^2 - (a^2 - x^2)}{\sqrt{a^2 - x^2}} \cdot dx$$

$$= \sqrt{a^2 - x^2} \cdot x + \int \left[\frac{a^2}{\sqrt{a^2 - x^2}} - \frac{(a^2 - x^2)}{\sqrt{a^2 - x^2}} \right] \cdot dx$$

$$= x \cdot \sqrt{a^2 - x^2} + a^2 \int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx - \int \sqrt{a^2 - x^2} \cdot dx$$

$$I = x \cdot \sqrt{a^2 - x^2} + a^2 \int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx - I$$

$$\therefore I + I = x \cdot \sqrt{a^2 - x^2} + a^2 \cdot \sin^{-1} \left(\frac{x}{a} \right) + c$$

$$\therefore I = \frac{x}{2} \cdot \sqrt{a^2 - x^2} + \frac{a^2}{2} \cdot \sin^{-1} \left(\frac{x}{a} \right) + c$$

$$\therefore \int \sqrt{a^2 - x^2} \cdot dx = \frac{x}{2} \cdot \sqrt{a^2 - x^2} + \frac{a^2}{2} \cdot \sin^{-1} \left(\frac{x}{a} \right) + c$$

e.g. $\int \sqrt{9 - x^2} \cdot dx = \frac{x}{2} \cdot \sqrt{9 - x^2} + \frac{9}{2} \cdot \sin^{-1} \left(\frac{x}{3} \right) + c$

with reference to the above example solve these :

7. $\int \sqrt{a^2 + x^2} \cdot dx = \frac{x}{2} \cdot \sqrt{x^2 + a^2} + \frac{a^2}{2} \cdot \log \left(x + \sqrt{x^2 + a^2} \right) + c$

8. $\int \sqrt{x^2 - a^2} \cdot dx = \frac{x}{2} \cdot \sqrt{x^2 - a^2} - \frac{a^2}{2} \cdot \log \left(x + \sqrt{x^2 - a^2} \right) + c$

9. $\int x \cdot \sin^{-1} x \cdot dx$

Solution : I = $\int \sin^{-1} x \cdot x \cdot dx$ by LIATE

$$\begin{aligned} &= \sin^{-1} x \cdot \int x \cdot dx - \int \frac{d}{dx} \cdot \sin^{-1} x \cdot \int x \cdot dx \cdot dx \\ &= \sin^{-1} x \cdot \frac{x^2}{2} - \int \frac{1}{\sqrt{1-x^2}} \cdot \frac{x^2}{2} \cdot dx \\ &= \frac{1}{2} x^2 \cdot \sin^{-1} x - \frac{1}{2} \int \frac{x^2}{\sqrt{1-x^2}} \cdot dx \\ &= \frac{1}{2} x^2 \cdot \sin^{-1} x - \frac{1}{2} \int \frac{1 - (1-x^2)}{\sqrt{1-x^2}} \cdot dx \\ &= \frac{1}{2} x^2 \cdot \sin^{-1} x - \frac{1}{2} \int \left[\frac{1}{\sqrt{1-x^2}} - \frac{(1-x^2)}{\sqrt{1-x^2}} \right] \cdot dx \\ &= \frac{1}{2} x^2 \cdot \sin^{-1} x - \frac{1}{2} \int \frac{dx}{\sqrt{1-x^2}} + \frac{1}{2} \int \sqrt{1-x^2} \cdot dx \\ &= \frac{1}{2} x^2 \cdot \sin^{-1} x - \frac{1}{2} \sin^{-1} x + \frac{1}{2} \left[\frac{x}{2} \sqrt{1-x^2} + \frac{1}{2} \sin^{-1}(x) \right] + c \\ &= \frac{1}{2} x^2 \cdot \sin^{-1} x + \frac{1}{4} x \sqrt{1-x^2} - \frac{1}{4} \sin^{-1} x + c \end{aligned}$$

$$\therefore \int x \cdot \sin^{-1} x \cdot dx = \frac{1}{2} x^2 \cdot \sin^{-1} x + \frac{1}{4} x \sqrt{1-x^2} - \frac{1}{4} \sin^{-1} x + c$$

Activity :

10. $\int \cos^{-1} \sqrt{x} \cdot dx$

Solution : put $\sqrt{x} = t$

$$\therefore x = t^2$$

differentiating w.r.t. x

$$\therefore 1 \cdot dx = 2t \cdot dt$$

$$I = \int \cos^{-1} t \cdot 2t \cdot dt$$

refer previous (example no. 9) example and solve it.

11. $\int \sqrt{4 + 3x - 2x^2} \cdot dx$

Solution : $I = \int \sqrt{4 - 2x^2 + 3x} \cdot dx$

$$= \int \sqrt{4 - 2 \left(x^2 - \frac{3}{2}x \right)} \cdot dx$$

$$= \int \sqrt{2} \cdot \sqrt{2 - \left(x^2 - \frac{3}{2}x \right)} \cdot dx$$

$$\therefore \left\{ \left(\frac{1}{2} \text{ coefficient of } x \right)^2 = \left[\frac{1}{2} \left(-\frac{3}{2} \right) \right]^2 = \left(-\frac{3}{4} \right)^2 = \frac{9}{16} \right\}$$

$$I = \sqrt{2} \cdot \int \sqrt{2 - \left(x^2 - \frac{3}{2}x + \frac{9}{16} - \frac{9}{16} \right)} \cdot dx$$

$$= \sqrt{2} \cdot \int \sqrt{2 - \left(x^2 - \frac{3}{2}x + \frac{9}{16} \right) + \frac{9}{16}} \cdot dx$$

$$= \sqrt{2} \cdot \int \sqrt{\left(\frac{\sqrt{41}}{4} \right)^2 - \left(x - \frac{3}{4} \right)^2} \cdot dx$$

$$\therefore \int \sqrt{a^2 - x^2} \cdot dx = \frac{x}{2} \cdot \sqrt{a^2 - x^2} + \frac{a^2}{2} \cdot \sin^{-1} \left(\frac{x}{a} \right) + c$$

$$= \sqrt{2} \cdot \left[\frac{\left(x - \frac{3}{4} \right)}{2} \cdot \sqrt{\left(\frac{\sqrt{41}}{4} \right)^2 - \left(x - \frac{3}{4} \right)^2} + \frac{\left(\frac{\sqrt{41}}{4} \right)^2}{2} \cdot \sin^{-1} \left(\frac{x - \frac{3}{4}}{\frac{\sqrt{41}}{4}} \right) \right] + c$$

$$= \sqrt{2} \left[\frac{4x - 3}{8} \cdot \sqrt{2 + \frac{3}{2}x - x^2} + \frac{41}{32} \cdot \sin^{-1} \left(\frac{4x - 3}{\sqrt{41}} \right) \right] + c$$

$$\therefore \int \sqrt{4 + 3x - 2x^2} \cdot dx = \frac{4x - 3}{8} \cdot \sqrt{4 + 3x - 2x^2} + \frac{41}{16\sqrt{2}} \cdot \sin^{-1} \left(\frac{4x - 3}{\sqrt{41}} \right) + c$$

Note that :

3.3.2 :

To evaluate the integral of type $\int (px + q) \sqrt{ax^2 + bx + c} \cdot dx$

we express the term $px + q = A \cdot \frac{d}{dx} (ax^2 + bx + c) + B \dots$ for constants A, B .

Then the integral will be evaluated by the usual known methods.



3.3.3 Integral of the type $\int e^x [f(x) + f'(x)] \cdot dx = e^x \cdot f(x) + c$

Let $e^x \cdot f(x) = t$

Differentiating w. r. t. x

$$\left[e^x [f'(x) + f(x)] \right] = \frac{dt}{dx}$$

$$e^x [f(x) + f'(x)] = \frac{dt}{dx}$$

By definition of integration,

$$\therefore \int e^x [f(x) + f'(x)] \cdot dx = t + c$$

$$\therefore \int e^x [f(x) + f'(x)] \cdot dx = e^x \cdot f(x) + c$$

e.g. $\int e^x [\tan x + \sec^2 x] \cdot dx = e^x \cdot \tan x + c$

$$\left(\because \frac{d}{dx} \tan x = \sec^2 x \right)$$



SOLVED EXAMPLES

1. $\int e^x \left(\frac{2 + \sin 2x}{1 + \cos 2x} \right) \cdot dx$

Solution :

$$\begin{aligned} I &= \int e^x \left(\frac{2 + 2 \sin x \cdot \cos x}{2 \cdot \cos^2 x} \right) \cdot dx \\ &= \int e^x \left(\frac{1}{\cos^2 x} + \frac{\sin x \cdot \cos x}{\cos^2 x} \right) \cdot dx \\ &= \int e^x [\sec^2 x + \tan x] \cdot dx \\ &= \int e^x [\tan x + \sec^2 x] \cdot dx \\ \therefore f(x) = \tan x &\Rightarrow f'(x) = \sec^2 x \\ \therefore \int e^x [f(x) + f'(x)] \cdot dx &= e^x \cdot f(x) + c \end{aligned}$$

$$I = e^x \cdot \tan x + c$$

$$\therefore \int e^x \left(\frac{2 + \sin 2x}{1 + \cos 2x} \right) \cdot dx = e^x \cdot \tan x + c$$

2. $\int e^x \left[\frac{x+2}{(x+3)^2} \right] \cdot dx$

Solution :

$$\begin{aligned} I &= \int e^x \left[\frac{x+3-1}{(x+3)^2} \right] \cdot dx \\ &= \int e^x \left[\frac{x+3}{(x+3)^2} + \frac{-1}{(x+3)^2} \right] \cdot dx \\ &= \int e^x \left[\frac{1}{x+3} + \frac{-1}{(x+3)^2} \right] \cdot dx \\ \therefore f(x) = \frac{1}{x+3} &\Rightarrow f'(x) = \frac{-1}{(x+3)^2} \\ \therefore \int e^x [f(x) + f'(x)] \cdot dx &= e^x \cdot f(x) + c \end{aligned}$$

$$= e^x \cdot \left(\frac{1}{x+3} \right) + c$$

$$= \frac{e^x}{x+3} + c$$

$$\therefore \int e^x \left[\frac{x+2}{(x+3)^2} \right] \cdot dx = \frac{e^x}{x+3} + c$$

$$3. \int e^{\tan^{-1} x} \cdot \left(\frac{1+x+x^2}{1+x^2} \right) \cdot dx$$

Solution : put $\tan^{-1} x = t$

$$\therefore x = \tan t$$

differentiating w. r. t. x

$$\therefore \frac{1}{1+x^2} \cdot dx = 1 \cdot dt$$

$$I = \int e^t \cdot [1 + \tan t + \tan^2 t] \cdot dt$$

$$= \int e^t \cdot [\tan t + (1 + \tan^2 t)] \cdot dt$$

$$= \int e^t \cdot [\tan t + \sec^2 t] \cdot dt$$

Here $f(t) = \tan t$

$$\Rightarrow f'(t) = \sec^2 t$$

$$I = e^t \cdot f(t) + c$$

$$= e^t \cdot \tan t + c$$

$$= e^{\tan^{-1} x} \cdot x + c$$

$$\therefore \int e^{\tan^{-1} x} \cdot \left(\frac{1+x+x^2}{1+x^2} \right) \cdot dx = e^{\tan^{-1} x} \cdot x + c$$

$$4. \int \frac{(x^2+1) \cdot e^x}{(x+1)^2} \cdot dx$$

Solution :

$$I = \int e^x \left[\frac{x^2+1}{(x+1)^2} \right] \cdot dx$$

$$= \int e^x \left[\frac{x^2-1+2}{(x+1)^2} \right] \cdot dx$$

$$= \int e^x \left[\frac{x^2-1}{(x+1)^2} + \frac{2}{(x+1)^2} \right] \cdot dx$$

$$= \int e^x \left[\frac{x-1}{x+1} + \frac{2}{(x+1)^2} \right] \cdot dx$$

Here $f(x) = \frac{x-1}{x+1}$

$$\Rightarrow f'(x) = \frac{(x+1)(1) - (x-1)(1)}{(x+1)^2} = \frac{2}{(x+1)^2}$$

$$\therefore \int [f(x) + f'(x)] \cdot dx = e^x \cdot f(x) + c$$

$$I = e^x \cdot \left(\frac{x-1}{x+1} \right) + c$$

$$\therefore \int \frac{(x^2+1) \cdot e^x}{(x+1)^2} \cdot dx = e^x \cdot \left(\frac{x-1}{x+1} \right) + c$$

EXERCISE 3.3

I. Evaluate the following :

1. $\int x^2 \cdot \log x \cdot dx$

2. $\int x^2 \cdot \sin 3x \cdot dx$

3. $\int x \cdot \tan^{-1} x \cdot dx$

4. $\int x^2 \cdot \tan^{-1} x \cdot dx$

5. $\int x^3 \cdot \tan^{-1} x \cdot dx$

6. $\int (\log x)^2 \cdot dx$

7. $\int \sec^3 x \cdot dx$

8. $\int x \cdot \sin^2 x \cdot dx$

9. $\int x^3 \cdot \log x \cdot dx$

10. $\int e^{2x} \cdot \cos 3x \cdot dx$

11. $\int x \cdot \sin^{-1} x \cdot dx$

12. $\int x^2 \cdot \cos^{-1} x \cdot dx$

13. $\int \frac{\log(\log x)}{x} \cdot dx$

14. $\int \frac{t \cdot \sin^{-1} t}{\sqrt{1-t^2}} \cdot dt$

15. $\int \cos \sqrt{x} \cdot dx$

16. $\int \sin \theta \cdot \log(\cos \theta) \cdot d\theta$

17. $\int x \cdot \cos^3 x \cdot dx$

18. $\int \frac{\sin(\log x)^2}{x} \cdot \log x \cdot dx$

19. $\int \frac{\log x}{x} \cdot dx$

20. $\int x \cdot \sin 2x \cdot \cos 5x \cdot dx$

21. $\int \cos(\sqrt[3]{x}) \cdot dx$

II. Integrate the following functions w. r. t. x :

- | | | |
|---|---------------------------------|----------------------------|
| 1. $e^{2x} \cdot \sin 3x$ | 2. $e^{-x} \cdot \cos 2x$ | 3. $\sin (\log x)$ |
| 4. $\sqrt{5x^2 + 3}$ | 5. $x^2 \cdot \sqrt{a^2 - x^6}$ | 6. $\sqrt{(x-3)(7-x)}$ |
| 7. $\sqrt{4^x(4^x + 4)}$ | 8. $(x+1)\sqrt{2x^2 + 3}$ | 9. $x \sqrt{5 - 4x - x^2}$ |
| 10. $\sec^2 x \cdot \sqrt{\tan^2 x + \tan x - 7}$ | 11. $\sqrt{x^2 + 2x + 5}$ | 12. $\sqrt{2x^2 + 3x + 4}$ |

III. Integrate the following functions w. r. t. x :

- | | | |
|---|---|--|
| 1. $(2 + \cot x - \operatorname{cosec}^2 x) \cdot e^x$ | 2. $\left(\frac{1 + \sin x}{1 + \cos x}\right) \cdot e^x$ | 3. $e^x \cdot \left(\frac{1}{x} - \frac{1}{x^2}\right)$ |
| 4. $\left(\frac{x}{(x+1)^2}\right) \cdot e^x$ | 5. $\frac{e^x}{x} [x (\log x)^2 + 2 (\log x)]$ | 6. $e^{5x} \cdot \left(\frac{5x \cdot \log x + 1}{x}\right)$ |
| 7. $e^{\sin^{-1} x} \cdot \left(\frac{x + \sqrt{1-x^2}}{\sqrt{1-x^2}}\right)$ | 8. $\log (1+x)^{(1+x)}$ | |
| 9. $\operatorname{cosec} (\log x) [1 - \cot (\log x)]$ | | |

3.4 Integration by partial fraction :

If $f(x)$ and $g(x)$ are two polynomials then $\frac{f(x)}{g(x)}$, $g(x) \neq 0$ is called a rational algebraic function.

$\frac{f(x)}{g(x)}$ is called a proper rational function provided degree of $f(x) <$ degree of $g(x)$; otherwise it is called **improper rational function**.

If degree of $f(x) \geq$ degree of $g(x)$ i.e. $\frac{f(x)}{g(x)}$ is an improper rational function then express it as in the form $\text{Quotient} + \frac{\text{Remainder}}{g(x)}$, $g(x) \neq 0$ where $\frac{\text{Remainder}}{g(x)}$ is proper rational function.

Lets see the three different types of the proper rational function $\frac{f(x)}{g(x)}$, $g(x) \neq 0$ where the denominator $g(x)$ is expressed as

- a non-repeated linear factors
- repeated Linear factors and
- product of Linear factor and non-repeated quadratic factor.

No.	Rational form	Partial form
(i)	$\frac{px^2 + qx + r}{(x - a)(x - b)(x - c)}$	$\frac{A}{(x - a)} + \frac{B}{(x - b)} + \frac{C}{(x - c)}$
(ii)	$\frac{px^2 + qx + r}{(x - a)^2(x - b)}$	$\frac{A}{(x - a)} + \frac{B}{(x - a)^2} + \frac{C}{(x - b)}$
(iii)	$\frac{px^2 + qx + r}{(x - a)(x^2 + bx + c)}$	$\frac{A}{(x - a)} + \frac{Bx + C}{x^2 + bx + c}$

Type (i) : $\int \frac{px^2 + qx + r}{(x - a)(x - b)(x - c)} \cdot dx$ i.e. denominator is expressed as non-repeated Linear factors.



SOLVED EXAMPLES

1. $\int \frac{3x^2 + 4x - 5}{(x^2 - 1)(x + 2)} \cdot dx$

Solution : $I = \int \frac{3x^2 + 4x - 5}{(x - 1)(x + 1)(x + 2)} \cdot dx$

Consider, $\frac{3x^2 + 4x - 5}{(x - 1)(x + 1)(x + 2)} = \frac{A}{(x - 1)} + \frac{B}{(x + 1)} + \frac{C}{(x + 2)}$

$$= \frac{A(x + 1)(x + 2) + B(x - 1)(x + 2) + C(x - 1)(x + 1)}{(x - 1)(x + 1)(x + 2)}$$

$\therefore 3x^2 + 4x - 5 = A(x + 1)(x + 2) + B(x - 1)(x + 2) + C(x - 1)(x + 1)$

at $x = 1$, $3(1)^2 + 4(1) - 5 = A(2)(3) + B(0) + C(0)$

$2 = 6A \Rightarrow A = \frac{1}{3}$

at $x = -1$, $3(-1)^2 + 4(-1) - 5 = A(0) + B(-2)(1) + C(0)$

$-6 = -2B \Rightarrow B = 3$

at $x = -2$, $3(-2)^2 + 4(-2) - 5 = A(0) + B(0) + C(-3)(-1)$

$-1 = 3C \Rightarrow C = -\frac{1}{3}$

Thus, $\frac{3x^2 + 4x - 5}{(x - 1)(x + 1)(x + 2)} = \frac{\left(\frac{1}{3}\right)}{(x - 1)} + \frac{3}{(x + 1)} + \frac{\left(-\frac{1}{3}\right)}{(x + 2)}$

$\therefore I = \int \left[\frac{\left(\frac{1}{3}\right)}{(x - 1)} + \frac{3}{(x + 1)} + \frac{\left(-\frac{1}{3}\right)}{(x + 2)} \right] \cdot dx = \frac{1}{3} \log(x - 1) + 3 \log(x + 1) - \frac{1}{3} \log(x + 2) + c$

$= \frac{1}{3} \log \left[\frac{(x - 1)(x + 1)^9}{(x + 2)} \right] + c \quad \therefore \int \frac{3x^2 + 4x - 5}{(x^2 - 1)(x + 2)} \cdot dx = \frac{1}{3} \log \left[\frac{(x - 1)(x + 1)^9}{(x + 2)} \right] + c$



$$2. \int \frac{2x^2 - 3}{(x^2 - 5)(x^2 + 4)} \cdot dx$$

Solution : Consider, $\frac{2x^2 - 3}{(x^2 - 5)(x^2 + 4)}$

Let $x^2 = m$

$$\therefore = \frac{2m - 3}{(m - 5)(m + 4)} \dots \text{proper rational function.}$$

$$\text{Now, } \frac{2m - 3}{(m - 5)(m + 4)} = \frac{A}{(m - 5)} + \frac{B}{(m + 4)} = \frac{A(m + 4) + B(m - 5)}{(m - 5)(m + 4)}$$

$$\therefore 2m - 3 = A(m + 4) + B(m - 5)$$

at $m = 5$, $2(5) - 3 = A(9) + B(0)$

$$7 = 9A \Rightarrow A = \frac{7}{9}$$

at $m = -4$, $2(-4) - 3 = A(0) + B(-9)$

$$-11 = -9B \Rightarrow B = \frac{11}{9}$$

$$\text{Thus, } \frac{2m - 3}{(m - 5)(m + 4)} = \frac{\left(\frac{7}{9}\right)}{(m - 5)} + \frac{\left(\frac{11}{9}\right)}{(m + 4)} \quad \text{i.e. } \frac{2x^2 - 3}{(x^2 - 5)(x^2 + 4)} = \frac{\left(\frac{7}{9}\right)}{x^2 - 5} + \frac{\left(\frac{11}{9}\right)}{x^2 + 4}$$

$$\therefore I = \int \left[\frac{\left(\frac{7}{9}\right)}{x^2 - 5} + \frac{\left(\frac{11}{9}\right)}{x^2 + 4} \right] \cdot dx$$

$$= \frac{7}{9} \cdot \int \frac{1}{x^2 - (\sqrt{5})^2} \cdot dx + \frac{11}{9} \cdot \int \frac{1}{x^2 + (2)^2} \cdot dx$$

$$= \frac{7}{9} \cdot \frac{1}{2(\sqrt{5})} \cdot \log \left[\frac{x - \sqrt{5}}{x + \sqrt{5}} \right] + \frac{11}{9} \cdot \frac{1}{2} \cdot \tan^{-1} \left(\frac{x}{2} \right) + c$$

$$\therefore I = \frac{7}{18(\sqrt{5})} \cdot \log \left[\frac{x - \sqrt{5}}{x + \sqrt{5}} \right] + \frac{11}{18} \cdot \tan^{-1} \left(\frac{x}{2} \right) + c$$

$$\therefore \int \frac{2x^2 - 3}{(x^2 - 5)(x^2 + 4)} \cdot dx = \frac{7}{18(\sqrt{5})} \cdot \log \left[\frac{x - \sqrt{5}}{x + \sqrt{5}} \right] + \frac{11}{18} \cdot \tan^{-1} \left(\frac{x}{2} \right) + c$$



$$4. \int \frac{1}{2 \cos x + \sin 2x} \cdot dx$$

$$\text{Solution : } I = \int \frac{1}{2 \cos x + \sin 2x} \cdot dx = \int \frac{1}{2 \cos x + 2 \sin x \cdot \cos x} \cdot dx = \int \frac{1}{2 (\cos x) (1 + \sin x)} \cdot dx$$

$$= \frac{1}{2} \cdot \int \frac{\cos x}{\cos^2 x (1 + \sin x)} \cdot dx = \frac{1}{2} \cdot \int \frac{\cos x}{(1 - \sin^2 x) (1 + \sin x)} \cdot dx$$

$$\text{put } \sin x = t \quad \therefore \cos x \cdot dx = 1 \cdot dt$$

$$= \frac{1}{2} \cdot \int \frac{1}{(1 - t^2) (1 + t)} \cdot dt = \frac{1}{2} \cdot \int \frac{1}{(1 - t) (1 + t) (1 + t)} \cdot dt = \frac{1}{2} \cdot \int \frac{1}{(1 - t) (1 + t)^2} \cdot dt$$

$$\text{Consider, } \frac{1}{(1 - t) (1 + t)^2} = \frac{A}{(1 - t)} + \frac{B}{(1 + t)} + \frac{C}{(1 + t)^2} = \frac{A(1 + t)^2 + B(1 - t)(1 + t) + C(1 - t)}{(1 - t) (1 + t)^2}$$

$$\therefore 1 = A(1 + t)^2 + B(1 - t)(1 + t) + C(1 - t)$$

$$\text{at } t = 1, \quad 1 = A(2)^2 + B(0) + C(0)$$

$$1 = 4A \quad \Rightarrow \quad A = \frac{1}{4}$$

$$\text{at } t = -1, \quad 1 = A(0) + B(0) + C(2)$$

$$1 = 2C \quad \Rightarrow \quad C = \frac{1}{2}$$

$$\text{at } t = 0, \quad 1 = A(1)^2 + B(1)(1) + C(1)$$

$$1 = A + B + C$$

$$1 = \frac{1}{4} + B + \frac{1}{2} \quad \Rightarrow \quad B = \frac{1}{4}$$

$$\text{Thus, } \frac{1}{(1 - t) (1 + t)^2} = \frac{\left(\frac{1}{4}\right)}{(1 - t)} + \frac{\left(\frac{1}{4}\right)}{(1 + t)} + \frac{\left(\frac{1}{2}\right)}{(1 + t)^2}$$

$$\therefore I = \int \left[\frac{\left(\frac{1}{4}\right)}{(1 - t)} + \frac{\left(\frac{1}{4}\right)}{(1 + t)} + \frac{\left(\frac{1}{2}\right)}{(1 + t)^2} \right] \cdot dt = \frac{1}{2} \left[\frac{1}{4} \log(1 - t) \cdot \frac{1}{(-1)} + \frac{1}{4} \log(1 + t) + \frac{1}{2} \cdot \frac{(-1)}{(1 + t)} \right] + c$$

$$= \frac{1}{2} \left[\frac{1}{4} \log(1 - t) \cdot \frac{1}{(-1)} + \frac{1}{4} \log(1 + t) + \frac{1}{2} \cdot \frac{-1}{1 + t} \right] + c$$

$$= \frac{1}{8} \left[-\log(1 - \sin x) + \log(1 + \sin x) - \frac{2}{1 + \sin x} \right] + c = \frac{1}{8} \left[\log \left(\frac{1 + \sin x}{1 - \sin x} \right) - \frac{2}{1 + \sin x} \right] + c$$

$$\therefore \int \frac{1}{2 \cos x + \sin 2x} \cdot dx = \frac{1}{8} \left[\log \left(\frac{1 + \sin x}{1 - \sin x} \right) - \frac{2}{1 + \sin x} \right] + c$$

5. $\int \frac{\tan \theta + \tan^3 \theta}{1 + \tan^3 \theta} \cdot d\theta$

Solution : $I = \int \frac{(\tan \theta) (1 + \tan^2 \theta)}{1 + \tan^3 \theta} \cdot d\theta = \int \frac{(\tan \theta) \cdot (1 + \tan^2 \theta)}{1 + \tan^3 \theta} \cdot d\theta = \int \frac{\tan \theta \cdot \sec^2 \theta}{1 + \tan^3 \theta} \cdot d\theta$

put $\tan \theta = x \quad \therefore \sec^2 \theta \cdot d\theta = 1 \cdot dx$

$$= \int \frac{x}{1+x^3} \cdot dx = \int \frac{x}{(1+x)(1-x+x^2)} \cdot dx$$

Consider, $\frac{x}{(1+x)(1-x+x^2)} = \frac{A}{1+x} + \frac{Bx+C}{1-x+x^2}$

$$= \frac{A(1-x+x^2) + Bx + C(1+x)}{(1+x)(1-x+x^2)}$$

$$\therefore x = A(1-x+x^2) + (Bx+C)(1+x) = A - Ax + Ax^2 + Bx + Bx^2 + C + Cx$$

$$0x^2 + 1x + 0 = (A+B)x^2 + (-A+B+C)x + (A+C)$$

comparing the co-efficients of like powers of variables.

$$0 = A + B \quad \dots \text{(I)}$$

$$1 = -A + B + C \quad \dots \text{(II)} \quad \text{and}$$

$$0 = A + C \quad \dots \text{(III)}$$

Solving these equations, we get $A = -\frac{1}{3}$; $B = \frac{1}{3}$ and $C = \frac{1}{3}$

Thus, $\frac{x}{(1+x)(1-x+x^2)} = \frac{\left(-\frac{1}{3}\right)}{1+x} + \frac{\left(\frac{1}{3}x + \frac{1}{3}\right)}{(1-x+x^2)}$

$$\therefore I = \int \left[\frac{\left(-\frac{1}{3}\right)}{1+x} + \frac{\left(\frac{1}{3}x + \frac{1}{3}\right)}{(1-x+x^2)} \right] \cdot dx = -\frac{1}{3} \int \frac{1}{1+x} \cdot dx + \frac{1}{3} \int \frac{x+1}{1-x+x^2} \cdot dx$$

$$= -\frac{1}{3} \int \frac{1}{1+x} \cdot dx + \frac{1}{3} \cdot \frac{1}{(2)} \int \frac{2x-1+3}{x^2-x+1} \cdot dx \quad \because \frac{d}{dx} x^2 - x + 1 = 2x - 1$$

$$= -\frac{1}{3} \int \frac{1}{1+x} \cdot dx + \frac{1}{3} \cdot \frac{1}{2} \int \frac{2x-1+3}{x^2-x+1} \cdot dx$$

$$= -\frac{1}{3} \int \frac{1}{1+x} \cdot dx + \frac{1}{6} \int \frac{2x-1}{x^2-x+1} \cdot dx + \frac{1}{6} \int \frac{3}{x^2-x+1} \cdot dx$$

$$= I_1 + I_2 + I_3 \quad \dots \text{(IV)}$$



$$\begin{aligned} \therefore I_1 &= -\frac{1}{3} \int \frac{1}{1+x} \cdot dx = -\frac{1}{3} [\log(1+x)] \\ &= -\frac{1}{3} \log(1 + \tan \theta) \quad \dots \text{(V)} \end{aligned}$$

$$\begin{aligned} \therefore I_2 &= \frac{1}{6} \int \frac{2x-1}{x^2-x+1} \cdot dx = \frac{1}{6} [\log(x^2-x+1)] \\ &= \frac{1}{6} \log(\tan^2 \theta - \tan \theta + 1) \quad \dots \text{(VI)} \end{aligned}$$

$$\begin{aligned} \therefore I_3 &= \frac{1}{6} \int \frac{3}{x^2-x+1} \cdot dx \\ &= \frac{1}{2} \int \frac{1}{x^2-x+\frac{1}{4}-\frac{1}{4}+1} \cdot dx \quad \because \left\{ \left(\frac{1}{2} \text{ coefficient of } x \right)^2 = \left(\frac{1}{2} (-1) \right)^2 = \left(-\frac{1}{2} \right)^2 = \frac{1}{4} \right\} \\ &= \frac{1}{2} \int \frac{1}{\left(x - \frac{1}{2} \right)^2 + \left(\frac{\sqrt{3}}{2} \right)^2} \cdot dx \\ &= \frac{1}{2} \cdot \left[\frac{1}{\left(\frac{\sqrt{3}}{2} \right)} \right] \cdot \tan^{-1} \left[\frac{x - \frac{1}{2}}{\left(\frac{\sqrt{3}}{2} \right)} \right] + c \\ &= \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{2x-1}{\sqrt{3}} \right) + c \\ \therefore I_3 &= \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{2 \tan \theta - 1}{\sqrt{3}} \right) + c \quad \dots \text{(VII)} \end{aligned}$$

$$\therefore \int \frac{\tan \theta + \tan^3 \theta}{1 + \tan^3 \theta} \cdot d\theta = -\frac{1}{3} \log(1 + \tan \theta) + \frac{1}{6} \log(\tan^2 \theta - \tan \theta + 1) + \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{2 \tan \theta - 1}{\sqrt{3}} \right) + c$$

EXERCISE 3.4

I. Integrate the following w. r. t. x :

1. $\frac{x^2+2}{(x-1)(x+2)(x+3)}$

2. $\frac{x^2}{(x^2+1)(x^2-2)(x^2+3)}$

3. $\frac{12x+3}{6x^2+13x-63}$

- | | | |
|---|---|---|
| 4. $\frac{2x}{4 - 3x - x^2}$ | 5. $\frac{x^2 + x - 1}{x^2 + x - 6}$ | 6. $\frac{6x^3 + 5x^2 - 7}{3x^2 - 2x - 1}$ |
| 7. $\frac{12x^2 - 2x - 9}{(4x^2 - 1)(x + 3)}$ | 8. $\frac{1}{x(x^5 + 1)}$ | 9. $\frac{2x^2 - 1}{x^4 + 9x^2 + 20}$ |
| 10. $\frac{x^2 + 3}{(x^2 - 1)(x^2 - 2)}$ | 11. $\frac{2x}{(2 + x^2)(3 + x^2)}$ | 12. $\frac{2^x}{4^x - 3 \cdot 2^x - 4}$ |
| 13. $\frac{3x - 2}{(x + 1)^2(x + 3)}$ | 14. $\frac{5x^2 + 20x + 6}{x^3 + 2x^2 + x}$ | 15. $\frac{1}{x(1 + 4x^3 + 3x^6)}$ |
| 16. $\frac{1}{x^3 - 1}$ | 17. $\frac{(3 \sin x - 2) \cdot \cos x}{5 - 4 \sin x - \cos^2 x}$ | 18. $\frac{1}{\sin x + \sin 2x}$ |
| 19. $\frac{1}{2 \sin x + \sin 2x}$ | 20. $\frac{1}{\sin 2x + \cos x}$ | 21. $\frac{1}{\sin x \cdot (3 + 2 \cos x)}$ |
| 22. $\frac{5 \cdot e^x}{(e^x + 1)(e^{2x} + 9)}$ | 23. $\frac{2 \log x + 3}{x(3 \log x + 2)[(\log x)^2 + 1]}$ | |

3.5 Something Interesting :

Students/ now familiar with the integration by parts.

The result is $\int u \cdot v \cdot dx = u \cdot \int v \cdot dx - \int \left(\frac{d}{dx} \cdot u \right) (\int v \cdot dx) \cdot dx$,

u and v are differentiable functions of x and $u \cdot v$ follows L I A T E order.

This result can be extended to the generalisation as -

$$\int u \cdot v \cdot dx = u \cdot v_1 - u' \cdot v_2 + u'' \cdot v_3 - u''' \cdot v_4 + \dots$$

(') dash indicates the derivative.

(₁) subscript indicates the integration.

This result is more useful where the first function (u) is a polynomial, because $\frac{d^n u}{dx^n} = 0$ for some n .

For example : $\int x^2 \cdot \cos 3x \cdot dx$

$$\begin{aligned} &= x^2 \cdot \left(\sin 3x \cdot \frac{1}{3} \right) - (2x) \left(-\cos 3x \cdot \frac{1}{3} \cdot \frac{1}{3} \right) + (2) \left(-\sin 3x \cdot \frac{1}{3} \cdot \frac{1}{9} \right) - (0) \\ &= \frac{1}{3} x^2 \cdot \sin 3x + \frac{2}{9} x \cdot \cos 3x - \frac{2}{27} \sin 3x + c \end{aligned}$$

verify this example with usual rule of integration by parts.





Let us Remember

✿ We can always add arbitrary constant c to the integration obtained :

(I) i.e. $\frac{d}{dx} \cdot g(x) = f(x) \Rightarrow \int f(x) \cdot dx = g(x) + c$

$f(x)$ is integrand, $g(x)$ is integral of $f(x)$ with respect to x , c is arbitrary constant.

(II) $\int f(ax + b) \cdot dx = g(ax + b) \cdot \frac{1}{a} + c$

(III) (1) $\int [f(x)]^n \cdot f'(x) \cdot dx = \frac{[f(x)]^{n+1}}{n+1} + c$ (2) $\int \frac{f'(x)}{f(x)} \cdot dx = \log(f(x)) + c$

(3) $\int \frac{f'(x)}{\sqrt{f(x)}} \cdot dx = 2\sqrt{f(x)} + c$

(IV) (1) $\int x^n \cdot dx = \frac{x^{n+1}}{n+1} + c$

(2) $\int \frac{1}{\sqrt{x}} \cdot dx = 2\sqrt{x} + c$

(3) $\int \text{constant } (k) \cdot dx = kx + c$

(4) $\int a^x \cdot dx = \frac{a^x}{\log a} + c$

(5) $\int e^x \cdot dx = e^x + c$

(6) $\int \frac{1}{x} \cdot dx = \log(x) + c$

(7) $\int \sin x \cdot dx = -\cos x + c$

(8) $\int \cos x \cdot dx = \sin x + c$

(9) $\int \tan x \cdot dx = \log(\sec x) + c$

(10) $\int \cot x \cdot dx = \log(\sin x) + c$

(11) $\int \sec x \cdot dx = \log(\sec x + \tan x) + c$ (12) $\int \text{cosec } x \cdot dx = \log(\text{cosec } x - \cot x) + c$

$= \log \left[\tan \left(\frac{x}{2} + \frac{\pi}{4} \right) \right] + c$

$= \log \left[\tan \left(\frac{x}{2} \right) \right] + c$

(13) $\int \sec^2 x \cdot dx = \tan x + c$

(14) $\int \text{cosec}^2 x \cdot dx = -\cot x + c$

(15) $\int \sec x \cdot \tan x \cdot dx = \sec x + c$

(16) $\int \text{cosec } x \cdot \cot x \cdot dx = -\text{cosec } x + c$

(17) $\int \frac{1}{\sqrt{1-x^2}} \cdot dx = \sin^{-1} x + c$

(18) $\int \frac{-1}{\sqrt{1-x^2}} \cdot dx = \cos^{-1} x + c$

(19) $\int \frac{1}{1+x^2} \cdot dx = \tan^{-1} x + c$

(20) $\int \frac{-1}{1+x^2} \cdot dx = \cot^{-1} x + c$

(21) $\int \frac{1}{x \cdot \sqrt{x^2-1}} \cdot dx = \sec^{-1} x + c$

(22) $\int \frac{-1}{x \cdot \sqrt{x^2-1}} \cdot dx = \text{cosec}^{-1} x + c$

(23) $\int \frac{1}{x^2 + a^2} \cdot dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + c$ (24) $\int \frac{1}{x^2 - a^2} \cdot dx = \frac{1}{2a} \log \left(\frac{x-a}{x+a} \right) + c$

(25) $\int \frac{1}{a^2 - x^2} \cdot dx = \frac{1}{2a} \log \left(\frac{a+x}{a-x} \right) + c$ (26) $\int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx = \sin^{-1} \left(\frac{x}{a} \right) + c$

(27) $\int \frac{1}{\sqrt{x^2 - a^2}} \cdot dx = \log |x + \sqrt{x^2 - a^2}| + c$ (28) $\int \frac{1}{\sqrt{x^2 + a^2}} \cdot dx = \log |x + \sqrt{x^2 + a^2}| + c$

(29) $\int \frac{1}{x\sqrt{x^2 - a^2}} \cdot dx = \frac{1}{a} \sec^{-1} \left(\frac{x}{a} \right) + c$

(30) $\int \sqrt{a^2 - x^2} \cdot dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) + c$

(31) $\int \sqrt{a^2 + x^2} \cdot dx = \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \log (x + \sqrt{x^2 + a^2}) + c$

(32) $\int \sqrt{x^2 - a^2} \cdot dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log (x + \sqrt{x^2 - a^2}) + c$

(V) If u and v are differentiable functions of x then $\int u \cdot v \cdot dx = u \cdot \int v \cdot dx - \int \left(\frac{d}{dx} \cdot u \right) (\int v \cdot dx) \cdot dx$ where $u \cdot v$ follows the L I A T E order.

(VI) $\int e^x [f(x) + f'(x)] \cdot dx = e^x \cdot f(x) + c$

(VII) For the integration of type $\int \frac{f(x)}{g(x)} \cdot dx$, $g(x) \neq 0$ where $\frac{f(x)}{g(x)}$ proper rational function.

- (i) non-repeated linear factors
- (ii) repeated Linear factors and
- (iii) product of Linear factor and non-repeated quadratic factor.

(VIII) $\int \frac{1}{x^2 + a^2} \cdot dx$ $\int \frac{1}{ax^2 + bx + c} \cdot dx$

$\int \frac{1}{x^2 - a^2} \cdot dx$ Method of completing square

$\int \frac{1}{a^2 - x^2} \cdot dx$ Divide Nr and Dr by $\cos^2 x$

$\int \frac{1}{a \sin^2 x + b \cos^2 x + c} \cdot dx$

$\int \frac{1}{a \sin x + b \cos x + c} \cdot dx$

put $\tan \left(\frac{x}{2} \right) = t$

$\int \frac{px + q}{ax^2 + bx + c} \cdot dx$

$px + q = A \frac{d}{dx} (ax^2 + bx + c) + B$

(VIII) $\int \frac{1}{\sqrt{x^2 + a^2}} \cdot dx$ $\int \frac{1}{\sqrt{ax^2 + bx + c}} \cdot dx$

$\int \frac{1}{\sqrt{x^2 - a^2}} \cdot dx$ $\int \frac{px + q}{\sqrt{ax^2 + bx + c}} \cdot dx$

$\int \frac{1}{\sqrt{a^2 - x^2}} \cdot dx$

MISCELLANEOUS EXERCISE 3

(I) Choose the correct option from the given alternatives :

(1) $\int \frac{1+x+\sqrt{x+x^2}}{\sqrt{x}+\sqrt{1+x}} \cdot dx =$

- (A) $\frac{1}{2} \sqrt{x+1} + c$ (B) $\frac{2}{3} (x+1)^{\frac{3}{2}} + c$ (C) $\sqrt{x+1} + c$ (D) $2(x+1)^{\frac{3}{2}} + c$

(2) $\int \frac{1}{x+x^5} \cdot dx = f(x) + c$, then $\int \frac{x^4}{x+x^5} \cdot dx =$

- (A) $\log x - f(x) + c$ (B) $f(x) + \log x + c$ (C) $f(x) - \log x + c$ (D) $\frac{1}{5} x^5 f(x) + c$

(3) $\int \frac{\log(3x)}{x \log(9x)} \cdot dx =$

- (A) $\log(3x) - \log(9x) + c$ (B) $\log(x) - (\log 3) \cdot \log(\log 9x) + c$
 (C) $\log 9 - (\log x) \cdot \log(\log 3x) + c$ (D) $\log(x) + (\log 3) \cdot \log(\log 9x) + c$

(4) $\int \frac{\sin^m x}{\cos^{m+2} x} \cdot dx =$

- (A) $\frac{\tan^{m+1} x}{m+1} + c$ (B) $(m+2) \tan^{m+1} x + c$ (C) $\frac{\tan^m x}{m} + c$ (D) $(m+1) \tan^{m+1} x + c$

(5) $\int \tan(\sin^{-1} x) \cdot dx =$

- (A) $(1-x^2)^{-\frac{1}{2}} + c$ (B) $(1-x^2)^{\frac{1}{2}} + c$ (C) $\frac{\tan^m x}{\sqrt{1-x^2}} + c$ (D) $-\sqrt{1-x^2} + c$

(6) $\int \frac{x - \sin x}{1 - \cos x} \cdot dx =$

- (A) $x \cot\left(\frac{x}{2}\right) + c$ (B) $-x \cot\left(\frac{x}{2}\right) + c$ (C) $\cot\left(\frac{x}{2}\right) + c$ (D) $x \tan\left(\frac{x}{2}\right) + c$

(7) If $f(x) = \frac{\sin^{-1} x}{\sqrt{1-x^2}}$, $g(x) = e^{\sin^{-1} x}$, then $\int f(x) \cdot g(x) \cdot dx =$

- (A) $e^{\sin^{-1} x} \cdot (\sin^{-1} x - 1) + c$ (B) $e^{\sin^{-1} x} \cdot (1 - \sin^{-1} x) + c$
 (C) $e^{\sin^{-1} x} \cdot (\sin^{-1} x + 1) + c$ (D) $e^{\sin^{-1} x} \cdot (\sin^{-1} x - 1) + c$

(8) If $\int \tan^3 x \cdot \sec^3 x \cdot dx = \left(\frac{1}{m}\right) \sec^m x - \left(\frac{1}{n}\right) \sec^n x + c$, then $(m, n) =$

- (A) (5, 3) (B) (3, 5) (C) $\left(\frac{1}{5}, \frac{1}{3}\right)$ (D) (4, 4)

- (9) $\int \frac{1}{\cos x - \cos^2 x} \cdot dx =$
 (A) $\log (\operatorname{cosec} x - \cot x) + \tan \left(\frac{x}{2} \right) + c$ (B) $\sin 2x - \cos x + c$
 (C) $\log (\sec x + \tan x) - \cot \left(\frac{x}{2} \right) + c$ (D) $\cos 2x - \sin x + c$
- (10) $\int \frac{\sqrt{\cot x}}{\sin x \cdot \cos x} \cdot dx =$
 (A) $2 \sqrt{\cot x} + c$ (B) $-2 \sqrt{\cot x} + c$ (C) $\frac{1}{2} \sqrt{\cot x} + c$ (D) $\sqrt{\cot x} + c$
- (11) $\int \frac{e^x (x-1)}{x^2} \cdot dx =$
 (A) $\frac{e^x}{x} + c$ (B) $\frac{e^x}{x^2} + c$ (C) $\left(x - \frac{1}{x} \right) e^x + c$ (D) $x e^{-x} + c$
- (12) $\int \sin (\log x) \cdot dx =$
 (A) $\frac{x}{2} [\sin (\log x) - \cos (\log x)] + c$ (B) $\frac{x}{2} [\sin (\log x) + \cos (\log x)] + c$
 (C) $\frac{x}{2} [\cos (\log x) - \sin (\log x)] + c$ (D) $\frac{x}{4} [\cos (\log x) - \sin (\log x)] + c$
- (13) $\int x^x (1 + \log x) \cdot dx =$
 (A) $\frac{1}{2} (1 + \log x)^2 + c$ (B) $x^{2x} + c$ (C) $x^x \log x + c$ (D) $x^x + c$
- (14) $\int \cos^{-\frac{3}{7}} x \cdot \sin^{-\frac{11}{7}} x \cdot dx =$
 (A) $\log \left(\sin^{-\frac{4}{7}} x \right) + c$ (B) $\frac{4}{7} \tan^{\frac{4}{7}} x + c$
 (C) $-\frac{7}{4} \tan^{-\frac{4}{7}} x + c$ (D) $\log \left(\cos^{\frac{3}{7}} x \right) + c$
- (15) $2 \int \frac{\cos^2 x - \sin^2 x}{\cos^2 x + \sin^2 x} \cdot dx =$
 (A) $\sin 2x + c$ (B) $\cos 2x + c$ (C) $\tan 2x + c$ (D) $2 \sin 2x + c$
- (16) $\int \frac{dx}{\cos x \sqrt{\sin^2 x - \cos^2 x}} \cdot dx =$
 (A) $\log \left(\tan x - \sqrt{\tan^2 x - 1} \right) + c$ (B) $\sin^{-1} (\tan x) + c$
 (C) $1 + \sin^{-1} (\cot x) + c$ (D) $\log \left(\tan x + \sqrt{\tan^2 x - 1} \right) + c$

- (17) $\int \frac{\log x}{(\log ex)^2} \cdot dx =$
 (A) $\frac{x}{1 + \log x} + c$ (B) $x(1 + \log x) + c$ (C) $\frac{1}{1 + \log x} + c$ (D) $\frac{1}{1 - \log x} + c$
- (18) $\int [\sin(\log x) + \cos(\log x)] \cdot dx =$
 (A) $x \cos(\log x) + c$ (B) $\sin(\log x) + c$ (C) $\cos(\log x) + c$ (D) $x \sin(\log x) + c$
- (19) $\int \frac{\cos 2x - 1}{\cos 2x + 1} \cdot dx =$
 (A) $\tan x - x + c$ (B) $x + \tan x + c$ (C) $x - \tan x + c$ (D) $-x - \cot x + c$
- (20) $\int \frac{e^{2x} + e^{-2x}}{e^x} \cdot dx =$
 (A) $e^x - \frac{1}{3e^{3x}} + c$ (B) $e^x + \frac{1}{3e^{3x}} + c$ (C) $e^{-x} + \frac{1}{3e^{3x}} + c$ (D) $e^{-x} - \frac{1}{3e^{3x}} + c$

(II) Integrate the following with respect to the respective variable :

- | | | |
|--|---|---|
| (1) $(x - 2)^2 \sqrt{x}$ | (2) $\frac{x^7}{x + 1}$ | (3) $(6x + 5)^{\frac{3}{2}}$ |
| (4) $\frac{t^3}{(t + 1)^2}$ | (5) $\frac{3 - 2 \sin x}{\cos^2 x}$ | (6) $\frac{\sin^6 \theta + \cos^6 \theta}{\sin^2 \theta \cdot \cos^2 \theta}$ |
| (7) $\cos 3x \cdot \cos 2x \cdot \cos x$ | (8) $\frac{\cos 7x - \cos 8x}{1 + 2 \cos 5x}$ | (9) $\cot^{-1}\left(\frac{1 + \sin x}{\cos x}\right)$ |

(III) Integrate the following :

- | | | |
|--|---|--|
| (1) $\frac{(1 + \log x)^3}{x}$ | (2) $\cot^{-1}(1 - x + x^2)$ | (3) $\frac{1}{x \cdot \sin^2(\log x)}$ |
| (4) $\sqrt{x} \sec\left(x^{\frac{3}{2}}\right) \cdot \tan\left(x^{\frac{3}{2}}\right)$ | (5) $\log(1 + \cos x) - x \cdot \tan\left(\frac{x}{2}\right)$ | (6) $\frac{x^2}{\sqrt{1 - x^6}}$ |
| (7) $\frac{1}{(1 - \cos 4x)(3 - \cot 2x)}$ | (8) $\log(\log x) + (\log x)^{-2}$ | (9) $\frac{1}{2 \cos x + 3 \sin x}$ |
| (10) $\frac{1}{x^3 \cdot \sqrt{x^2 - 1}}$ | (11) $\frac{3x + 1}{\sqrt{-2x^2 + x + 3}}$ | (12) $\log(x^2 + 1)$ |
| (13) $e^{2x} \cdot \sin x \cdot \cos x$ | (14) $\frac{x^2}{(x - 1)(3x - 1)(3x - 2)}$ | (15) $\frac{1}{\sin x + \sin 2x}$ |
| (16) $\sec^2 x \cdot \sqrt{7 + 2 \tan x - \tan^2 x}$ | (17) $\frac{x + 5}{x^3 + 3x^2 - x - 3}$ | (18) $\frac{1}{x \cdot (x^5 + 1)}$ |
| (19) $\frac{\sqrt{\tan x}}{\sin x \cdot \cos x}$ | (20) $\sec^4 x \cdot \operatorname{cosec}^2 x$ | |



4. DEFINITE INTEGRATION



Let us Study

- Definite integral as limit of sum.
- Fundamental theorem of integral calculus.
- Methods of evaluation and properties of definite integral.

4.1 Definite integral as limit of sum :

In the last chapter, we studied various methods of finding the primitives or indefinite integrals of given function. We shall now interpret the definite integrals denoted by $\int_a^b f(x) dx$, read as the integral from a to b of the function $f(x)$ with respect to x . Here $a < b$, are real numbers and $f(x)$ is defined on $[a, b]$. At present, we assume that $f(x) \geq 0$ on $[a, b]$ and $f(x)$ is continuous.

$\int_a^b f(x) dx$ is defined as the area of the region bounded by $y = f(x)$, X-axis and the ordinates $x = a$ and $x = b$. If $g(x)$ is the primitive of $f(x)$ then the area is $g(b) - g(a)$.

The reason of the above definition will be clear from the figure 4.1. and the discussion that follows here. We are using the mean value theorem learnt earlier. Divide the interval $[a, b]$ into n equal parts by




Fig. 4.1

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$

Draw the curve $y = f(x)$ in $[a, b]$ and divide the interval $[a, b]$ into n equal parts by

$$a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b.$$

Divide the region whose area is measured into their strips as above.

Note that, the area of each strip can be approximated by the area of a rectangle $M_r M_{r+1} QP$ as shown in the figure 4.1, which is $(x_{r+1} - x_r) \times f(T)$ where T is a point on the curve $y = f(x)$ between P and Q .

The mean value theorem states that if $g(x)$ is the primitive of $f(x)$,

$$g(x_{r+1}) - g(x_r) = (x_{r+1} - x_r) \cdot f(t_r) \quad \text{where } x_r < t_r < x_{r+1}.$$

Now we can replace $f(T)$ by $f(t_r)$ given here and express the approximation of the area of the shaded region as $\sum_{r=0}^{n-1} (x_{r+1} - x_r) \cdot f(t_r)$ where $x_r < t_r < x_{r+1}$.


Now we can replace $f(T)$ by $f(t_r)$ given here and express the approximation of the area of the shaded region as

$$\sum_{r=0}^{n-1} (x_{r+1} - x_r) \cdot f(t_r) = \sum_{r=0}^{n-1} g(x_{r+1}) - g(x_r) = g(b) - g(a)$$

Thus taking limit as $n \rightarrow \infty$

$$\begin{aligned} g(b) - g(a) &= \lim_{n \rightarrow \infty} \sum (x_{r+1} - x_r) \cdot f(t_r) \\ &= \lim_{n \rightarrow \infty} S_n \\ &= \int_a^b f(x) dx \end{aligned}$$

The word 'to integrate' means 'to find the sum of'. The technique of integration is very useful in finding plane areas, length of arcs, volume of solid revolution etc...



SOLVED EXAMPLES

Ex. 1 : $\int_1^2 (2x + 5) dx$

Solution : Given, $\int_1^2 (2x + 5) dx = \int_a^b f(x) dx$

$$f(x) = 2x + 5 \quad a = 1 ; b = 2$$

$$\begin{aligned} \Rightarrow f(a + rh) &= f(1 + rh) & \text{and } h &= \frac{b - a}{n} \\ &= 2(1 + rh) + 5 \\ &= 2 + 2rh + 5 & h &= \frac{2 - 1}{n} \\ &= 7 + 2rh & \therefore nh &= 1 \end{aligned}$$

We know $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot f(a + rh)$

$$\begin{aligned}
\therefore \int_1^2 (2x + 5) dx &= \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot (7 + 2rh) \\
&= \lim_{n \rightarrow \infty} \sum_{r=1}^n (7h + 2rh^2) \\
&= \lim_{n \rightarrow \infty} \left(7h \sum_{r=1}^n 1 + 2h^2 \sum_{r=1}^n r \right) \\
&= \lim_{n \rightarrow \infty} \left[7h \cdot (n) + 2h^2 \left(\frac{n(n+1)}{2} \right) \right] \\
&= \lim_{n \rightarrow \infty} \left[7nh + h^2 n^2 \left(1 + \frac{1}{n} \right) \right] \\
&= \lim_{n \rightarrow \infty} \left[7(1) + (1)^2 \left(1 + \frac{1}{n} \right) \right] \\
&= 7 + 1(1 + 0) = 8
\end{aligned}$$

Ex. 2 : $\int_2^3 7^x \cdot dx$

Solution :

Given, $\int_2^3 7^x \cdot dx = \int_a^b f(x) dx$

$f(x) = 7^x \quad a = 2 ; b = 3$

$\Rightarrow f(a + rh) = f(1 + rh)$
 $= 7^{2+rh}$
 $= 7^2 \cdot 7^{rh}$

and $h = \frac{b-a}{n}$
 $h = \frac{3-2}{n} \quad \therefore nh = 1$

We know $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot f(a + rh)$

$$\begin{aligned}
\therefore \int_1^3 7^x \cdot dx &= \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot (7^2 \cdot 7^{r \cdot h}) \\
&= \lim_{n \rightarrow \infty} 7^2 \cdot \sum_{r=1}^n h \cdot 7^{r \cdot h} \\
&= \lim_{n \rightarrow \infty} 7^2 \cdot h \cdot [7^h + 7^{2h} + 7^{3h} + 7^{4h} + \dots + 7^{nh}] \\
&= \lim_{n \rightarrow \infty} 7^2 \cdot h \cdot \left(\frac{7^h [(7^h)^n - 1]}{7^h - 1} \right) = \lim_{n \rightarrow \infty} 7^2 \cdot \left(\frac{7^h (7^{nh} - 1)}{7^h - 1} \right) \\
&= \lim_{n \rightarrow \infty} 7^2 \cdot \left(\frac{7^h (7^{(1)} - 1)}{7^h - 1} \right) \\
&= \frac{7^2 \cdot 7^0 \cdot (7 - 1)}{\log 7} = \frac{(49)(1)(6)}{\log 7} = \frac{294}{\log 7}
\end{aligned}$$

Ex. 3 : $\int_0^4 (x - x^2) \cdot dx$

Solution : $\int_0^4 (x - x^2) \cdot dx = \int_a^b f(x) dx$

$$\begin{aligned} f(x) &= x - x^2 & a &= 0 ; b = 4 \\ \Rightarrow f(a + rh) &= f(0 + rh) & \text{and } h &= \frac{b - a}{n} \\ &= f(rh) & & h = \frac{4 - 0}{n} \\ &= (rh) - (rh)^2 & & \\ &= rh - r^2h^2 & \therefore & nh = 4 \end{aligned}$$

We know $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot [f(a + rh)]$

$$\begin{aligned} \therefore \int_0^4 (x - x^2) \cdot dx &= \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot (rh - r^2h^2) \\ &= \lim_{n \rightarrow \infty} \sum_{r=1}^n (rh^2 - r^2h^3) \\ &= \lim_{n \rightarrow \infty} \left(h^2 \cdot \sum_{r=1}^n r - h^3 \cdot \sum_{r=1}^n r^2 \right) \\ &= \lim_{n \rightarrow \infty} \left[h^2 \left(\frac{n(n+1)}{2} \right) - h^3 \left(\frac{n(n+1)(2n+1)}{6} \right) \right] \\ &= \lim_{n \rightarrow \infty} \left[\frac{h^2 \cdot n \cdot n \left(1 + \frac{1}{n} \right)}{2} - \frac{h^3 \cdot n \cdot n \left(1 + \frac{1}{n} \right) n \left(2 + \frac{1}{n} \right)}{6} \right] \\ &= \lim_{n \rightarrow \infty} \left[\frac{(nh)^2 \left(1 + \frac{1}{n} \right)}{2} - \frac{(nh)^3 \left(1 + \frac{1}{n} \right) \left(2 + \frac{1}{n} \right)}{6} \right] \\ &= \lim_{n \rightarrow \infty} \left[\frac{(4)^2 \left(1 + \frac{1}{n} \right)}{2} - \frac{(4)^3 \left(1 + \frac{1}{n} \right) \left(2 + \frac{1}{n} \right)}{6} \right] \\ &= \frac{(4)^2 \cdot (1 + 0)}{2} - \frac{(4)^3 (1 + 0) (2 + 0)}{6} \\ &= 8 - \frac{(64)(2)}{6} \\ &= -\frac{40}{3} \end{aligned}$$

Ex. 4 : $\int_0^{\frac{\pi}{2}} \sin x \cdot dx$

Solution : $\int_0^{\frac{\pi}{2}} \sin x \cdot dx = \int_0^{\frac{\pi}{2}} f(x) dx$
 $f(x) = \sin x \quad a = 0 ; b = \frac{\pi}{2}$
 $\Rightarrow f(a + rh) = \sin(a + rh)$
 $= \sin(0 + rh)$ and $h = \frac{b - a}{n} = \frac{\frac{\pi}{2} - 0}{n}$
 $= \sin rh \quad \therefore nh = \frac{\pi}{2}$

We know $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot [f(a + rh)]$

$\therefore \int_0^{\frac{\pi}{2}} \sin x \cdot dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot \sin rh$
 $= \lim_{n \rightarrow \infty} h \cdot \sum_{r=1}^n \sin rh$
 $= \lim_{n \rightarrow \infty} h \cdot [\sin h + \sin 2h + \sin 3h + \dots + \sin nh] \quad \dots \text{(I)}$

Consider,

$$\sum_{r=1}^n \sin rh = \sin h + \sin 2h + \sin 3h + \dots + \sin nh$$

$$= 2 \sin \frac{h}{2} \cdot \sin h + 2 \sin \frac{h}{2} \cdot \sin 2h + 2 \sin \frac{h}{2} \cdot \sin 3h + \dots + 2 \sin \frac{h}{2} \cdot \sin nh$$

$$\therefore 2 \sin A \cdot \sin B = \cos(A - B) - \cos(A + B)$$

$$2 \sin \frac{h}{2} \cdot \sum_{r=1}^n \sin rh = \left[\left(\cos \frac{h}{2} - \cos \frac{3h}{2} \right) + \left(\cos \frac{3h}{2} - \cos \frac{5h}{2} \right) + \left(\cos \frac{5h}{2} - \cos \frac{7h}{2} \right) + \dots \right.$$

$$\left. + \dots + \left(\cos \left(\frac{2n-1}{2} h \right) - \cos \left(\frac{2n+1}{2} h \right) \right) \right]$$

$$= \left[\cos \frac{h}{2} - \cos \left(\frac{2n+1}{2} h \right) \right]$$

$$= \left[\cos \frac{h}{2} - \cos \left(\frac{2nh}{2} + \frac{h}{2} \right) \right]$$

$$= \left[\cos \frac{h}{2} - \cos \left(\frac{\pi}{2} + \frac{h}{2} \right) \right] \quad \because nh = \frac{\pi}{2}$$

$$= \left(\cos \frac{h}{2} + \sin \frac{h}{2} \right)$$

$$\therefore \sum_{r=1}^n \sin rh = \frac{\cos \frac{h}{2} + \sin \frac{h}{2}}{2 \sin \frac{h}{2}}$$

Now from I,

$$\int_0^{\pi/2} \sin x \cdot dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n h \cdot \sin rh$$

$$= \lim_{n \rightarrow \infty} h \cdot \left[\frac{\cos \frac{h}{2} + \sin \frac{h}{2}}{2 \sin \frac{h}{2}} \right]$$

$$\therefore nh = \frac{\pi}{4} \text{ as } n \rightarrow \infty \Rightarrow h \rightarrow 0 \left(\frac{1}{n} \rightarrow 0 \right)$$

$$= \lim_{\substack{n \rightarrow \infty \\ h \rightarrow 0}} \left[\frac{\cos \frac{h}{2} + \sin \frac{h}{2}}{2 \cdot \sin \frac{h}{2}} \right]$$

$$= \frac{\cos 0 + \sin 0}{\left(\frac{1}{2} \right)}$$

$$= \frac{1 + 0}{2 \cdot \frac{1}{2}} = 1$$

$$\therefore \int_0^{\pi/2} \sin x \cdot dx = 1$$

EXERCISE 4.1

I. Evaluate the following integrals as limit of sum.

$$(1) \int_1^3 (3x - 4) \cdot dx$$

$$(2) \int_0^4 x^2 \cdot dx$$

$$(3) \int_0^2 e^x \cdot dx$$

$$(4) \int_0^2 (3x^2 - 1) \cdot dx$$

$$(5) \int_1^3 x^3 \cdot dx$$

4.2 Fundamental theorem of integral calculus :

Let f be the continuous function defined on $[a, b]$ and if $\int f(x) dx = g(x) + c$

$$\begin{aligned} \text{then } \int_a^b f(x) dx &= [g(x) + c]_a^b \\ &= [(g(b) + c) - (g(a) + c)] \\ &= g(b) + c - g(a) - c \\ &= g(b) - g(a) \end{aligned}$$

$$\text{Thus } \int_a^b f(x) dx = g(b) - g(a)$$

$$\begin{aligned} \text{Ex. : } \int_2^5 (x^2 - x) dx &= \left[\left(\frac{x^3}{3} - \frac{x^2}{2} \right) \right]_2^5 \\ &= \left[\left(\frac{5^3}{3} - \frac{5^2}{2} \right) - \left(\frac{2^3}{3} - \frac{2^2}{2} \right) \right] \\ &= \frac{125}{3} - \frac{25}{2} - \frac{8}{3} + \frac{4}{2} \\ &= \frac{117}{3} - \frac{21}{2} = \frac{234 - 83}{6} \\ \therefore \int_2^5 (x^2 - x) dx &= \frac{151}{6} \end{aligned}$$

In $\int_a^b f(x) dx$ a is called as a lower limit and b is called as an upper limit.

Now let us discuss some fundamental properties of definite integration.

These properties are very useful in evaluation of the definite integral.

4.2.1

$$\text{Property I : } \int_a^a f(x) dx = 0$$

$$\text{Let } \int f(x) dx = g(x) + c$$

$$\begin{aligned} \therefore \int_a^a f(x) dx &= [g(x) + c]_a^a \\ &= [(g(a) + c) - (g(a) + c)] \\ &= 0 \end{aligned}$$

$$\text{Property II : } \int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\text{Let } \int f(x) dx = g(x) + c$$

$$\begin{aligned} \therefore \int_a^b f(x) dx &= [g(x) + c]_a^b \\ &= [(g(b) + c) - (g(a) + c)] \\ &= g(b) - g(a) \\ &= - [g(a) - g(b)] \\ &= - \int_b^a f(x) dx \end{aligned}$$

$$\text{Thus } \int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\begin{aligned} \text{Ex. } \int_1^3 x dx &= \left[\frac{x^2}{2} \right]_1^3 \\ &= \frac{3^2}{2} - \frac{1^2}{2} = \frac{9}{2} - \frac{1}{2} = 4 \end{aligned}$$

$$\begin{aligned} \text{Ex. } \int_3^1 x dx &= \left[\frac{x^2}{2} \right]_3^1 \\ &= \frac{1^2}{2} - \frac{3^2}{2} = \frac{1}{2} - \frac{9}{2} = -4 \end{aligned}$$

Property III : $\int_a^b f(x) dx = \int_a^b f(t) dt$

Let $\int f(x) dx = g(x) + c$

L.H.S. : $\int_a^b f(x) dx = [g(x) + c]_a^b$
 $= [(g(b) + c) - (g(a) + c)]$
 $= g(b) - g(a) \dots \dots (i)$

R.H.S. : $\int_a^b f(t) dt = [g(t) + c]_a^b$
 $= [(g(b) + c) - (g(a) + c)]$
 $= g(b) - g(a) \dots \dots (ii)$

from (i) and (ii)

$$\int_a^b f(x) dx = \int_a^b f(t) dt$$

i.e. definite integration is independent of the variable.

Ex. $\int_{\pi/6}^{\pi/3} \cos x \cdot dx = \left[\sin x \right]_{\pi/6}^{\pi/3}$
 $= \sin \frac{\pi}{3} - \sin \frac{\pi}{6}$
 $= \frac{\sqrt{3}}{2} - \frac{1}{2}$
 $= \frac{\sqrt{3} - 1}{2}$

Ex. $\int_{\pi/6}^{\pi/3} \cos t \cdot dt = \left[\sin t \right]_{\pi/6}^{\pi/3}$
 $= \sin \frac{\pi}{3} - \sin \frac{\pi}{6}$
 $= \frac{\sqrt{3}}{2} - \frac{1}{2}$
 $= \frac{\sqrt{3} - 1}{2}$

Property IV : $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ where $a < c < b$ i.e. $c \in [a, b]$

Let $\int f(x) dx = g(x) + c$

Consider R.H.S. : $\int_a^c f(x) dx + \int_c^b f(x) dx$
 $= [g(x) + c]_a^c + [g(x) + c]_c^b$
 $= [(g(c) + c) - (g(a) + c)] + [(g(b) + c) - (g(c) + c)]$
 $= g(c) + c - g(a) - c + g(b) + c - g(c) - c$
 $= g(b) - g(a)$
 $= [g(x) + c]_a^b$
 $= \int_a^b f(x) dx : \text{L.H.S.}$

Thus $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ where $a < c < b$

$$\text{Ex. : } \int_{-1}^5 (2x + 3) \cdot dx = \int_{-1}^3 (2x + 3) \cdot dx + \int_3^5 (2x + 3) \cdot dx$$

$$\text{L.H.S. : } \int_{-1}^5 (2x + 3) \cdot dx$$

$$= \left[2 \frac{x^2}{2} + 3x \right]_{-1}^5$$

$$= \left[x^2 + 3x \right]_{-1}^5$$

$$= [(5)^2 + 3(5)] - [(-1)^2 + 3(-1)]$$

$$= (25 + 15) - (1 - 3)$$

$$= 40 + 2 = 42$$

$$\text{R.H.S. : } \int_{-1}^3 (2x + 3) \cdot dx + \int_3^5 (2x + 3) \cdot dx$$

$$= \left[x^2 + 3x \right]_{-1}^3 + \left[x^2 + 3x \right]_3^5$$

$$= [((3)^2 + 3(3)) - ((-1)^2 + 3(-1))] +$$

$$[((5)^2 + 3(5)) - ((3)^2 + 3(3))]$$

$$= [(9 + 9) - (1 - 3)] + [(25 + 15) - (9 - 9)]$$

$$= 18 + 2 + 40 - 18$$

$$= 42$$

$$\text{Property V : } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

$$\text{Let } \int f(x) dx = g(x) + c$$

$$\text{Consider R.H.S. : } \int_a^b f(a + b - x) dx$$

$$\text{put } a + b - x = t \quad \text{i.e.} \quad x = a + b - t$$

$$\therefore -dx = dt \Rightarrow dx = -dt$$

$$\text{As } x \rightarrow a \Rightarrow t \rightarrow b \quad \text{and} \quad x \rightarrow b \Rightarrow t \rightarrow a$$

$$\text{therefore } = \int_b^a f(t) (-dt)$$

$$= -\int_b^a f(t) dt$$

$$= \int_a^b f(t) dt \dots \left(\because \int_a^b f(x) dx = -\int_b^a f(x) dx \right)$$

$$= \int_a^b f(x) dx \quad \dots \quad \text{as definite integration is independent of the variable.}$$

$$= \text{L. H. S.}$$

$$\text{Thus } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

Ex. :

$$\int_{\pi/6}^{\pi/3} \sin^2 x \cdot dx$$

$$\text{I} = \int_{\pi/6}^{\pi/3} \sin^2 x \cdot dx \quad \dots \text{(i)}$$

$$= \int_{\pi/6}^{\pi/3} \sin^2 \left(\frac{\pi}{6} + \frac{\pi}{3} - x \right)$$

$$= \int_{\pi/6}^{\pi/3} \sin^2 \left(\frac{\pi}{2} - x \right)$$

$$\text{I} = \int_{\pi/6}^{\pi/3} \cos^2 x \cdot dx \quad \dots \text{(ii)}$$

adding (i) and (ii)

$$2\text{I} = \int_{\pi/6}^{\pi/3} \sin^2 x \cdot dx + \int_{\pi/6}^{\pi/3} \cos^2 x \cdot dx$$

$$2\text{I} = \int_{\pi/6}^{\pi/3} (\sin^2 x + \cos^2 x) \cdot dx$$

$$2\text{I} = \int_{\pi/6}^{\pi/3} 1 \cdot dx = \left[x \right]_{\pi/6}^{\pi/3}$$

$$2\text{I} = \frac{\pi}{3} - \frac{\pi}{6} = \frac{\pi}{6} \quad \therefore \text{I} = \frac{\pi}{12}$$

$$\int_{\pi/6}^{\pi/3} \sin^2 x \cdot dx = \frac{\pi}{12}$$

Property VI : $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

Let $\int f(x) dx = g(x) + c$

Consider R.H.S. : $\int_0^a f(a-x) dx$

put $a-x=t$ i.e. $x=a-t$

$\therefore -dx = dt \Rightarrow dx = -dt$

As x varies from 0 to a , t varies from a to 0

therefore I = $\int_0^a f(t) (-dt)$
 $= -\int_a^0 f(t) dt$
 $= \int_0^a f(t) dt \dots \left(\int_a^b f(x) dx = -\int_b^a f(x) dx \right)$
 $= \int_0^a f(x) dx \dots$ as definite integration is independent of the variable.
 = L. H. S.

Thus

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

Ex. : $\int_0^{\pi/4} \log(1 + \tan x) \cdot dx$

Let $\int_0^{\pi/4} \log(1 + \tan x) \cdot dx \dots (i)$

$$I = \int_0^{\pi/4} \log \left[1 + \tan \left(\frac{\pi}{4} - x \right) \right]$$

$$= \int_0^{\pi/4} \log \left[1 + \frac{\tan \frac{\pi}{4} - \tan x}{1 + \tan \frac{\pi}{4} \cdot \tan x} \right] \cdot dx$$

$$= \int_0^{\pi/4} \log \left[1 + \frac{1 - \tan x}{1 + \tan x} \right] \cdot dx$$

$$= \int_0^{\pi/4} \log \left[\frac{1 + \tan x + 1 - \tan x}{1 + \tan x} \right] \cdot dx$$

$$= \int_0^{\pi/4} \log \left[\frac{2}{1 + \tan x} \right] \cdot dx$$

$$= \int_0^{\pi/4} [\log 2 - \log(1 + \tan x)] \cdot dx$$

$$= \int_0^{\pi/4} (\log 2) \cdot dx - \int_0^{\pi/4} \log(1 + \tan x) \cdot dx$$

$I = (\log 2) \int_0^{\pi/4} 1 \cdot dx - I \dots$ by eq. (i)

$$I + I = (\log 2) \left[x \right]_0^{\pi/4}$$

$$2I = (\log 2) \left[\frac{\pi}{4} - 0 \right]$$

$\therefore I = \frac{\pi}{8} (\log 2)$

Thus

$$\int_0^{\pi/4} \log(1 + \tan x) \cdot dx = \frac{\pi}{8} (\log 2)$$

Property VII :

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$$

$$\begin{aligned} \text{R.H.S. : } & \int_0^a f(x) dx + \int_0^a f(2a-x) dx \\ & = I_1 + I_2 \quad \dots \text{(i)} \end{aligned}$$

$$\text{Consider } I_2 = \int_0^a f(2a-x) dx$$

$$\text{put } 2a-x=t \quad \text{i.e.} \quad x=2a-t$$

$$\therefore -1 dx = 1 dt \Rightarrow dx = -dt$$

As x varies from 0 to $2a$, t varies from $2a$ to 0

$$\begin{aligned} I & = \int_{2a}^0 f(t) (-dt) \\ & = -\int_{2a}^0 f(t) dt \\ & = \int_0^{2a} f(t) dt \dots \left(\int_a^b f(x) dx = -\int_b^a f(x) dx \right) \\ & = \int_0^{2a} f(x) dx \dots \left(\int_a^b f(x) dx = \int_a^b f(t) dt \right) \end{aligned}$$

$$\therefore \int_0^a f(x) dx = \int_0^{2a} f(x) dx$$

from eq. (i)

$$\begin{aligned} \int_0^a f(x) dx + \int_0^a f(2a-x) dx & = \int_0^a f(x) dx + \int_0^{2a} f(x) dx \\ & = \int_0^{2a} f(x) dx : \text{L.H.S} \end{aligned}$$

Thus,

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a-x) dx$$

Property VIII :

$$\begin{aligned} \int_{-a}^a f(x) dx & = 2 \cdot \int_0^a f(x) dx, \text{ if } f(x) \text{ even function} \\ & = 0, \text{ if } f(x) \text{ is odd function} \end{aligned}$$

$$f(x) \text{ even function if } f(-x) = f(x)$$

and $f(x)$ odd function if $f(-x) = -f(x)$

$$\int_{-a}^a f(x) dx = \int_{-a}^0 f(x) dx + \int_0^a f(x) dx \quad \dots \text{(i)}$$

$$\text{Consider } \int_{-a}^0 f(x) dx$$

$$\text{put } x = -t \quad \therefore dx = -dt$$

As x varies from $-a$ to 0 , t varies from a to 0

$$\begin{aligned} I & = \int_a^0 f(-t) (-dt) = -\int_a^0 f(-t) dt \\ & = \int_0^a f(-t) dt \dots \left(\int_a^b f(x) dx = -\int_b^a f(x) dx \right) \\ & = \int_0^a f(-x) dx \dots \left(\int_a^b f(x) dx = \int_a^b f(t) dt \right) \end{aligned}$$

Equation (i) becomes

$$\begin{aligned} \int_{-a}^a f(x) dx & = \int_0^a f(-x) dx + \int_0^a f(x) dx \\ & = \int_0^a [f(-x) + f(x)] dx \end{aligned}$$

If $f(x)$ is odd function then $f(-x) = -f(x)$, hence

$$\int_{-a}^a f(x) dx = 0$$

If $f(x)$ is even function then $f(-x) = f(x)$, hence

$$\int_{-a}^a f(x) dx = 2 \cdot \int_0^a f(x) dx$$

Hence :

$$\begin{aligned} \int_{-a}^a f(x) dx & = 2 \cdot \int_0^a f(x) dx, \text{ if } f(x) \text{ even function} \\ & = 0, \text{ if } f(x) \text{ is odd function} \end{aligned}$$

Ex. :

$$1. \int_{-\pi/4}^{\pi/4} x^3 \cdot \sin^4 x \cdot dx$$

Let $f(x) = x^3 \cdot \sin^4 x$

$$\begin{aligned} f(-x) &= (-x)^3 \cdot [\sin(-x)]^4 = -x^3 \cdot [-\sin x]^4 = -x^3 \cdot \sin^4 x \\ &= -f(x) \end{aligned}$$

$f(x)$ is odd function.

$$\therefore \int_{-\pi/4}^{\pi/4} x^3 \cdot \sin^4 x \cdot dx = 0$$

$$2. \int_{-1}^1 \frac{x^2}{1+x^2} \cdot dx$$

Let $f(x) = \frac{x^2}{1+x^2}$

$$\begin{aligned} f(-x) &= \frac{(-x)^2}{1+(-x)^2} \\ &= \frac{x^2}{1+x^2} \\ &= f(x) \end{aligned}$$

$f(x)$ is even function.

$$\begin{aligned} \int_{-1}^1 \frac{x^2}{1+x^2} \cdot dx &= 2 \int_0^1 \frac{x^2}{1+x^2} \cdot dx \\ &= 2 \int_0^1 \frac{1+x^2-1}{1+x^2} \cdot dx \\ &= 2 \int_0^1 \left[1 - \frac{1}{1+x^2}\right] \cdot dx \\ &= 2 \left[x - \tan^{-1}x\right]_0^1 \\ &= 2 \{(1 - \tan^{-1}1) - (0 - \tan^{-1}0)\} \\ &= 2 \left\{1 - \frac{\pi}{4} - 0\right\} \\ &= 2 \left(1 - \frac{\pi}{4}\right) = \left(\frac{4 - \pi}{2}\right) \end{aligned}$$

$$\therefore \int_{-1}^1 \frac{x^2}{1+x^2} \cdot dx = \frac{4 - \pi}{2}$$



SOLVED EXAMPLES

Ex. 1 : $\int_1^3 \frac{1}{\sqrt{2+x} + \sqrt{x}} \cdot dx$

Solution :
$$\begin{aligned} &= \int_1^3 \left(\frac{1}{\sqrt{2+x} + \sqrt{x}}\right) \left(\frac{\sqrt{2+x} - \sqrt{x}}{\sqrt{2+x} - \sqrt{x}}\right) \cdot dx \\ &= \int_1^3 \left(\frac{\sqrt{2+x} - \sqrt{x}}{2+x-x}\right) \cdot dx \\ &= \frac{1}{2} \cdot \int_1^3 (\sqrt{2+x} - \sqrt{x}) \cdot dx \\ &= \frac{1}{2} \cdot \left[\frac{(2+x)^{3/2}}{3/2} - \frac{x^{3/2}}{3/2}\right]_1^3 \end{aligned}$$

$$\begin{aligned} &= \frac{1}{3} \cdot \left[(2+x)^{3/2} - (x)^{3/2}\right]_1^3 \\ &= \frac{1}{3} \left\{[(2+3)^{3/2} - (3)^{3/2}] - [(2+1)^{3/2} - (1)^{3/2}]\right\} \\ &= \frac{1}{3} \left\{5^{3/2} - 3^{3/2} - 3^{3/2} + 1^{3/2}\right\} \\ &= \frac{1}{3} \left\{5^{3/2} - 2(3)^{3/2} + 1\right\} \\ \therefore \int_1^3 \frac{1}{\sqrt{2+x} + \sqrt{x}} \cdot dx &= \frac{1}{3} \left[5^{3/2} - 2(3)^{3/2} + 1\right] \end{aligned}$$

Ex. 2 : $\int_0^{\pi/2} \sqrt{1 - \cos 4x} \cdot dx$

Solution : Let $I = \int_0^{\pi/2} \sqrt{1 - \cos 4x} \cdot dx$

$$\begin{aligned} I &= \int_0^{\pi/2} \sqrt{2 \sin^2 2x} \cdot dx \\ &\left(\because 1 - \cos A = 2 \sin^2 \frac{A}{2} \right) \\ &= \sqrt{2} \cdot \int_0^{\pi/2} \sin 2x \cdot dx \\ &= \sqrt{2} \cdot \left[\frac{-\cos 2x}{2} \right]_0^{\pi/2} \\ &= \frac{\sqrt{2}}{2} \cdot \left[\cos 2 \cdot \frac{\pi}{2} - \cos 0 \right] \\ &= -\frac{\sqrt{2}}{2} \cdot [\cos \pi - \cos 0] \\ &= -\frac{\sqrt{2}}{2} \cdot (-1 - 1) = \sqrt{2} \end{aligned}$$

$\therefore \int_0^{\pi/2} \sqrt{1 - \cos 4x} \cdot dx = \sqrt{2}$

Ex. 4 : $\int_0^{\pi/4} \frac{\sec^2 x}{2 \tan^2 x + 5 \tan x + 1} \cdot dx$

Solution : Let $I = \int_0^{\pi/4} \frac{\sec^2 x}{2 \tan^2 x + 5 \tan x + 1} \cdot dx$

put $\tan x = t \quad \therefore \sec^2 x \cdot dx = 1 \cdot dt$

As x varies from 0 to $\frac{\pi}{4}$

t varies from 0 to 1

$$\begin{aligned} &= \int_0^1 \frac{1}{2t^2 + 4t + 1} \cdot dt \\ &= \frac{1}{2} \cdot \int_0^1 \frac{1}{t^2 + 2t + \frac{1}{2}} \cdot dt \\ &= \frac{1}{2} \cdot \int_0^1 \frac{1}{t^2 + 2t + 1 - 1 + \frac{1}{2}} \cdot dt \\ &= \frac{1}{2} \cdot \int_0^1 \frac{1}{(t+1)^2 - \left(\frac{1}{\sqrt{2}}\right)^2} \cdot dt \end{aligned}$$

Ex. 3 : $\int_0^{\pi/2} \cos^3 x \cdot dx$

Solution : Let $I = \int_0^{\pi/2} \cos^3 x \cdot dx$

$$\begin{aligned} &= \int_0^{\pi/2} \frac{1}{4} [\cos 3x + 3 \cos x] \cdot dx \\ &= \frac{1}{4} \left[\sin 3x \cdot \frac{1}{3} + 3 \sin x \right]_0^{\pi/2} \\ &= \frac{1}{4} \left[\left(\frac{1}{3} \sin 3 \cdot \frac{\pi}{2} + 3 \sin \frac{\pi}{2} \right) - \left(\frac{1}{3} \sin 3(0) + 3 \sin(0) \right) \right] \\ &= \frac{1}{4} \left[\frac{1}{3} \sin \frac{3\pi}{2} + 3 \sin \frac{\pi}{2} - \frac{1}{3} \sin 0 + 3 \sin 0 \right] \\ &= \frac{1}{4} \left[\frac{1}{3} (-1) + 3(1) - 0 \right] \\ &= \frac{1}{4} \left[-\frac{1}{3} + 3 \right] = \frac{1}{4} \left[\frac{8}{3} \right] = \frac{2}{3} \end{aligned}$$

$\therefore \int_0^{\pi/2} \cos^3 x \cdot dx = \frac{2}{3}$

$$\begin{aligned} &= \frac{1}{2} \cdot \frac{1}{2 \left(\frac{1}{\sqrt{2}}\right)} \left[\log \left[\frac{(t+1) - \frac{1}{\sqrt{2}}}{(t+1) + \frac{1}{\sqrt{2}}} \right] \right]_0^1 \\ &= \frac{\sqrt{2}}{4} \log \left[\left(\frac{\sqrt{2}t + \sqrt{2} - 1}{\sqrt{2}t + \sqrt{2} + 1} \right) \right]_0^1 \\ &= \frac{\sqrt{2}}{4} \left[\log \left(\frac{\sqrt{2}(1) + \sqrt{2} - 1}{\sqrt{2}(1) + \sqrt{2} + 1} \right) - \log \left(\frac{\sqrt{2}(0) + \sqrt{2} - 1}{\sqrt{2}(0) + \sqrt{2} + 1} \right) \right] \\ &= \frac{\sqrt{2}}{4} \left[\log \left(\frac{2\sqrt{2} - 1}{2\sqrt{2} + 1} \right) - \log \left(\frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right) \right] \\ &= \frac{\sqrt{2}}{4} \log \left[\left(\frac{2\sqrt{2} - 1}{2\sqrt{2} + 1} \right) \div \left(\frac{\sqrt{2} - 1}{\sqrt{2} + 1} \right) \right] \\ &= \frac{\sqrt{2}}{4} \log \left[\frac{3 + \sqrt{2}}{3 - \sqrt{2}} \right] \end{aligned}$$

Ex. 5 : $\int_1^2 \frac{\log x}{x^2} \cdot dx$

Solution : Let $I = \int_1^2 (\log x) \left(\frac{1}{x^2}\right) \cdot dx$

$$= \left[(\log x) \cdot \int \frac{1}{x^2} \cdot dx \right]_1^2 - \int_1^2 \frac{d}{dx} \log x \cdot \int \frac{1}{x^2} \cdot dx \cdot dx$$

$$= \left[(\log x) \cdot \left(-\frac{1}{x}\right) \right]_1^2 - \int_1^2 \frac{1}{x} \cdot \left(-\frac{1}{x}\right) \cdot dx$$

$$= \left[-\frac{1}{x} \log x \right]_1^2 + \int_1^2 \frac{1}{x^2} \cdot dx$$

$$= \left[-\frac{1}{x} \log x \right]_1^2 + \left[-\frac{1}{x} \right]_1^2$$

$$= \left[\left(-\frac{1}{2} \log 2\right) - \left(-\frac{1}{1} \log 1\right) \right] + \left[\left(-\frac{1}{2}\right) - \left(-\frac{1}{1}\right) \right]$$

$$= -\frac{1}{2} \log 2 - 0 - \frac{1}{2} + 1 = \frac{1}{2} - \frac{1}{2} \log 2 \quad \because \log 1 = 0$$

$$\therefore \int_1^2 \frac{\log x}{x^2} \cdot dx = \frac{1}{2} (1 - \log 2)$$

Ex. 6 : $\int_0^{\pi/2} \frac{\cos x}{1 + \cos x + \sin x} \cdot dx$

Solution : Let $I = \int_0^{\pi/2} \frac{\cos x}{1 + \cos x + \sin x} \cdot dx$

$$= \int_0^{\pi/2} \frac{\cos^2\left(\frac{x}{2}\right) - \sin^2\left(\frac{x}{2}\right)}{2 \cos^2\left(\frac{x}{2}\right) + 2 \sin\left(\frac{x}{2}\right) \cdot \cos\left(\frac{x}{2}\right)} \cdot dx$$

$$= \int_0^{\pi/2} \frac{\left[\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right] \left[\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right]}{2 \left[\cos\left(\frac{x}{2}\right)\right] \left[\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right]} \cdot dx$$

$$= \int_0^{\pi/2} \left[\frac{\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right)} \right] \cdot dx = \int_0^{\pi/2} \left[1 - \tan\left(\frac{x}{2}\right) \right] \cdot dx$$

$$\begin{aligned}
&= \frac{1}{2} \cdot \left[x - \log \left(\sec \frac{x}{2} \right) \cdot \frac{1}{\frac{1}{2}} \right]_0^{\pi/2} \\
&= \frac{1}{2} \cdot \left[\frac{\pi}{2} - 2 \cdot \log \left(\sec \frac{\pi}{4} \right) - (0 - 2 \log \sec 0) \right] \\
&= \frac{1}{2} \cdot \left[\frac{\pi}{2} - 2 \log \sqrt{2} - 0 + 2(0) \right] = \frac{1}{2} \cdot \left[\frac{\pi}{2} - 2 \log \sqrt{2} \right] = \frac{\pi}{4} - \log \sqrt{2} \\
\therefore \int_0^{\pi/2} \frac{\sec^2 x}{1 + \cos x + \sin x} \cdot dx &= \frac{\pi}{4} - \log \sqrt{2}
\end{aligned}$$

Ex. 7 : $\int_0^{1/2} \frac{1}{(1-2x^2)\sqrt{1-x^2}} \cdot dx$

Solution : Let $I = \int_0^{1/2} \frac{1}{(1-2x^2)\sqrt{1-x^2}} \cdot dx$

put $x = \sin \theta \quad \therefore \quad 1 \cdot dx = \cos \theta \cdot d\theta$

As x varies from 0 to $\frac{1}{2}$, θ varies from 0 to $\frac{\pi}{6}$

$$= \int_0^{\pi/6} \frac{\cos \theta}{(1-2\sin^2 \theta)\sqrt{1-\sin^2 \theta}} \cdot d\theta = \int_0^{\pi/6} \frac{\cos \theta}{(\cos 2\theta)\sqrt{\cos^2 \theta}} \cdot d\theta$$

$$= \int_0^{\pi/6} \frac{1}{\cos 2\theta} \cdot d\theta$$

$$= \int_0^{\pi/6} \sec 2\theta \cdot d\theta$$

$$= \left[\log (\sec 2\theta + \tan 2\theta) \cdot \frac{1}{2} \right]_0^{\pi/6}$$

$$= \frac{1}{2} \cdot \left[\log \left(\sec 2 \left(\frac{\pi}{6} \right) + \tan 2 \left(\frac{\pi}{6} \right) \right) - \log (\sec 0 + \tan 0) \right]$$

$$= \frac{1}{2} \cdot \left[\log \left(\sec \frac{\pi}{3} + \tan \frac{\pi}{3} \right) - \log (1 + 0) \right] \quad \because \log 1 = 0$$

$$= \frac{1}{2} \cdot [\log (2 + \sqrt{3}) - 0]$$

$$= \frac{1}{2} \log (2 + \sqrt{3})$$

$$\therefore \int_0^{1/2} \frac{1}{(1-2x^2)\sqrt{1-x^2}} \cdot dx = \frac{1}{2} \log (2 + \sqrt{3})$$

Ex. 8 : $\int_0^2 \frac{2^x}{2^x(1+4^x)} \cdot dx$

Solution : Let $I = \int_0^2 \frac{2^x}{2^x(1+4^x)} \cdot dx$

put $2^x = t \quad \therefore \quad 2^x \cdot \log 2 \cdot dx = 1 \cdot dt$

As x varies from 0 to 2, t varies from 1 to 4

$$= \int_1^4 \frac{1}{\log 2} \cdot \frac{1}{t(1+t^2)} \cdot dt$$

$$= \frac{1}{\log 2} \cdot \int_1^4 \frac{1}{t(1+t^2)} \cdot dt$$

$$= \frac{1}{\log 2} \cdot \int_1^4 \frac{1+t^2-t^2}{t(1+t^2)} \cdot dt$$

may be solved by method of partial fraction

$$= \frac{1}{\log 2} \cdot \int_1^4 \left[\frac{1+t^2}{t(1+t^2)} - \frac{t^2}{t(1+t^2)} \right] \cdot dt$$

$$= \frac{1}{\log 2} \cdot \int_1^4 \left[\frac{1}{t} - \frac{t}{1+t^2} \right] \cdot dt$$

$$= \frac{1}{\log 2} \cdot \left[\int_1^4 \frac{1}{t} \cdot dt - \frac{1}{2} \int_1^4 \frac{2t}{1+t^2} \cdot dt \right]$$

$$= \frac{1}{\log 2} \cdot \left[\log(t) - \frac{1}{2} \log(1+t^2) \right]_1^4$$

$$= \frac{1}{\log 2} \cdot \left[\left(\log 4 - \frac{1}{2} \log 17 \right) - \left(\log 1 - \frac{1}{2} \log 2 \right) \right]$$

$$= \frac{1}{\log 2} \cdot \left[\log 4 - \frac{1}{2} \log 17 + \frac{1}{2} \log 2 \right]$$

$$\because \log 1 = 0$$

$$= \frac{1}{\log 2} \cdot \left[\log \frac{4\sqrt{2}}{\sqrt{17}} \right]$$

$$\therefore \int_0^2 \frac{2^x}{2^x(1+4^x)} \cdot dx = \frac{1}{(\log 2)} \cdot \left[\log \frac{4\sqrt{2}}{\sqrt{17}} \right]$$

$$= \log_2 \left(\frac{4\sqrt{2}}{\sqrt{17}} \right)$$

Ex. 9 : $\int_{-1}^1 |5x-3| \cdot dx$

Solution : Let $I = \int_{-1}^1 |5x-3| \cdot dx$

$$|5x-3| = -(5x-3) \text{ for } (5x-3) < 0 \text{ i.e. } x < \frac{3}{5}$$

$$= (5x-3) \text{ for } (5x-3) > 0 \text{ i.e. } x > \frac{3}{5}$$

$$= \int_{-1}^{\frac{3}{5}} |5x-3| \cdot dx + \int_{\frac{3}{5}}^1 |5x-3| \cdot dx$$

$$= \int_{-1}^{\frac{3}{5}} -(5x-3) \cdot dx + \int_{\frac{3}{5}}^1 (5x-3) \cdot dx$$

$$= \left[-\left(5 \frac{x^2}{2} - 3x\right) \right]_{-1}^{\frac{3}{5}} + \left[\left(5 \frac{x^2}{2} - 3x\right) \right]_{\frac{3}{5}}^1$$

$$= \left[3x - \frac{5}{2}x^2 \right]_{-1}^{\frac{3}{5}} + \left[\frac{5}{2}x^2 - 3x \right]_{\frac{3}{5}}^1$$

$$= \left[\left(3 \left(\frac{3}{5} \right) - \frac{5}{2} \left(\frac{3}{5} \right)^2 \right) - \left(3(-1) - \frac{5}{2}(-1)^2 \right) \right] + \left[\left(\frac{5}{2}(1)^2 - 3(1) \right) - \left(\frac{5}{2} \left(\frac{3}{5} \right)^2 - 3 \left(\frac{3}{5} \right) \right) \right]$$



$$= \left[\left(\frac{9}{5} - \frac{9}{10} \right) - \left(-3 - \frac{5}{2} \right) \right] + \left[\left(\frac{5}{2} - 3 \right) - \left(\frac{9}{10} - \frac{9}{5} \right) \right]$$

$$= \frac{9}{5} - \frac{9}{10} + 3 + \frac{5}{2} + \frac{5}{2} - 3 - \frac{9}{10} + \frac{9}{5} = 2 \left(\frac{9}{5} - \frac{9}{10} + \frac{5}{2} \right) = 2 \left(\frac{18 - 9 + 25}{5} \right) = \frac{34}{5}$$

$$\therefore \int_{-1}^1 |5x - 3| \cdot dx = \frac{34}{5}$$

Ex. 10 : $\int_0^{\pi/2} \frac{1}{1 + \sqrt[3]{\tan x}} \cdot dx$

Solution : Let $I = \int_0^{\pi/2} \frac{1}{1 + \sqrt[3]{\tan x}} \cdot dx$

$$= \int_0^{\pi/2} \left[\frac{1}{1 + \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\cos x}}} \right] \cdot dx$$

$$= \int_0^{\pi/2} \frac{\sqrt[3]{\cos x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} \cdot dx \quad \dots (i)$$

By property $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

$$I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos \left(\frac{\pi}{2} - x \right)}}{\sqrt[3]{\cos \left(\frac{\pi}{2} - x \right)} + \sqrt[3]{\sin \left(\frac{\pi}{2} - x \right)}} \cdot dx$$

$$= \int_0^{\pi/2} \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} \cdot dx \quad \dots (ii)$$

adding (i) and (ii)

$$I + I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} \cdot dx + \int_0^{\pi/2} \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} \cdot dx$$

$$2I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} \cdot dx$$

$$2I = \int_0^{\pi/2} 1 \cdot dx$$

$$I = \frac{1}{2} \left[x \right]_0^{\pi/2} = \frac{1}{2} \left[\frac{\pi}{4} - 0 \right] = \frac{\pi}{4}$$

$$\therefore \int_0^{\pi/2} \frac{1}{1 + \sqrt[3]{\tan x}} \cdot dx = \frac{\pi}{4}$$

with the help of the above solved/ illustrative example verify whether the following examples evaluates their definite integrate to be equal to / as $\frac{\pi}{4}$

$$\int_0^{\pi/2} \frac{1}{1 + \cot^3 x} \cdot dx ;$$

$$\int_0^{\pi/2} \frac{\sin x}{\sin x + \cos x} \cdot dx ;$$

$$\int_0^{\pi/2} \frac{\sec x}{\sec x + \operatorname{cosec} x} \cdot dx ;$$

$$\int_0^{\pi/2} \frac{\sin^4 x}{\sin^4 x + \cos^4 x} \cdot dx ;$$

$$\int_0^{\pi/2} \frac{\operatorname{cosec}^{\frac{5}{2}} x}{\operatorname{cosec}^{\frac{5}{2}} x + \sec^{\frac{5}{2}} x} \cdot dx$$

Ex. 11 : $\int_3^8 \frac{(11-x)^2}{x^2 + (1-x)^2} \cdot dx$

Solution : Let $I = \int_3^8 \frac{(11-x)^2}{x^2 + (1-x)^2} \cdot dx \quad \dots (i)$

By property $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

$$I = \int_3^8 \frac{[11-(8+3-x)]^2}{[8+3-x]^2 + [11-(8+3-x)]^2} \cdot dx = \int_3^8 \frac{[11-(11-x)]^2}{(11-x)^2 + [11-(11-x)]^2} \cdot dx$$

$$= \int_3^8 \frac{x^2}{(11-x)^2 + x^2} \cdot dx \quad \dots (ii)$$

adding (i) and (ii)

$$I + I = \int_3^8 \frac{(11-x)^2}{x^2 + (1-x)^2} \cdot dx + \int_3^8 \frac{x^2}{(11-x)^2 + x^2} \cdot dx$$

$$2I = \int_3^8 \frac{(11-x)^2 + x^2}{x^2 + (11-x)^2} \cdot dx$$

$$I = \frac{1}{2} \int_3^8 1 \cdot dx$$

$$I = \frac{1}{2} \left[x \right]_3^8 = \frac{1}{2} [8 - 3] = \frac{5}{2}$$

$$\therefore \int_3^8 \frac{(11-x)^2}{x^2 + (1-x)^2} \cdot dx = \frac{5}{2}$$

Note that : In general $\int_a^b \frac{f(x)}{f(x) + f(a+b-x)} \cdot dx = \frac{1}{2} (b-a)$

verify the generalisation for the following examples :

$$\int_1^2 \frac{\sqrt{x}}{\sqrt{3-x} + \sqrt{x}} \cdot dx ;$$

$$\int_2^7 \frac{x^3}{(9-x)^3 + x^3} \cdot dx ;$$

$$\int_4^9 \frac{x^{\frac{1}{4}}}{(13-x)^{\frac{1}{4}} + x^{\frac{1}{4}}} \cdot dx$$

$$\int_{\pi/6}^{\pi/3} \frac{1}{1 + \sqrt{\cot x}} \cdot dx$$

$$\int_{\pi/6}^{\pi/3} \frac{1}{1 + \sqrt{\operatorname{cosec} x}} \cdot dx$$

Ex. 12 : $\int_0^{\pi} x \cdot \sin^2 x \cdot dx$

Solution :

Consider, $I = \int_0^{\pi} x \cdot \sin^2 x \cdot dx \dots (i)$

$$I = \int_0^{\pi} (\pi - x) \cdot [\sin(\pi - x)]^2 x \cdot dx$$

$$I = \int_0^{\pi} (\pi - x) \cdot \sin^2 x \cdot dx$$

$$I = \int_0^{\pi} \pi \cdot \sin^2 x \cdot dx - \int_0^{\pi} x \cdot \sin^2 x \cdot dx$$

$$I = \pi \cdot \int_0^{\pi} \frac{1}{2} (1 - \cos 2x) \cdot dx - I \dots \text{by (i)}$$

$$I + I = \frac{\pi}{2} \int_0^{\pi} (1 - \cos 2x) \cdot dx$$

$$2I = \frac{\pi}{2} \left[x - \sin 2x \cdot \frac{1}{2} \right]_0^{\pi}$$

$$I = \frac{\pi}{4} \left[\left(\pi - \frac{1}{2} \sin 2\pi \right) - \left(0 - \frac{1}{2} \sin 0 \right) \right]$$

$$= \frac{\pi}{4} [\pi] \quad \because \sin 0 = 0; \sin 2\pi = 0$$

$$= \frac{\pi^2}{4}$$

$$\therefore \int_0^{\pi} x^2 \cdot \sin^2 x \cdot dx = \frac{\pi^2}{4}$$

Ex. 13 : Evaluate the integral $\int_0^{\pi} \cos^2 x \cdot dx$ using the result/ property.

Solution :

$$\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a - x) dx$$

Let, $I = \int_0^{\pi} \cos^2 x \cdot dx$

$$= \int_0^{2\left(\frac{\pi}{2}\right)} \cos^2 x \cdot dx$$

$$= \int_0^{\pi/2} \cos^2 x \cdot dx + \int_0^{\pi/2} \left[\cos \left(2\frac{\pi}{2} - x \right) \right]^2 \cdot dx$$

$$= \int_0^{\pi/2} \cos^2 x \cdot dx + \int_0^{\pi/2} \cos^2 x \cdot dx$$

$$\because \cos(\pi - x) = -\cos x$$

$$= 2 \cdot \int_0^{\pi/2} \cos^2 x \cdot dx$$

$$= \int_0^{\pi/2} (1 + \cos 2x) \cdot dx$$

$$= \left[x + \sin 2x \cdot \frac{1}{2} \right]_0^{\pi/2}$$

$$= \left[\left(\frac{\pi}{2} + \frac{1}{2} \sin 2\frac{\pi}{2} \right) - \left(0 + \frac{1}{2} \sin 2(0) \right) \right]$$

$$= \frac{\pi}{2} + 0 \quad \because \sin 0 = 0; \sin \pi = 0$$

$$= \frac{\pi}{2}$$

$$\therefore \int_0^{\pi} \cos^2 x \cdot dx = \frac{\pi}{2}$$

Ex. 14 : $\int_{-\pi}^{\pi} \frac{x(1 + \sin x)}{1 + \cos^2 x} \cdot dx$

Solution : Let $I = \int_{-\pi}^{\pi} \frac{x(1 + \sin x)}{1 + \cos^2 x} \cdot dx$

$$= \left[\left(\int_{-\pi}^{\pi} \frac{x}{1 + \cos^2 x} \cdot dx \right) + \left(\int_{-\pi}^{\pi} \frac{x \cdot \sin x}{1 + \cos^2 x} \cdot dx \right) \right]$$

The function $\frac{x}{1 + \cos^2 x}$ is odd function and the function $\frac{x \cdot \sin x}{1 + \cos^2 x}$ is even function.

$$\int_{-a}^a f(x) dx = 2 \cdot \int_0^a f(x) dx, \text{ if } f(x) \text{ even function}$$

$$= 0, \text{ if } f(x) \text{ is odd function}$$

$$\therefore I = 0 + 2 \cdot \int_0^{\pi} \frac{x \cdot \sin x}{1 + \cos^2 x} \cdot dx$$

$$\therefore I = 2 \cdot \int_0^{\pi} \frac{x \cdot \sin x}{1 + \cos^2 x} \cdot dx \quad \dots (i)$$

$$= 2 \cdot \int_0^{\pi} \frac{(\pi - x) \cdot \sin(\pi - x)}{1 + [\cos(\pi - x)]^2} \cdot dx$$

$$= 2 \cdot \int_0^{\pi} \frac{(\pi - x) \cdot \sin x}{1 + (-\cos x)^2} \cdot dx$$

$$= 2\pi \cdot \int_0^{\pi} \frac{\pi \cdot \sin x - x \cdot \sin x}{1 + \cos^2 x} \cdot dx$$

$$= 2\pi \cdot \int_0^{\pi} \frac{\sin x}{1 + \cos^2 x} - 2 \cdot \int_0^{\pi} \frac{x \cdot \sin x}{1 + \cos^2 x} \cdot dx$$

$$I = 2\pi \cdot \int_0^{\pi} \frac{\sin x}{1 + \cos^2 x} - I \quad \dots \text{ by eq.(i)}$$

$$I + I = 2\pi \cdot \int_0^{\pi} \frac{\sin x}{1 + \cos^2 x} \quad \dots (ii)$$

put $\cos x = t \quad \therefore -\sin x \cdot dx = + dt$

As x varies from 0 to π , t varies from 1 to -1

$$2I = 2\pi \cdot \int_{-1}^1 \frac{-1}{1 + t^2} \cdot dt$$

$$I = \pi \cdot 2 \int_0^1 \frac{1}{1 + t^2} \cdot dt \quad \left(\text{where } \frac{1}{1 + t^2} \text{ is even function.} \right)$$



$$\begin{aligned}
 I &= 2\pi \cdot \left[\tan^{-1} t \right]_0^1 \\
 &= 2\pi [\tan^{-1}(1) - \tan^{-1}(0)] \\
 &= 2\pi \left(\frac{\pi}{4} - 0 \right) = \frac{\pi^2}{2}
 \end{aligned}$$

$$\therefore \int_{-\pi}^{\pi} \frac{x(1 + \sin x)}{1 + \cos^2 x} \cdot dx = \frac{\pi^2}{2}$$

Ex. 15 : $\int_0^3 x [x] \cdot dx$, where $[x]$ denote greatest integrate function not greater than x .

Solution : Let $I = \int_0^3 x [x] \cdot dx$

$$\begin{aligned}
 I &= \int_0^1 x [x] \cdot dx + \int_1^2 x [x] \cdot dx + \int_2^3 x [x] \cdot dx \\
 &= \int_0^1 x(0) \cdot dx + \int_1^2 x(1) \cdot dx + \int_2^3 x(2) \cdot dx \\
 &= 0 + \left[\frac{x^2}{2} \right]_1^2 + \left[x^2 \right]_2^3 \\
 &= 0 + \left(\frac{4}{2} - \frac{1}{2} \right) + (9 - 4) \\
 &= \frac{3}{2} + 5 = \frac{13}{2}
 \end{aligned}$$

$$\therefore \int_0^3 x [x] \cdot dx = \frac{13}{2}$$

EXERCISE 4.2

I. Evaluate :

(1) $\int_1^9 \frac{x+1}{\sqrt{x}} \cdot dx$

(2) $\int_2^3 \frac{1}{x^2 + 5x + 6} \cdot dx$

(8) $\int_0^{\pi/4} \sqrt{1 + \sin 2x} \cdot dx$

(9) $\int_0^{\pi/4} \sin^4 x \cdot dx$

(3) $\int_0^{\pi/4} \cot^2 \cdot dx$

(4) $\int_{-\pi/4}^{\pi/4} \frac{1}{1 - \sin x} \cdot dx$

(10) $\int_{-4}^2 \frac{1}{x^2 + 4x + 13} \cdot dx$

(11) $\int_0^4 \frac{1}{\sqrt{4x - x^2}} \cdot dx$

(5) $\int_3^5 \frac{1}{\sqrt{2x+3} - \sqrt{2x-3}} \cdot dx$

(12) $\int_0^1 \frac{1}{\sqrt{3+2x-x^2}} \cdot dx$

(13) $\int_0^{\pi/2} x \cdot \sin x \cdot dx$

(6) $\int_0^1 \frac{x^2-2}{x^2+1} \cdot dx$

(7) $\int_0^{\pi/4} \sin 4x \sin 3x \cdot dx$

(14) $\int_0^1 x \cdot \tan^{-1} x \cdot dx$

(15) $\int_0^{\infty} x \cdot e^{-x} \cdot dx$

II. Evaluate :

$$(1) \int_0^{\frac{1}{\sqrt{2}}} \frac{\sin^{-1} x}{(1-x^2)^{\frac{3}{2}}} dx$$

$$(2) \int_0^{\frac{\pi}{4}} \frac{\sec^2 x}{3 \tan^2 x + 4 \tan x + 1} dx$$

$$(3) \int_0^{4\pi} \frac{\sin 2x}{\sin^4 x + \cos^4 x} dx$$

$$(4) \int_0^{2\pi} \sqrt{\cos x} \cdot \sin^3 x \cdot dx$$

$$(5) \int_0^{\frac{\pi}{2}} \frac{1}{5 + 4 \cos x} dx$$

$$(6) \int_0^{\frac{\pi}{4}} \frac{\cos x}{4 - \sin^2 x} dx$$

$$(7) \int_0^{\frac{\pi}{2}} \frac{\cos x}{(1 + \sin x)(2 + \sin x)} dx$$

$$(8) \int_{-1}^1 \frac{1}{a^2 e^x + b^2 e^{-x}} dx$$

$$(9) \int_0^{\pi} \frac{1}{3 + 2 \sin x + \cos x} dx$$

$$(10) \int_0^{\frac{\pi}{4}} \sec^4 x \cdot dx$$

$$(11) \int_0^1 \sqrt{\frac{1-x}{1+x}} dx$$

$$(12) \int_0^{\pi} \sin^3 x (1 + 2 \cos x) (1 + \cos x)^2 \cdot dx$$

$$(13) \int_0^{\frac{\pi}{2}} \sin 2x \cdot \tan^{-1}(\sin x) \cdot dx$$

$$(14) \int_{\frac{1}{\sqrt{2}}}^1 \frac{(e^{\cos^{-1} x})(\sin^{-1} x)}{\sqrt{1-x^2}} dx$$

$$(15) \int_2^3 \frac{\cos(\log x)}{x} dx$$

III. Evaluate :

$$(1) \int_0^a \frac{1}{x + \sqrt{a^2 - x^2}} dx$$

$$(2) \int_0^{\frac{\pi}{2}} \log \tan x \cdot dx$$

$$(3) \int_0^1 \log \left(\frac{1}{x} - 1 \right) dx$$

$$(4) \int_0^{\frac{\pi}{2}} \frac{\sin x - \cos x}{1 + \sin x \cdot \cos x} dx$$

$$(5) \int_0^3 x^2 (3-x)^{\frac{5}{2}} dx$$

$$(6) \int_{-3}^3 \frac{x^3}{9-x^2} dx$$

$$(7) \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \log \left(\frac{2 + \sin x}{2 - \sin x} \right) dx$$

$$(8) \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{x + \frac{\pi}{4}}{2 - \cos 2x} dx$$

$$(9) \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} x^3 \cdot \sin^4 x \cdot dx$$

$$(10) \int_0^1 \frac{\log(x+1)}{x^2+1} dx$$

$$(11) \int_{-1}^1 \frac{x^3+2}{\sqrt{x^2+4}} dx$$

$$(12) \int_{-a}^a \frac{x+x^3}{16-x^2} dx$$

$$(13) \int_0^1 t^2 \sqrt{1-t} \cdot dx$$

$$(14) \int_0^{\pi} x \cdot \sin x \cdot \cos^2 x \cdot dx$$

$$(15) \int_0^1 \frac{\log x}{\sqrt{1-x^2}} dx$$

Note that :

To evaluate the integrals of the type $\int_0^{\pi/2} \sin^n x \cdot dx$ and $\int_0^{\pi/2} \cos^n x \cdot dx$, the results used are known as

'reduction formulae' which are stated as follows :

$$\int_0^{\pi/2} \sin^n x \cdot dx = \frac{(n-1) \cdot (n-3) \cdot (n-5) \dots \frac{4}{5} \frac{2}{3}}, \quad \text{if } n \text{ is odd.}$$

$$= \frac{(n-1) \cdot (n-3) \cdot (n-5) \dots \frac{3}{4} \frac{1}{2} \cdot \frac{\pi}{2}}, \quad \text{if } n \text{ is even.}$$

$$\int_0^{\pi/2} \cos^n x \cdot dx = \int_0^{\pi/2} \left[\cos \left(\frac{\pi}{2} - x \right) \right]^n \cdot dx \quad \dots \text{ by property}$$

$$= \int_0^{\pi/2} [\sin x]^n \cdot dx$$

$$= \int_0^{\pi/2} \sin^n x \cdot dx$$

$$\int_0^{\pi/2} \sin^7 x \cdot dx = \frac{(7-1) \cdot (7-3) \cdot (7-5)}{7 \cdot (7-2) \cdot (7-4)}$$

$$= \frac{(7-1) \cdot (7-3) \cdot (7-5)}{7 \cdot (7-2) \cdot (7-4)}$$

$$= \frac{6 \cdot 4 \cdot 2}{7 \cdot 5 \cdot 3} = \frac{16}{35}$$

$$\int_0^{\pi/2} \cos^8 x \cdot dx = \frac{(8-1) \cdot (8-3) \cdot (8-5) \cdot (8-7)}{8 \cdot (8-2) \cdot (8-4) \cdot (8-6)} \cdot \frac{\pi}{2}$$

$$= \frac{(8-1) \cdot (8-3) \cdot (8-5) \cdot (8-7)}{8 \cdot (8-2) \cdot (8-4) \cdot (8-6)} \cdot \frac{\pi}{2}$$

$$= \frac{7 \cdot 5 \cdot 3 \cdot 1}{8 \cdot 6 \cdot 4 \cdot 2} \cdot \frac{\pi}{2}$$

$$= \frac{35\pi}{256}$$



Let us Remember

$$\ast \sum_{r=0}^{n-1} (x_{r+1} - x_r) \cdot f(t_r) = \sum_{r=0}^{n-1} g(x_{r+1}) - g(x_r) = g(b) - g(a)$$

Thus taking limit as $n \rightarrow \infty$

$$g(b) - g(a) = \lim_{n \rightarrow \infty} \sum (x_{r+1} - x_r) \cdot f(t_r) = \lim_{n \rightarrow \infty} S_n = \int_a^b f(x) dx$$

$$\ast \text{Fundamental theorem of integral calculus : } \int_a^b f(x) dx = g(b) - g(a)$$

$$\text{Property I : } \int_a^a f(x) dx = 0$$

$$\text{Property II : } \int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\text{Property III : } \int_a^b f(x) dx = \int_a^b f(t) dt$$

$$\text{Property IV : } \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx \quad \text{where } a < c < b \text{ i.e. } c \in [a, b]$$

$$\text{Property V : } \int_a^b f(x) dx = \int_a^b f(a + b - x) dx$$

$$\text{Property VI : } \int_0^a f(x) dx = \int_0^a f(a - x) dx$$

$$\text{Property VII : } \int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a - x) dx$$

$$\text{Property VIII : } \int_{-a}^a f(x) dx = 2 \cdot \int_0^a f(x) dx \quad , \text{ if } f(x) \text{ even function}$$

$$= 0 \quad , \text{ if } f(x) \text{ is odd function}$$

$f(x)$ even function if $f(-x) = f(x)$ and $f(x)$ odd function if $f(-x) = -f(x)$

\ast 'Reduction formulae' which are stated as follows :

$$\int_0^{\pi/2} \sin^n x \cdot dx = \frac{(n-1)(n-3)(n-5) \dots 4}{n(n-2)(n-4) \dots 5} \cdot \frac{2}{3}, \quad \text{if } n \text{ is odd.}$$

$$= \frac{(n-1)(n-3)(n-5) \dots 3}{n(n-2)(n-4) \dots 4} \cdot \frac{1}{2} \cdot \frac{\pi}{2}, \quad \text{if } n \text{ is even.}$$

$$\int_0^{\pi/2} \cos^n x \cdot dx = \int_0^{\pi/2} \left[\cos \left(\frac{\pi}{2} - 0 \right) \right]^n \cdot dx = \int_0^{\pi/2} [\sin x]^n \cdot dx = \int_0^{\pi/2} \sin^n x \cdot dx$$

MISCELLANEOUS EXERCISE 4

(I) Choose the correct option from the given alternatives :

- (1) $\int_2^3 \frac{dx}{x(x^3-1)} =$
 (A) $\frac{1}{3} \log\left(\frac{208}{189}\right)$ (B) $\frac{1}{3} \log\left(\frac{189}{208}\right)$ (C) $\log\left(\frac{208}{189}\right)$ (D) $\log\left(\frac{189}{208}\right)$
- (2) $\int_0^{\pi/2} \frac{\sin^2 x \cdot dx}{(1 + \cos x)^2} =$
 (A) $\frac{4 - \pi}{2}$ (B) $\frac{\pi - 4}{2}$ (C) $4 - \frac{\pi}{2}$ (D) $\frac{4 + \pi}{2}$
- (3) $\int_0^{\log 5} \frac{e^x \sqrt{e^x - 1}}{e^x + 3} \cdot dx =$
 (A) $3 + 2\pi$ (B) $4 - \pi$ (C) $2 + \pi$ (D) $4 + \pi$
- (4) $\int_0^{\pi/2} \sin^6 x \cos^2 x \cdot dx =$
 (A) $\frac{7\pi}{256}$ (B) $\frac{3\pi}{256}$ (C) $\frac{5\pi}{256}$ (D) $\frac{-5\pi}{256}$
- (5) If $\int_0^1 \frac{dx}{\sqrt{1+x} - \sqrt{x}} = \frac{k}{3}$, then k is equal to
 (A) $\sqrt{2}(2\sqrt{2} - 2)$ (B) $\frac{\sqrt{2}}{3}(2 - 2\sqrt{2})$ (C) $\frac{2\sqrt{2} - 2}{3}$ (D) $4\sqrt{2}$
- (6) $\int_1^2 \frac{1}{x^2} e^{\frac{1}{x}} \cdot dx =$
 (A) $\sqrt{e} + 1$ (B) $\sqrt{e} - 1$ (C) $\sqrt{e}(\sqrt{e} - 1)$ (D) $\frac{\sqrt{e} - 1}{e}$
- (7) If $\int_2^e \left[\frac{1}{\log x} - \frac{1}{(\log x)^2} \right] \cdot dx = a + \frac{b}{\log 2}$, then
 (A) $a = e, b = -2$ (B) $a = e, b = 2$ (C) $a = -e, b = 2$ (D) $a = -e, b = -2$
- (8) Let $I_1 = \int_e^{e^2} \frac{dx}{\log x}$ and $I_2 = \int_1^2 \frac{e^x}{x} \cdot dx$, then
 (A) $I_1 = \frac{1}{3} I_2$ (B) $I_1 + I_2 = 0$ (C) $I_1 = 2I_2$ (D) $I_1 = I_2$

(9) $\int_0^9 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{9-x}} \cdot dx =$

- (A) 9 (B) $\frac{9}{2}$ (C) 0 (D) 1

(10) The value of $\int_{-\pi/4}^{\pi/4} \log \left(\frac{2 + \sin \theta}{2 - \sin \theta} \right) \cdot d\theta$ is

- (A) 0 (B) 1 (C) 2 (D) π

(II) Evaluate the following :

(1) $\int_0^{\pi/2} \frac{\cos x}{3 \cdot \cos x + \sin x} \cdot dx$

(2) $\int_{\pi/4}^{\pi/2} \frac{\cos \theta}{\left[\cos \frac{\theta}{2} + \sin \frac{\theta}{2} \right]^3} \cdot d\theta$

(3) $\int_0^1 \frac{1}{1 + \sqrt{x}} \cdot dx$

(4) $\int_0^{\pi/4} \frac{\tan^3 x}{1 + \cos 2x} \cdot dx$

(5) $\int_0^1 t^5 \cdot \sqrt{1-t^2} \cdot dt$

(6) $\int_0^1 (\cos^{-1} x)^2 \cdot dx$

(7) $\int_{-1}^1 \frac{1+x^3}{9-x^2} \cdot dx$

(8) $\int_0^{\pi} x \cdot \sin x \cdot \cos^4 x \cdot dx$

(9) $\int_0^{\pi} \frac{x}{1 + \sin^2 x} \cdot dx$

(10) $\int_1^{\infty} \frac{1}{\sqrt{x}(1+x)} \cdot dx$

(III) Evaluate :

(1) $\int_0^1 \left(\frac{1}{1+x^2} \right) \sin^{-1} \left(\frac{2x}{1+x^2} \right) \cdot dx$

(2) $\int_0^{\pi/2} \frac{1}{6 - \cos x} \cdot dx$

(3) $\int_0^a \frac{1}{a^2 + ax - x^2} \cdot dx$

(4) $\int_{\pi/5}^{3\pi/10} \frac{\sin x}{\sin x + \cos x} \cdot dx$

(5) $\int_0^1 \sin^{-1} \left(\frac{2x}{1+x^2} \right) \cdot dx$

(6) $\int_0^{\pi/4} \frac{\cos 2x}{1 + \cos 2x + \sin 2x} \cdot dx$

(7) $\int_0^{\pi/2} (2 \cdot \log \sin x - \log \sin 2x) \cdot dx$

(8) $\int_0^{\pi} (\sin^{-1} x + \cos^{-1} x)^3 \cdot \sin^3 x \cdot dx$

(9) $\int_0^4 [\sqrt{x^2 + 2x + 3}]^{-1} \cdot dx$

(10) $\int_{-2}^3 |x-2| \cdot dx$

(IV) Evaluate the following :

(1) If $\int_0^a \sqrt{x} \cdot dx = 2a \cdot \int_0^{\frac{\pi}{2}} \sin^3 x \cdot dx$ then find the value of $\int_a^{a+1} x \cdot dx$.

(2) If $\int_0^k \frac{1}{2+8x^2} \cdot dx = \frac{\pi}{16}$. Find k .

(3) If $f(x) = a + bx + cx^2$, show that $\int_0^1 f(x) = \frac{1}{6} \left[f(0) + 4f\left(\frac{1}{2}\right) + f(1) \right]$.



5. APPLICATION OF DEFINITE INTEGRATION

Let us Study

- Area under the curve
 - Area bounded by the curve, axis and given lines
 - Area between two curves.

Let us Recall

- In previous chapter, we have studied definition of definite integral as limit of a sum. Geometrically $\int_a^b f(x) \cdot dx$ gives the area A under the curve $y = f(x)$ with $f(x) > 0$ and bounded by the X-axis and the lines $x = a$, $x = b$; and is given by

$$\int_a^b f(x) dx = \phi(b) - \phi(a)$$

where $\int f(x) dx = \phi(x)$

This is also known as fundamental theorem of integral calculus.

We shall find the area under the curve by using definite integral.

5.1 Area under the curve :

For evaluation of area bounded by certain curves, we need to know the nature of the curves and their graphs. We should also be able to draw sketch of the curves.

5.1.1 Area under a curve :

The curve $y = f(x)$ is continuous in $[a, b]$ and $f(x) \geq 0$ in $[a, b]$.

1. The area shaded in figure 5.2 is bounded by the curve $y = f(x)$, X-axis and the lines $x = a$, $x = b$ and is given by the definite integral $\int_{x=a}^{x=b} (y) \cdot dx$

A = area of the shaded region.

$$A = \int_a^b f(x) \cdot dx$$




Fig. 5.1




Fig. 5.2




Fig. 5.3

2. The area A, bounded by the curve $x = g(y)$, Y axis and the lines $y = c$ and $y = d$ is given by

$$A = \int_{y=c}^d x \cdot dy$$

$$= \int_{y=c}^d g(y) \cdot dx$$

SOLVED EXAMPLE




Fig. 5.4

- Ex. 1 :** Find the area bounded by the curve $y = x^2$, the Y axis the X axis and $x = 3$.

Solution : The required area $A = \int_{x=0}^3 y \cdot dx$

$$A = \int_0^3 x^2 \cdot dx$$

$$= \left[\frac{x^3}{3} \right]_0^3$$

$$A = 9 - 0$$

$$= 9 \text{ sq.units}$$

5.1.2 Area between two curves :




Fig. 5.5

Let $y = f(x)$ and $y = g(x)$ be the equations of the two curves as shown in fig 5.5.

Let A be the area bounded by the curves $y = f(x)$ and $y = g(x)$

$$A = | A_1 - A_2 | \quad \text{where}$$

A_1 = Area bounded by the curve $y = f(x)$, X-axis and $x = a, x = b$.

A_2 = Area bounded by the curve $y = g(x)$, X-axis and $x = a, x = b$.

The point of intersection of the curves $y = f(x)$ and $y = g(x)$ can be obtained by solving their equations simultaneously.

∴ The required area $A = \left| \int_a^b f(x) dx - \int_a^b g(x) dx \right|$

SOLVED EXAMPLES

Ex. 1 : Find the area of the region bounded by the curves $y^2 = 9x$ and $x^2 = 9y$.

Solution : The equations of the curves are

$$y^2 = 9x \dots\dots (I)$$

and $x^2 = 9y \dots\dots (II)$

Squaring equation (II)

$$x^4 = 81y^2$$

$$x^4 = 81(9x) \dots \text{by (I)}$$

$$x^4 = 729x$$

$$\therefore x(x^3 - 9^3) = 0$$

i.e. $x(x^3 - 9^3) = 0$

$$\Rightarrow x = 0 \quad \text{or} \quad x = 9$$

From equation (II), $y = 0$ or $y = 9$

∴ The points of intersection of the curves are (0, 0), (9, 9).

$$\begin{aligned} \therefore \text{Required area } A &= \int_0^9 \sqrt{9x} dx - \int_0^9 \frac{x^2}{9} dx \\ &= \left[3 \cdot \frac{2}{3} \cdot x^{\frac{3}{2}} \right]_0^9 - \left[\frac{1}{9} \cdot \frac{x^3}{3} \right]_0^9 \\ &= 2 \cdot 9^{\frac{3}{2}} - 27 \\ A &= 54 - 27 \\ &= 27 \text{ sq.units} \end{aligned}$$





Fig. 5.6

Now, we will see how to find the area bounded by the curve $y = f(x)$, X-axis and lines $x = a, x = b$ if $f(x)$ is negative i.e. $f(x) \leq 0$ in $[a, b]$.



Ex. 2 : Find the area bounded by the curve $y = -x^2$, X-axis and lines $x = 1$ and $x = 4$.

Solution : Let A be the area bounded by the curve $y = -x^2$, X-axis and $1 \leq x \leq 4$.




Fig. 5.7

$$\begin{aligned} \text{The required area } A &= \int_1^4 y \, dx \\ &= \int_1^4 -x^2 \, dx \\ &= \left[-\frac{x^3}{3} \right]_1^4 \\ &= -\frac{64}{3} + \frac{1}{3} \\ A &= -21, \end{aligned}$$

But we consider the area to be positive.

$$\therefore A = \left| -21 \right| \text{sq.units} = 21 \text{ square units.}$$

Thus, if $f(x) \leq 0$ or $f(x) \geq 0$ in $[a, b]$ then the area enclosed between $y = f(x)$, X-axis and $x = a, x = b$ is $\left| \int_a^b f(x) \cdot dx \right|$.




Fig. 5.8

If the area A is divided into two parts A_1 and A_2 such that

A_1 is the part of $a \leq x \leq t$ where $f(x) \leq 0$ and

A_2 is the part of $a \leq x \leq t$ where $f(x) \geq 0$

then in A_1 , the required area is below the X-axis

and in A_2 , the required area is above the X-axis.

Now the total area $A = A_1 + A_2$

$$= \left| \int_a^t f(x) \, dx \right| + \left| \int_t^b f(x) \, dx \right|$$

Ex. 3 : Find the area bounded by the line $y = x$, X axis and the lines $x = -1$ and $x = 4$.

Solution : Consider the area A, bounded by straight line $y = x$, X axis and $x = -1, x = 4$.





Fig. 5.9

From figure 5.9, A is divided into A_1 and A_2

$$\begin{aligned} \text{The required area } A_1 &= \int_{-1}^0 y \, dx = \int_{-1}^0 x \, dx \\ &= \left[\frac{x^2}{2} \right]_{-1}^0 \\ &= 0 - \frac{1}{2} \\ A_1 &= -\frac{1}{2} \text{ square units.} \end{aligned}$$



But area is always positive.


$$\therefore A_1 = \left| -\frac{1}{2} \right| \text{sq.units} = \frac{1}{2} \text{ square units.}$$

$$A_2 = \int_0^4 y \, dx = \int_0^4 x \, dx = \left[\frac{x^2}{2} \right]_0^4 = \frac{4^2}{2} = 8 \text{ square units.}$$

$$\therefore \text{Required area } A = A_1 + A_2 = \frac{1}{2} + 8 = \frac{17}{2} \text{ sq.units}$$

Ex. 4 : Find the area enclosed between the X-axis and the curve $y = \sin x$ for values of x between 0 to 2π .

Solution : The area enclosed between the curve and the X-axis consists of equal area lying alternatively above and below X-axis which are respectively positive and negative.



1) Area $A_1 =$ area lying above the X-axis

$$= \int_0^{\pi} \sin x \cdot dx = [-\cos x]_0^{\pi}$$

$$= -[\cos \pi - \cos 0] = -(-1 - 1)$$

$$A_1 = 2$$

2) Area $A_2 =$ area lying below the X-axis $= \int_{\pi}^{2\pi} \sin x \, dx = [-\cos x]_{\pi}^{2\pi} = [-\cos 2\pi - \cos \pi]$

$$= -[-1 - (-1)]$$

$$A_2 = -2$$

$$\therefore \text{Total area} = A_1 + |A_2| = 2 + |(-2)| = 4 \text{ sq.units.}$$

Activity :

Ex. 5 : Find the area enclosed between $y = \sin x$ and X-axis between 0 and 4π .

Ex. 6 : Find the area enclosed between $y = \cos x$ and X-axis between the limits :

(i) $0 \leq x \leq \frac{\pi}{2}$

(ii) $\frac{\pi}{2} \leq x \leq \pi$

(iii) $0 \leq x \leq \pi$





SOLVED EXAMPLES

Ex. 1 : Using integration, find the area of the region bounded by the line $2y + x = 8$, X-axis and the lines $x = 2$ and $x = 4$.

Solution : The required region is bounded by the lines $2y + x = 8$, and $x = 2, x = 4$ and X-axis.




Fig. 5.11

$$\therefore y = \frac{1}{2}(8 - x) \text{ and the limits are } x = 2, x = 4.$$

$$\begin{aligned} \text{Required area} &= \text{Area of the shaded region} \\ &= \int_{x=2}^4 y \, dx \\ &= \int_2^4 \frac{1}{2}(8 - x) \, dx \\ &= \frac{1}{2} \left[8x - \frac{x^2}{2} \right]_2^4 \\ &= \frac{1}{2} \left[\left(8 \cdot 4 - \frac{4^2}{2} \right) - \left(8 \cdot 2 - \frac{2^2}{2} \right) \right] \\ &= 5 \text{ sq. units.} \end{aligned}$$

Ex. 2 : Find the area of the regions bounded by the following curve, the X-axis and the given lines :

(i) $y = x^2, x = 1, x = 2$

(ii) $y^2 = 4x, x = 1, x = 4, y \geq 0$

(iii) $y = \sin x, x = -\frac{\pi}{2}, x = \frac{\pi}{2}$

Solution : Let A be the required area




Fig. 5.12

$$\begin{aligned} \text{(i) } A &= \int_1^3 y \, dx \\ &= \int_1^3 x^2 \, dx \\ &= \frac{1}{3} \left[x^3 \right]_1^3 \\ &= \frac{1}{3} [27 - 1] \\ A &= \frac{26}{3} \text{ sq. units.} \end{aligned}$$



$$\begin{aligned}
 \text{(ii) } A &= \int_1^4 y \, dx \\
 &= \int_1^4 2\sqrt{x} \, dx \\
 &= 2 \int_1^4 x^{\frac{1}{2}} \, dx \\
 &= 2 \cdot \frac{2}{3} \left[x^{\frac{3}{2}} \right]_1^4 = \frac{4}{3} \left[4^{\frac{3}{2}} - 1 \right]
 \end{aligned}$$

$$A = \frac{28}{3} \text{ sq. units.}$$




Fig. 5.13

$$\begin{aligned}
 \text{(iii) } A &= \int_{-\pi/2}^{\pi/2} y \, dx \\
 &= \int_{-\pi/2}^{\pi/2} \sin x \, dx \\
 &= \left| \int_{-\pi/2}^0 \sin x \cdot dx \right| + \left| \int_0^{\pi/2} \sin x \cdot dx \right| \\
 &= \left| [-\cos x]_{-\pi/2}^0 \right| + \left| [-\cos x]_0^{\pi/2} \right| \\
 &= \left| -[\cos 0 - \cos(\frac{\pi}{2})] \right| + \left| -[\cos(\frac{\pi}{2}) + \cos 0] \right| \\
 &= |[-1 - 0] + [0 + 1]| = 1 + 1
 \end{aligned}$$

$$A = 2 \text{ sq. units.}$$




Fig. 5.14

Ex. 3 : Find the area of the region bounded by the parabola $y^2 = 16x$ and the line $x = 4$.

Solution : $y^2 = 16x \Rightarrow y = \pm 4\sqrt{x}$

$$\begin{aligned}
 A &= \text{Area POCP} + \text{Area QOCQ} \\
 &= 2 (\text{Area POCP}) \\
 &= 2 \int_0^4 y \cdot dx \\
 &= 2 \int_0^4 4\sqrt{x} \cdot dx \\
 A &= 8 \cdot \frac{2}{3} \left[x^{\frac{3}{2}} \right]_0^4 = \frac{16}{3} \times 8
 \end{aligned}$$

$$A = \frac{128}{3} \text{ sq. units.}$$




Fig. 5.15

Ex. 4 : Find the area of the region bounded by the curves $x^2 = 16y$, $y = 1$, $y = 4$ and the Y-axis, lying in the first quadrant.

Solution :




Fig. 5.16

$$\begin{aligned} \text{Required area} &= \int_1^4 x \, dy \\ A &= \int_1^4 \sqrt{16y} \, dy \\ &= 4 \int_1^4 \sqrt{y} \cdot dy \\ &= 4 \cdot \left[\frac{2}{3} \cdot y^{\frac{3}{2}} \right]_1^4 \\ &= \frac{8}{3} \times [8 - 1] \\ A &= \frac{56}{3} \text{ sq. units.} \end{aligned}$$

Ex. 5 : Find the area of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.

Solution : By the symmetry of the ellipse, required area of the ellipse is 4 times the area of the region OPQO. For this region the limit of integration are $x = 0$ and $x = a$.




Fig. 5.17

From the equation of ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y^2 = b^2 \cdot \left(\frac{a^2 - x^2}{a^2} \right)$$

$$y = \frac{b}{a} \cdot \sqrt{a^2 - x^2} \quad , \text{ In first quadrant, } y > 0$$

$$\begin{aligned} A &= 4 \int_{x=0}^a y \, dx \\ &= \int_0^a \frac{b}{a} \cdot \sqrt{a^2 - x^2} \, dx \\ &= \frac{4b}{a} \cdot \left[\frac{x}{a} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right]_0^a \\ &= \frac{4b}{a} \cdot \left[\frac{a^2}{2} \cdot \frac{\pi}{2} - 0 \right]_0^a \\ A &= \pi ab \text{ sq. units} \end{aligned}$$



Ex. 6 : Find the area of the region lying between the parabolas $y^2 = 4ax$ and $x^2 = 4ay$ where $a > 0$.

Solution : The equations of the parabolas are

$$y^2 = 4ax \quad \dots \text{(I)}$$

and $x^2 = 4ay \quad \dots \text{(II)}$

From (ii) $y = \frac{x^2}{4a}$ substitute in (I)

$$\left(\frac{x^2}{4a}\right)^2 = 4ax$$

$$\Rightarrow x^4 = 64a^3x$$

$$\therefore x(x^3 - 64a^3) = 0$$

$$\therefore x[x^3 - (4a^3)] = 0$$

$$\therefore x = 0 \text{ and } x = 4a \quad \therefore y = 0 \text{ and } y = 4a$$

The point of intersection of curves are O (0, 0), P (4a, 4a)

\therefore The required area is in the first quadrant and it is

A = area under the parabola ($y^2 = 4ax$) – area under the parabola ($x^2 = 4ay$)

$$\begin{aligned} A &= \int_0^{4a} \sqrt{4ax} \, dx - \int_0^{4a} \frac{x^2}{4a} \, dx &&= \sqrt{4a} \int_0^{4a} x^{\frac{1}{2}} \, dx - \int_0^{4a} \frac{x^2}{4a} \, dx \\ &= 2\sqrt{a} \cdot \left[\frac{2}{3} \cdot x^{\frac{3}{2}} \right]_0^{4a} - \frac{1}{4a} \cdot \left[\frac{x^3}{3} \right]_0^{4a} \\ &= \frac{4}{3} \sqrt{a} \cdot \left[4a \sqrt{4a} - \frac{1}{4a} \cdot 64a^3 \right] = \frac{32}{3} a^2 - \frac{16}{3} a^2 &&\therefore A = \frac{16}{3} a^2 \text{ sq. units.} \end{aligned}$$




Fig. 5.18

Ex. 7 : Find the area of the region bounded by the curve $y = x^2$ and the line $y = 4$.

Solution : Required area $A = 2 \times$ area of OPQO

$$\begin{aligned} \therefore A &= \int_0^4 x \cdot dy \\ A &= 2 \cdot \int_0^4 \sqrt{y} \cdot dy \\ &= 2 \cdot \left[\frac{2}{3} \cdot y^{\frac{3}{2}} \right]_0^4 = \left(\frac{4}{3} \times 4^{\frac{3}{2}} \right) \\ &= \frac{4}{3} \times 8 \\ A &= \frac{32}{3} \text{ sq. units.} \end{aligned}$$




Fig. 5.19

Ex. 8 : Find the area of sector bounded by the circle $x^2 + y^2 = 16$ and the line $y = x$ in the first quadrant.

Solution : Required area $A = A(\Delta OCB) + A(\text{region } ABC)$




Fig. 5.20

To find,

The point of intersection of $x^2 + y^2 = 16 \dots (I)$

and line $y = x \dots (II)$

Substitute (II) in (I)

$$x^2 + x^2 = 16$$

$$2x^2 = 16$$

$$x^2 = 8$$

$$x = \pm 2\sqrt{2}, \quad y = \pm 2\sqrt{2}$$

The point of intersection is $B(2\sqrt{2}, 2\sqrt{2})$

$$\begin{aligned} A &= \int_0^{2\sqrt{2}} x \cdot dx + \int_{2\sqrt{2}}^4 \sqrt{16-x^2} \cdot dx = \frac{1}{2} \left[x^2 \right]_0^{2\sqrt{2}} + \left[\frac{x}{2} \sqrt{16-x^2} + \frac{16}{2} \sin^{-1} \frac{x}{4} \right]_{2\sqrt{2}}^4 \\ &= \frac{1}{2} \cdot (2\sqrt{2})^2 + \left[8 \sin^{-1} 1 - \left(\frac{2\sqrt{2}}{2} \sqrt{8} + 8 \sin^{-1} \frac{1}{2} \right) \right] \\ &= 4 + 8 \cdot \frac{\pi}{2} - 4 - 8 \cdot \frac{\pi}{4} \quad \therefore A = 2\pi \text{ sq. units.} \end{aligned}$$

Note that, the required area is $\frac{1}{8}$ times the area of the circle given.

EXERCISE 5.1

- (1) Find the area of the region bounded by the following curves, X-axis and the given lines:
 - (i) $y = 2x, x = 0, x = 5$
 - (ii) $x = 2y, y = 0, y = 4$
 - (iii) $x = 0, x = 5, y = 0, y = 4$
 - (iv) $y = \sin x, x = 0, x = \frac{\pi}{2}$
 - (v) $xy = 2, x = 1, x = 4$
 - (vi) $y^2 = x, x = 0, x = 4$
 - (vii) $y^2 = 16x$ and $x = 0, x = 4$
- (2) Find the area of the region bounded by the parabola :
 - (i) $y^2 = 16x$ and its latus rectum.
 - (ii) $y = 4 - x^2$ and the X-axis
- (3) Find the area of the region included between:
 - (i) $y^2 = 2x$, line $y = 2x$
 - (ii) $y^2 = 4x$, line $y = x$
 - (iii) $y = x^2$ and the line $y = 4x$
 - (iv) $y^2 = 4ax$ and the line $y = x$
 - (v) $y = x^2 + 3$ and the line $y = x + 3$



Let us Remember

- ✿ The area A , bounded by the curve $y = f(x)$, X-axis and the lines $x = a$ and $x = b$ is given by

$$A = \int_a^b f(x) \cdot dx = \int_{x=a}^{x=b} f(x) \cdot dx$$

If the area A lies below the X-axis, then A is negative and in this case we take $|A|$.

- ✿ The area A of the region bounded by the curve $x = g(y)$, the Y axis, and the lines $y = c$ and $y = d$ is given by

$$A = \int_{y=c}^d f(x) \cdot dx = \int_{y=c}^{y=d} g(y) \cdot dx$$

- ✿ **Tracing of curve :**

- (i) X-axis is an axis of symmetry for a curve C , if $(x, y) \in C \Leftrightarrow (x, -y) \in C$.
- (ii) Y-axis is an axis of symmetry for a curve C , if $(x, y) \in C \Leftrightarrow (-x, y) \in C$.
- (iii) If replacing x and y by $-x$ and $-y$ respectively, the equation of the curve is unchanged then the curve is symmetric about X-axis and Y-axis.

MISCELLANEOUS EXERCISE 5

(I) Choose the correct option from the given alternatives :

- (1) The area bounded by the region $1 \leq x \leq 5$ and $2 \leq y \leq 5$ is given by
 (A) 12 sq. units (B) 8 sq. units (C) 25 sq. units (D) 32 sq. units
- (2) The area of the region enclosed by the curve $y = \frac{1}{x}$, and the lines $x = e$, $x = e^2$ is given by
 (A) 1 sq. unit (B) $\frac{1}{2}$ sq. unit (C) $\frac{3}{2}$ sq. units (D) $\frac{5}{2}$ sq. units
- (3) The area bounded by the curve $y = x^3$, the X-axis and the lines $x = -2$ and $x = 1$ is
 (A) -9 sq. units (B) $-\frac{15}{4}$ sq. units (C) $\frac{15}{4}$ sq. units (D) $\frac{17}{4}$ sq. units
- (4) The area enclosed between the parabola $y^2 = 4x$ and line $y = 2x$ is
 (A) $\frac{2}{3}$ sq. units (B) $\frac{1}{3}$ sq. units (C) $\frac{1}{4}$ sq. units (D) $\frac{3}{4}$ sq. units
- (5) The area of the region bounded between the line $x = 4$ and the parabola $y^2 = 16x$ is
 (A) $\frac{128}{3}$ sq. units (B) $\frac{108}{3}$ sq. units (C) $\frac{118}{3}$ sq. units (D) $\frac{218}{3}$ sq. units

- (6) The area of the region bounded by $y = \cos x$, Y-axis and the lines $x = 0$, $x = 2\pi$ is
 (A) 1 sq. unit (B) 2 sq. units (C) 3 sq. units (D) 4 sq. units
- (7) The area bounded by the parabola $y^2 = 8x$ the X-axis and the latus rectum is
 (A) $\frac{31}{3}$ sq. units (B) $\frac{32}{3}$ sq. units (C) $\frac{32\sqrt{2}}{3}$ sq. units (D) $\frac{16}{3}$ sq. units
- (8) The area under the curve $y = 2\sqrt{x}$, enclosed between the lines $x = 0$ and $x = 1$ is
 (A) 4 sq. units (B) $\frac{3}{4}$ sq. units (C) $\frac{2}{3}$ sq. units (D) $\frac{4}{3}$ sq. units
- (9) The area of the circle $x^2 + y^2 = 25$ in first quadrant is
 (A) $\frac{25\pi}{3}$ sq. units (B) 5π sq. units (C) 5 sq. units (D) 3 sq. units
- (10) The area of the region bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is
 (A) ab sq. units (B) πab sq. units (C) $\frac{\pi}{ab}$ sq. units (D) πa^2 sq. units
- (11) The area bounded by the parabola $y^2 = x$ and the line $2y = x$ is
 (A) $\frac{4}{3}$ sq. units (B) 1 sq. units (C) $\frac{2}{3}$ sq. units (D) $\frac{1}{3}$ sq. units
- (12) The area enclosed between the curve $y = \cos 3x$, $0 \leq x \leq \frac{\pi}{6}$ and the X-axis is
 (A) $\frac{1}{2}$ sq. units (B) 1 sq. units (C) $\frac{2}{3}$ sq. units (D) $\frac{1}{3}$ sq. units
- (13) The area bounded by $y = \sqrt{x}$ and line $x = 2y + 3$, X-axis in first quadrant is
 (A) $2\sqrt{3}$ sq. units (B) 9 sq. units (C) $\frac{34}{3}$ sq. units (D) 18 sq. units
- (14) The area bounded by the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the line $\frac{x}{a} + \frac{y}{b} = 1$ is
 (A) $\pi ab - 2 ab$ (B) $\frac{\pi ab}{4} - \frac{ab}{2}$ (C) $\pi ab - ab$ (D) πab
- (15) The area bounded by the parabola $y = x^2$ and the line $y = x$ is
 (A) $\frac{1}{2}$ (B) $\frac{1}{3}$ (C) $\frac{1}{6}$ (D) $\frac{1}{12}$
- (16) The area enclosed between the two parabolas $y^2 = 4x$ and $y = x$ is
 (A) $\frac{8}{3}$ (B) $\frac{32}{3}$ (C) $\frac{16}{3}$ (D) $\frac{4}{3}$



- (17) The area bounded by the curve $y = \tan x$, X-axis and the line $x = \frac{\pi}{4}$ is
 (A) $\frac{1}{3} \log 2$ (B) $\log 2$ (C) $2 \log 2$ (D) $3 \cdot \log 2$
- (18) The area of the region bounded by $x^2 = 16y$, $y = 1$, $y = 4$ and $x = 0$ in the first quadrant, is
 (A) $\frac{7}{3}$ (B) $\frac{8}{3}$ (C) $\frac{64}{3}$ (D) $\frac{56}{3}$
- (19) The area of the region included between the parabolas $y^2 = 4ax$ and $x^2 = 4ay$, ($a > 0$) is given by
 (A) $\frac{16 a^2}{3}$ (B) $\frac{8 a^2}{3}$ (C) $\frac{4 a^2}{3}$ (D) $\frac{32 a^2}{3}$
- (20) The area of the region included between the line $x + y = 1$ and the circle $x^2 + y^2 = 1$ is
 (A) $\frac{\pi}{2} - 1$ (B) $\pi - 2$ (C) $\frac{\pi}{4} - \frac{1}{2}$ (D) $\pi - \frac{1}{2}$

(II) Solve the following :

- (1) Find the area of the region bounded by the following curve, the X-axis and the given lines
 (i) $0 \leq x \leq 5, 0 \leq y \leq 2$ (ii) $y = \sin x, x = 0, x = \pi$ (iii) $y = \sin x, x = 0, x = \frac{\pi}{3}$
- (2) Find the area of the circle $x^2 + y^2 = 9$, using integration.
- (3) Find the area of the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ using integration.
- (4) Find the area of the region lying between the parabolas.
 (i) $y^2 = 4x$ and $x^2 = 4y$ (ii) $4y^2 = 9x$ and $3x^2 = 16y$ (iii) $y^2 = x$ and $x^2 = y$
- (5) Find the area of the region in first quadrant bounded by the circle $x^2 + y^2 = 4$ and the x axis and the line $x = y\sqrt{3}$.
- (6) Find the area of the region bounded by the parabola $y^2 = x$ and the line $y = x$ in the first quadrant.
- (7) Find the area enclosed between the circle $x^2 + y^2 = 1$ and the line $x + y = 1$, lying in the first quadrant.
- (8) Find the area of the region bounded by the curve $(y - 1)^2 = 4(x + 1)$ and the line $y = (x - 1)$.
- (9) Find the area of the region bounded by the straight line $2y = 5x + 7$, X-axis and $x = 2, x = 5$.
- (10) Find the area of the region bounded by the curve $y = 4x^2$, Y-axis and the lines $y = 1, y = 4$.



6. DIFFERENTIAL EQUATIONS



Let us Study

- Differential Equation
- Formation of differential equation
- Types of differential equation.
- Order and degree of differential equation
- Solution of differential equation
- Application of differential equation.



Let us Recall

- The differentiation and integration of functions and the properties of differentiation and integration.



Let us Learn

6.1.1 Introduction :

In physics, chemistry and other sciences we often have to build mathematical models which involves differential equations. We need to find functions which satisfy those differential equations.

6.1.2 Differential Equation :

Equation which contains the derivative of a function is called a **differential equation**. The following are differential equations.

$$(i) \quad \frac{dy}{dx} = \cos x$$

$$(ii) \quad \frac{d^2y}{dx^2} + k^2y = 0$$

$$(iii) \quad \left(\frac{d^2w}{dx^2} \right) - x^2 \frac{dw}{dx} + w = 0$$

$$(iv) \quad \frac{d^2y}{dt^2} + \frac{d^2x}{dt^2} = x, \text{ here } x \text{ and } y \text{ are functions of 't' .}$$

$$(v) \quad \frac{d^3y}{dx^3} + x \frac{dy}{dx} - 4xy = 0, \text{ here } x \text{ is a function of } y.$$

$$(vi) \quad r \frac{dr}{d\theta} + \cos \theta = 5$$

6.2 Order and degree of the differential equation :

The order of a differential equation is the highest order of the derivative appearing in the equation.

The degree of differential equation is the power of the highest ordered derivative present in the equation. To find the degree of the differential equation, we need to have a positive integer as the index of each derivative.



SOLVED EXAMPLES

Ex. 1 : Find order and degree of the following differential equations.

(i) $x^2 \frac{d^2y}{dx^2} + 3x \frac{dy}{dx} + 4y = 0$

Solution : It's order is 2 and degree is 1.

(ii) $\left(\frac{d^3y}{dx^3}\right)^2 + xy \frac{dy}{dx} - 2x + 3y + 7 = 0$

Solution : It's order is 3 and degree is 2.

(iii) $r \frac{dr}{d\theta} + \cos \theta = 5$

Solution : It's order is 1 and degree is 1.

(iv) $\left(\frac{d^2y}{dx^2}\right)^2 + \left(\frac{dy}{dx}\right)^2 = e^x$

Solution : It's order is 2 and degree is 2.

(v) $\frac{dy}{dx} + \frac{3xy}{\frac{dy}{dx}} = \cos x$

Solution : This equation expressed as

$$\left(\frac{dy}{dx}\right)^2 + 3xy = \cos x \left(\frac{dy}{dx}\right)$$

It's order is 1 and degree is 2.

(vi) $\sqrt{1 + \frac{1}{\left(\frac{dy}{dx}\right)^2}} = \left(\frac{d^2y}{dx^2}\right)^{\frac{3}{2}}$

Solution : This equation can be expressed as

$$1 + \frac{1}{\left(\frac{dy}{dx}\right)^2} = \left(\frac{d^2y}{dx^2}\right)^3$$

$$\therefore \left(\frac{dy}{dx}\right)^2 + 1 = \left(\frac{d^2y}{dx^2}\right)^3 \left(\frac{dy}{dx}\right)^2$$

It's order is 2 and degree is 3.

(vii) $\frac{d^4y}{dx^4} = \left[1 + \left(\frac{dy}{dx}\right)^2\right]^3$

Solution : It's order is 4 and degree is 1.

(viii) $e^{\frac{dy}{dx}} + \frac{dy}{dx} = x$

Solution : It's order is 1, but equation can not be expressed as a polynomial differential equation.

\therefore The degree is not defined.

(ix)
$$\begin{vmatrix} x^3 & y^3 & 3 \\ 2x^2 & 3y \frac{dy}{dx} & 0 \\ 5x & 2 \left[y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 \right] & 0 \end{vmatrix} = 0$$

Solution : $\therefore x^3 [0 - 0] - y^2 [0 - 0] + 3 \left\{ 4x^2 \left[y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 \right] - 15xy \frac{dy}{dx} \right\} = 0$

$\therefore 4x^2 y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 - 15xy \frac{dy}{dx} = 0 \quad \therefore$ Its order is 2 and degree is 1.

Notes : (1) $\frac{dy}{dx}$ is also denoted by y' , $\frac{d^2y}{dx^2}$ is also denoted by y'' , $\frac{d^3y}{dx^3}$ is also by y''' and so-on.

(2) The order and degree of a differential equation are always positive integers.



EXERCISE 6.1

(1) Determine the order and degree of each of the following differential equations.

(i) $\frac{d^2y}{dx^2} + x \left(\frac{dy}{dx}\right) + y = 2 \sin x$

(ii) $\sqrt[3]{1 + \left(\frac{dy}{dx}\right)^2} = \frac{d^2y}{dx^2}$

(iii) $\frac{dy}{dx} = \frac{2 \sin x + 3}{\frac{dy}{dx}}$

(iv) $\frac{d^2y}{dx^2} + \frac{dy}{dx} + x = \sqrt{1 + \frac{d^3y}{dx^3}}$

(v) $\frac{d^2y}{dt^2} + \left(\frac{dy}{dt}\right)^2 + 7x + 5 = 0$

(vi) $(y''')^2 + 3y'' + 3xy' + 5y = 0$

(vii) $\left(\frac{d^2y}{dx^2}\right)^2 + \cos\left(\frac{dy}{dx}\right) = 0$

(viii) $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}} = 8 \frac{d^2y}{dx^2}$

(ix) $\left(\frac{d^3y}{dx^3}\right)^{\frac{1}{2}} - \left(\frac{dy}{dx}\right)^{\frac{1}{3}} = 20$

(x) $x + \frac{d^2y}{dx^2} = \sqrt{1 + \left(\frac{d^2y}{dx^2}\right)^2}$

6.3 Formation of Differential Equation :

From the given information, we can form the differential equation. Sometimes we need to eliminate the arbitrary constants from a given relation. It may be done by differentiation.



SOLVED EXAMPLES

Ex. 1 : Obtain the differential equation by eliminating the arbitrary constants from the following :

(i) $y^2 = 4ax$

(ii) $y = Ae^{3x} + Be^{-3x}$

(iii) $y = (c_1 + c_2x) e^x$

(iv) $y = c^2 + \frac{c}{x}$

(v) $y = c_1e^{3x} + c_2e^{2x}$

Solution :

(i) $y^2 = 4ax \dots (1)$

Here a is the arbitrary constant, we differentiate *w. r. t. x*,

$\therefore \frac{dy}{dx} = 4a$

then eq. (1) gives

$y^2 = \left(\frac{dy}{dx}\right)x$ is required differential equation.

(ii) $y = Ae^{3x} + Be^{-3x} \dots (1)$

Here A and B are arbitrary constants.

Differentiate *w. r. t. x*, we get

$\therefore \frac{dy}{dx} = 3Ae^{3x} - 3Be^{-3x}$

again Differentiate *w. r. t. x*, we get

$\frac{d^2y}{dx^2} = 3 \times 3Ae^{3x} - 3 \times 3Be^{-3x}$

$= 9(Ae^{3x} + Be^{-3x}) = 9y \dots$ from eq.(1)

$\therefore \frac{d^2y}{dx^2} = 9y$

(iii) $y = (c_1 + c_2 x) e^x \dots (1)$

Here c_1 and c_2 are arbitrary constants.

Differentiate *w. r. t.* x , we get

$$\therefore \frac{dy}{dx} = (c_1 + c_2 x) e^x + c_2 e^x$$

$$\therefore \frac{dy}{dx} = y + c_2 e^x \dots (2) \quad \dots \text{from eq.(1)}$$

Again differentiate *w. r. t.* x , we get

$$\frac{d^2y}{dx^2} = \frac{dy}{dx} + c_2 e^x$$

$$\therefore c_2 e^x = \frac{d^2y}{dx^2} - \frac{dy}{dx}$$

put in eq.(2)

$$\frac{dy}{dx} = y + \frac{d^2y}{dx^2} - \frac{dy}{dx}$$

$$\therefore \frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + y = 0$$

(v) $y = c_1 e^{3x} + c_2 e^{2x} \dots (I)$

Differentiate *w. r. t.* x , we get

$$\therefore \frac{dy}{dx} = 3c_1 e^{3x} + 2c_2 e^{2x} \dots (II)$$

Again differentiate *w. r. t.* x , we get

$$\frac{d^2y}{dx^2} = 9c_1 e^{3x} + 4c_2 e^{2x} \dots (III)$$

As equations (I), (II) and (III) in $c_1 e^{3x}$ and $c_2 e^{2x}$ are consistent

$$\therefore \begin{vmatrix} y & 1 & 1 \\ \frac{dy}{dx} & 3 & 2 \\ \frac{d^2y}{dx^2} & 9 & 4 \end{vmatrix} = 0$$

$$\therefore y(12 - 18) - 1 \left(4 \frac{dy}{dx} - 2 \frac{d^2y}{dx^2} \right) + 1 \left(9 \frac{dy}{dx} - 3 \frac{d^2y}{dx^2} \right) = 0$$

$$\therefore -6y - 4 \frac{dy}{dx} + 2 \frac{d^2y}{dx^2} + 9 \frac{dy}{dx} - 3 \frac{d^2y}{dx^2} = 0$$

$$\therefore -\frac{d^2y}{dx^2} + 5 \frac{dy}{dx} - 6y = 0$$

(iv) $y = c^2 + \frac{c}{x} \dots (1)$

Differentiate *w. r. t.* x , we get

$$\therefore \frac{dy}{dx} = -\frac{c}{x^2}$$

$$\therefore c = -x^2 \frac{dy}{dx}$$

then eq.(1) gives

$$y = \left[-x^2 \left(\frac{dy}{dx} \right)^2 \right] - x^2 \frac{dy}{dx} \times \frac{1}{x}$$

$$\therefore y = x^4 \left(\frac{dy}{dx} \right)^2 - x \frac{dy}{dx}$$

$$\therefore x^4 \left(\frac{dy}{dx} \right)^2 - x \frac{dy}{dx} - y = 0$$

Ex. 2 : The rate of decay of the mass of a radioactive substance any time is k times its mass at that time, form the differential equation satisfied by the mass of the substance.

Solution : Let m be the mass of a radioactive substance time ' t '

$$\therefore \quad \text{The rate of of decay of mass is } \frac{dm}{dt}$$

$$\text{Here } \frac{dm}{dt} \propto m$$

$$\therefore \quad \frac{dm}{dt} = mk, \text{ where } k < 0$$

is the required differential equation.

Ex. 3 : Form the differential equation of family of circles above the X-axis and touching the X-axis at the origin

Solution : Let $c (a, b)$ be the centre of the circle touching X-axis at the origin ($b < 0$).

The radius of the circle of b .

The equation of the circle is

$$(x - 0)^2 + (y - b)^2 = b^2$$

$$\therefore x^2 + y^2 - 2by + b^2 = b^2$$

$$\therefore x^2 + y^2 - 2by = 0 \quad \dots (I)$$

Differentiate w. r. t. x , we get

$$2x + 2y \left(\frac{dy}{dx} \right) - 2b \left(\frac{dy}{dx} \right) = 0$$

$$\therefore x + (y - b) \frac{dy}{dx} = 0$$

$$\therefore \frac{x}{\left(\frac{dy}{dx} \right)} + (y - b) = 0$$

$$\therefore b = y + \frac{x}{\left(\frac{dy}{dx} \right)} \quad \dots (II)$$

From eq. (I) and eq. (II)

$$\therefore x^2 + y^2 - 2 \left[y + \frac{x}{\left(\frac{dy}{dx} \right)} \right] y = 0$$

$$\therefore x^2 + y^2 - 2y^2 - \frac{2xy}{\left(\frac{dy}{dx} \right)} = 0$$

$$\therefore x^2 - y^2 = \frac{2xy}{\left(\frac{dy}{dx} \right)}$$

$$\therefore (x^2 - y^2) \frac{dy}{dx} = 2xy$$

is the required differential equation.




Fig. 6.1

Activity : Form the differential equation of family of circles touching Y-axis at the origin and having their centres on the X-axis.

Ex. 4 : A particle is moving along the X-axis. Its acceleration at time t is proportional to its velocity at that time. Find the differential equation of the motion of the particle.

Solution : Let s be the displacement of the particle at time ' t '.

Its velocity and acceleration are $\frac{ds}{dt}$ and $\frac{d^2s}{dt^2}$ respectively.

$$\text{Here } \frac{d^2s}{dt^2} \propto \frac{ds}{dt}$$

$$\therefore \frac{d^2s}{dt^2} = k \frac{ds}{dt}, \quad (\text{where } k \text{ is constant } \neq 0)$$

is the required differential equation.

EXERCISE 6.2

(1) Obtain the differential equations by eliminating arbitrary constants c_1 and c_2 .

(i) $x^3 + y^3 = 4ax$

(ii) $Ax^2 + By^2 = 1$

(iii) $y = A \cos(\log x) + B \sin(\log x)$

(iv) $y^2 = (x + c)^3$

(v) $y = Ae^{5x} + Be^{-5x}$

(vi) $(y - a)^2 = 4(x - b)$

(vii) $y = a + \frac{a}{x}$

(viii) $y = c_1 e^{2x} + c_2 e^{5x}$

(ix) $c_1 x^3 + c_2 y^2 = 5$

(x) $y = e^{-2x} (A \cos x + B \sin x)$

(2) Form the differential equation of family of lines having intercepts a and b on the co-ordinate axes respectively.

(3) Find the differential equation of all parabolas having length of latus rectum $4a$ and axis is parallel to the X-axis.

(4) Find the differential equation of an ellipse whose major axis is twice its minor axis.

(5) Form the differential equation of family of lines parallel to the line $2x + 3y + 4 = 0$

(6) Find the differential equations of all circles having radius 9 and centre at point $A(h, k)$.

(7) Form the differential equation of all parabolas whose axis is the X-axis.

6.4 Solution of differential equation :

Verify that

$$y = a \sin x \text{ and } y = b \cos x$$

are solutions of the differential equation, where a and b are any constants.

Also $y = a \sin x + b \cos x$ is a solution of the equation.

Here $\sin x$ and $\cos x$ are particular solutions where as $a \sin x + b \cos x$ is the general solution which describes all possible solutions.

A solution which can be obtained from the general solution by giving particular values to the arbitrary constants is called a **particular solution**.

Therefore the differential equation has infinitely many solutions.



SOLVED EXAMPLES

Ex. 1 : Verify that : $y \sec x = \tan x + c$

is a solution of the differential equation

$$\frac{dy}{dx} + y \tan x = \sec x.$$

Solution : Here $y \sec x = \tan x + c$

Differentiate w. r. t. x , we get

$$y \sec x \tan x + \sec x \frac{dy}{dx} = \sec^2 x$$

$$\therefore \frac{dy}{dx} + y \tan x = \sec x$$

Hence $y \sec x = \tan x + c$

is a solution of the differential equation

$$\frac{dy}{dx} + y \tan x = \sec x$$

Ex. 2 : Verify that : $y = \log x + c$

is a solution of the differential equation

$$x \frac{d^2y}{dx^2} + \frac{dy}{dx} = 0.$$

Solution : Here $y = \log x + c$

Differentiate w. r. t. x , we get

$$\frac{dy}{dx} = \frac{1}{x}$$

$$\therefore x \frac{dy}{dx} = 1$$

Differentiate w. r. t. x , we get

$$x \frac{d^2y}{dx^2} + \frac{dy}{dx} \times 1 = 0$$

$$\therefore x \frac{d^2y}{dx^2} + \frac{dy}{dx} = 0$$

$y = \log x + c$ is the solution of

$$x \frac{d^2y}{dx^2} + \frac{dy}{dx} = 0.$$

Consider the example :

$$\frac{dy}{dx} = x^2y + y$$

$$\therefore \frac{1}{y} \cdot \frac{dy}{dx} = x^2 + 1$$

We can consider x and y both as variables and write this as

$$\frac{dy}{y} = (x^2 + 1) \cdot dx$$

Now we can integrate L.H.S. w. r. t. y and R.H.S. w. r. t. x , then we get

$$\therefore \log y = \frac{x^3}{3} + x + c$$

This integration is obtained by separating the variables.


It helps to examine the equation and find out if such a separation is possible.

The above method is known as the method of separation of variables.

In general, if the given differential equation can be written as

$$f(x) dx = g(y) dy$$

then this method is applicable.



SOLVED EXAMPLES

Ex. 1 : Find the general solution of the following differential equations :

(i) $\frac{dy}{dx} = x \sqrt{25 - x^2}$

(ii) $\frac{dx}{dt} = \frac{x \log x}{t}$

Solution :

(i) $\frac{dy}{dx} = x \sqrt{25 - x^2}$

$\therefore dy = x \sqrt{25 - x^2} \cdot dx$

Integrating both sides, we get

$\int dy = \int \sqrt{25 - x^2} \cdot x \cdot dx \quad \dots (I)$

Put $25 - x^2 = t$

$\therefore -2x \cdot dx = dt$

$\therefore x \cdot dx = -\frac{dt}{2}$

Eq. (I) becomes, $\int dy = \int \sqrt{t} \left(-\frac{dt}{2} \right)$

$\therefore 2 \int dy = -\int \sqrt{t} \cdot dt$

$\therefore 2 \int dy + \int t^{\frac{1}{2}} \cdot dt = 0$

$\therefore 2y + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} = c_1$

$\therefore 2y + \frac{2}{3} t^{\frac{3}{2}} = c_1$

$\therefore 6y + 2t^{\frac{3}{2}} = 3c_1$

$\therefore 6y + 2(25 - x^2)^{\frac{3}{2}} = c \quad \dots [c = 3c_1]$

(ii) $\frac{dx}{dt} = \frac{x \log x}{t}$

$\therefore \frac{dx}{x \log x} = \frac{dt}{t}$

Integrating both sides, we get

$\int \frac{dx}{x \log x} = \int \frac{dt}{t}$

$\therefore \log(\log x) = \log(t) + \log c$

$\therefore \log(\log x) = \log(tc)$

$\therefore \log x = ct$

$\therefore e^{ct} = x$

Ex. 2 : Find the particular solution with given initial conditions :

(i) $\frac{dy}{dx} = e^{2y} \cos x$ when $x = \frac{\pi}{6}, y = 0$

(ii) $\frac{y-1}{y+1} + \frac{x-1}{x+1} \cdot \frac{dy}{dx} = 0$, when $x = y = 2$

Solution :

(i) $\frac{dy}{dx} = e^{2y} \cos x$

$\therefore \frac{dy}{e^{2y}} = \cos x \cdot dx$

$\therefore e^{-2y} \cdot dy = \cos x \cdot dx$

Integrating both sides, we get

$\therefore \int e^{-2y} \cdot dy = \int \cos x \cdot dx$

$\therefore \frac{e^{-2y}}{-2} = \sin x + c \quad \dots (I)$

When $x = \frac{\pi}{6}, y = 0$. So eq. (1), becomes

$\therefore \frac{e^0}{-2} = \sin \frac{\pi}{6} + c \therefore -\frac{1}{2} = \frac{1}{2} + c$

$\therefore -\frac{1}{2} - \frac{1}{2} = c \quad \therefore c = -1$

(Given initial condition determines the value of c)

Put in eq. (1), we get

$\therefore \frac{e^{-2y}}{-2} = \sin x - 1$

$\therefore -e^{-2y} = 2\sin x - 2$

$\therefore e^{2y}(2\sin x - 2) + 1 = 0$ is the required particular solution.

(ii) $\frac{y-1}{y+1} + \frac{x-1}{x+1} \cdot \frac{dy}{dx} = 0$

$\therefore \frac{x+1}{x-1} \cdot dx + \frac{y+1}{y-1} \cdot dy = 0$

$\therefore \frac{(x-1)+2}{x-1} \cdot dx + \frac{(y-1)+2}{y-1} \cdot dy = 0$

$\therefore \left(1 + \frac{2}{x-1}\right) \cdot dx + \left(1 + \frac{2}{y-1}\right) \cdot dy = 0$

Integrating , we get

$\int dx + 2 \int \frac{dx}{x-1} + \int dy + 2 \int \frac{dy}{y-1} = 0$

$\therefore x + 2 \log(x-1) + y + 2 \log(y-1) = c$

$\therefore x + y + 2 \log[(x-1)(y-1)] = c \quad \dots (I)$

When $x = 2, y = 2$. So eq. (I), becomes

$\therefore 2 + 2 + 2 \log[(2-1)(2-1)] = c$

$\therefore 4 + 2 \log(1 \times 1) = c$

$\therefore 4 + 2 \log 1 = c$

$\therefore 4 + 2(0) = c$

$\therefore c = 4$ Put in eq. (I), we get

$\therefore x + y + 2 \log[(x-1)(y-1)] = 4$ is required particular solution.

Ex. 3 : Reduce each of the following differential equations to the separated variable form and hence find the general solution.

(i) $1 + \frac{dy}{dx} = \operatorname{cosec}(x + y)$

(ii) $\frac{dy}{dx} = (4x + y + 1)^2$

Solution :

(i) $1 + \frac{dy}{dx} = \operatorname{cosec}(x + y) \quad \dots \text{(I)}$

Put $x + y = u$

$\therefore 1 + \frac{dy}{dx} = \frac{du}{dx}$

Given differential equation becomes

$\frac{du}{dx} = \operatorname{cosec} u$

$\therefore \frac{du}{\operatorname{cosec} u} = dx$

$\therefore \sin u \cdot du = dx$

Integrating both sides, we get

$\therefore \int \sin u \cdot du = \int dx$

$\therefore -\cos u = x + c$

$\therefore x + \cos u + c = 0$

$\therefore x + \cos(x + y) + c = 0 \quad \dots (\because x + y = u)$

(ii) $\frac{dy}{dx} = (4x + y + 1)^2 \quad \dots \text{(I)}$

Put $4x + y + 1 = u$

$\therefore 4 + \frac{dy}{dx} = \frac{du}{dx}$

$\therefore \frac{dy}{dx} = \frac{du}{dx} - 4$

Given differential equation becomes

$\frac{du}{dx} - 4 = u^2$

$\therefore \frac{du}{dx} = u^2 + 4$

$\therefore \frac{du}{u^2 + 4} = dx$

Integrating both sides, we get

$\therefore \int \frac{du}{u^2 + 4} = \int dx$

$\therefore \frac{1}{2} \tan^{-1} \left(\frac{u}{2} \right) = x + c_1$

$\therefore \tan^{-1} \left(\frac{u}{2} \right) = 2x + 2c_1$

$\therefore \tan^{-1} \left(\frac{4x + y + 1}{2} \right) = 2x + c \quad \dots [2c_1 = c]$

EXERCISE 6.3

(1) In each of the following examples verify that the given expression is a solution of the corresponding differential equation.

(i) $xy = \log y + c ; \frac{dy}{dx} = \frac{y^2}{1 - xy}$

(ii) $y = (\sin^{-1} x)^2 + c ; (1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 2$

(iii) $y = e^{-x} + Ax + B ; e^x \frac{d^2y}{dx^2} = 1$

(iv) $y = x^m ; x^2 \frac{d^2y}{dx^2} - mx \frac{dy}{dx} + my = 0$

(v) $y = a + \frac{b}{x} ; x \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} = 0$

(vi) $y = e^{ax} ; x \frac{dy}{dx} = y \log y$

(2) Solve the following differential equations.

(i) $\frac{dy}{dx} = \frac{1+y^2}{1+x^2}$

(ii) $\log\left(\frac{dy}{dx}\right) = 2x + 3y$

(iii) $y - x \frac{dy}{dx} = 0$

(iv) $\sec^2 x \cdot \tan y \cdot dx + \sec^2 y \cdot \tan x \cdot dy = 0$

(v) $\cos x \cdot \cos y \cdot dy - \sin x \cdot \sin y \cdot dx = 0$

(vi) $\frac{dy}{dx} = -k$, where $k = \text{constant}$.

(vii) $\frac{\cos^2 y \cdot dy}{x} + \frac{\cos^2 x \cdot dx}{y} = 0$

(viii) $y^3 - \frac{dy}{dx} = x^2 \frac{dy}{dx}$

(ix) $2e^{x+2y} \cdot dx - 3dy = 0$

(x) $\frac{dy}{dx} = e^{x+y} + x^2 e^y$

(3) For each of the following differential equations find the particular solution satisfying the given condition.

(i) $3e^x \tan y \cdot dx + (1 + e^x) \sec^2 y \cdot dy = 0$, when $x = 0, y = \pi$.

(ii) $(x - y^2 x) \cdot dx - (y + x^2 y) \cdot dy = 0$, when $x = 2, y = 0$.

(iii) $y(1 + \log x) \frac{dx}{dy} - x \log x = 0$, $y = e^2$, when $x = e$.

(iv) $(e^y + 1) \cos x + e^y \sin x \frac{dy}{dx} = 0$, when $x = \frac{\pi}{6}, y = 0$.

(v) $(x + 1) \frac{dy}{dx} - 1 = 2e^{-y}$, $y = 0, x = 1$

(vi) $\cos\left(\frac{dy}{dx}\right) = a, a \in 1^2, y(0) = 2$

(4) Reduce each of the following differential to the variable separable form and hence solve.

(i) $\frac{dy}{dx} = \cos(x + y)$

(ii) $(x - y)^2 \frac{dy}{dx} = a^2$

(iii) $x + y \frac{dy}{dx} = \sec(x^2 + y^2)$

(iv) $\cos^2(x - 2y) = 1 - 2 \frac{dy}{dx}$

(v) $(2x - 2y + 3) dx - (x - y + 1) dy = 0$, when $x = 0, y = 1$.

6.4.1 Homogeneous differential :

Recall that the degree of a term is the sum of the degrees in all variables in the equation, eg. : degree of $3x^2y^2z$ is 5. If all terms have the same degree, the equation is called **homogeneous differential equation**.

For example : (i) $x + y \frac{dy}{dx} = 0$ is a homogeneous differential equation of degree 1.

(ii) $x^3y + xy^3 + x^2y^2 \frac{dy}{dx} = 0$ is a homogeneous differential equation of degree 4.

(iii) $x \frac{dy}{dx} + x^2y = 0$

(iv) $xy \frac{dy}{dx} + y^2 + 2x = 0$

(iii) and (iv) are not homogeneous differential equations.

To solve the homogeneous differential equation, we use the substitution $y = vx$ or $u = vy$.





SOLVED EXAMPLES

Ex. 1 : Solve the following differential equations :

(i) $x^2y \cdot dx - (x^3 + y^3) \cdot dy = 0$ (ii) $x \frac{dy}{dx} = x \tan \left(\frac{y}{x} \right) + y$ (iii) $\frac{dy}{dx} = \frac{y + \sqrt{x^2 + y^2}}{x}$

Solution :

(i) $x^2y \cdot dx - (x^3 + y^3) \cdot dy = 0$

$\therefore x^2y - (x^3 + y^3) \frac{dy}{dx} = 0 \quad \dots (I)$

This is homogeneous Differential equation.

Put $y = vx \quad \dots (II)$

Differentiate w. r. t. x, we get

$\therefore \frac{dy}{dx} = v + x \frac{dv}{dx} \quad \dots (III)$

Put (II) and (III) in Eq. (I), it becomes,

$x^2 \cdot vx - (x^3 + v^3x^3) \left(v + x \frac{dv}{dx} \right) = 0$

divide by x^3 , we get

$v - (1 + v^3) \left(v + x \frac{dv}{dx} \right) = 0$

$\therefore v - v - v^4 - v^3x \frac{dv}{dx} = 0$

$\therefore -x(1 + v^3) \frac{dv}{dx} = v^4$

$\therefore \frac{1 + v^3}{v^4} \cdot dv = - \frac{dx}{x}$

$\therefore \frac{1 + v^3}{v^4} \cdot dv = - \frac{dx}{x}$

$\therefore \left(\frac{1}{v^4} - \frac{v^3}{v^4} \right) dv + \frac{dx}{x} = 0$

integrating eq., we get

$\therefore \int v^{-4} dv + \int \frac{dv}{v} + \int \frac{dx}{x} = c_1$

$\therefore \frac{v^{-3}}{-3} + \log(v) + \log(x) = c_1$

$\therefore \log(vx) = c_1 + \frac{v^{-3}}{3} \quad \therefore \log(y) = c_1 + \frac{1}{3} \cdot \frac{v^{-3}}{x^3}$

$\therefore 3 \log(y) = 3c_1 + \frac{x^3}{y^3}$

$\therefore 3 \log y = \frac{x^3}{y^3} + c \quad \dots \text{where } c = 3c_1$

(ii) $x \frac{dy}{dx} = x \tan \left(\frac{y}{x} \right) + y \quad \dots (I)$

This is homogeneous Differential equation.

Put $y = vx \quad \dots (II)$

Differentiate w. r. t. x, we get

$\therefore \frac{dy}{dx} = v + x \frac{dv}{dx} \quad \dots (III)$

Put (II) and (III) in Eq. (I), it becomes,

$x \left(v + x \frac{dv}{dx} \right) = x \tan \left(\frac{vx}{x} \right) + vx$

divide by x, we get

$\therefore v + x \frac{dv}{dx} = \tan v + v$

$\therefore x \frac{dv}{dx} = \tan v$

$\therefore \frac{dv}{\tan v} = \frac{dx}{x}$

integrating eq., we get

$\therefore \int \cot v dv = \int \frac{dx}{x}$

$\therefore \log(\sin v) = \log(x) + \log c$

$\therefore \log(\sin v) = \log(x \times c)$

$\therefore \sin v = cx$

$\therefore \sin \left(\frac{y}{x} \right) = cx$ is the solution.

$$(iii) \frac{dy}{dx} = \frac{y + \sqrt{x^2 + y^2}}{x} \quad \dots (I)$$

Solution : It is homogeneous differential equation.

$$\text{Put } y = vx \quad \dots (II)$$

Differentiate w. r. t. x , we get

$$\therefore \frac{dy}{dx} = v + x \frac{dv}{dx} \quad \dots (III)$$

Put (II) and (III) in Eq. (I), it becomes,

$$v + x \frac{dv}{dx} = \frac{vx + \sqrt{x^2 + v^2x^2}}{x}$$

$$\therefore v + x \frac{dv}{dx} = v + \sqrt{1 + v^2}$$

$$\therefore x \frac{dv}{dx} = \sqrt{1 + v^2}$$

$$\therefore \frac{dv}{\sqrt{1 + v^2}} = \frac{dx}{x} \quad \dots (IV)$$

integrating eq. (IV), we get

$$\therefore \int \frac{dv}{\sqrt{1 + v^2}} = \int \frac{dx}{x}$$

$$\therefore \log (v + \sqrt{1 + v^2}) = \log (x) + \log c$$

$$\therefore \log (v + \sqrt{1 + v^2}) = \log (cx)$$

$$\therefore v + \sqrt{1 + v^2} = cx$$

$$\therefore \frac{y}{x} + \sqrt{1 + \frac{y^2}{x^2}} = cx$$

$$\therefore y + \sqrt{x^2 + y^2} = cx^2 \text{ is the solution.}$$

EXERCISE 6.4

I. Solve the following differential equations :

$$(1) x \sin \left(\frac{y}{x} \right) dy = \left[y \sin \left(\frac{y}{x} \right) - x \right] dx$$

$$(2) (x^2 - y^2) dx - 2xy \cdot dy = 0$$

$$(3) \left(1 + 2e^{\frac{x}{y}} \right) + 2e^{\frac{x}{y}} \left(1 - \frac{x}{y} \right) \frac{dy}{dx} = 0$$

$$(4) y^2 \cdot dx + (xy + x^2) dy = 0$$

$$(5) (x^2 - y^2) dx + 2xy \cdot dy = 0$$

$$(6) \frac{dy}{dx} + \frac{x - 2y}{2x - y} = 0$$

$$(7) x \frac{dy}{dx} - y + x \sin \left(\frac{y}{x} \right) = 0$$

$$(8) \left(1 + e^{\frac{x}{y}} \right) dx + e^{\frac{x}{y}} \left(1 - \frac{x}{y} \right) dy = 0$$

$$(9) y^2 - x^2 \frac{dy}{dx} = xy \frac{dy}{dx}$$

$$(10) xy \frac{dy}{dx} = x^2 + 2y^2, y(1) = 0$$

$$(11) x dy + 2y \cdot dx = 0, \text{ when } x = 2, y = 1$$

$$(12) x^2 \frac{dy}{dx} = x^2 + xy + y^2$$

$$(13) (9x + 5y) dy + (15x + 11y) dx = 0$$

$$(14) (x^2 + 3xy + y^2) dx - x^2 dy = 0$$

$$(15) (x^2 + y^2) dx - 2xy \cdot dy = 0$$

6.4.2 Linear Differential Equation :

The differential equation of the type, $\frac{dy}{dx} + Py = Q$ (where P, Q are functions of x .)

is called **linear differential equation**.

To get the solution of equation, multiply the equation by $e^{\int P dx}$, which is helping factor here.

We get,

$$e^{\int P dx} \left[\frac{dy}{dx} + Py \right] = Q \cdot e^{\int P dx}$$

Note that, $\frac{d}{dx} [y \cdot e^{\int P dx}] = \left[\frac{dy}{dx} + y \cdot P \right] \cdot e^{\int P dx}$

$\therefore \frac{d}{dx} [y \cdot e^{\int P dx}] = Q \cdot e^{\int P dx}$

$\therefore \int Q \cdot e^{\int P dx} \cdot dx = y \cdot e^{\int P dx}$


Hence, $y \cdot e^{\int P dx} = \int Q \cdot (e^{\int P dx}) dx + c$ is the solution of the given equation

Here $e^{\int P dx}$ is called the integrating factor. (I.F.)

Note : For the linear differential equation.

$\frac{dx}{dy} + py = Q$ (where P, Q are constants or functions of y) the general solution is

x (I.F.) = $\int Q \cdot$ (I.F.) $dy + c$, where I.F. (integrating factor) = $e^{\int P dy}$



SOLVED EXAMPLES

Ex. 1: Solve the following differential equations :

(i) $\frac{dy}{dx} + y = e^{-x}$

(ii) $x \sin \frac{dy}{dx} + (x \cos x + \sin y) = \sin x$

(iii) $(1 + y^2) dx = (\tan^{-1} y - x) dy$

Solution :

(i) $\frac{dy}{dx} + y = e^{-x} \dots$ (I)

This is linear differential equation of the form

$\frac{dy}{dx} + Py = Q$ where $P = 1, Q = e^{-x}$

It's Solution is

y (I.F.) = $\int Q \cdot$ (I.F.) $dx + c \dots$ (II)

where I.F. = $e^{\int P dx} = e^{\int dx} = e^x$

eq. (II) becomes,

$y \cdot e^x = \int e^{-x} \times e^x \cdot dx + c$

$\therefore y \cdot e^x = \int e^{-x+x} \cdot dx + c$

$\therefore y \cdot e^x = \int e^0 \cdot dx + c$

$\therefore y \cdot e^x = \int dx + c$

$\therefore y \cdot e^x = x + c$ is the general solution.

(ii) $x \sin x \frac{dy}{dx} + (x \cos x + \sin x) y = \sin x$

divide by $x \sin x$, we get

$$\frac{dy}{dx} + \left(\cot x + \frac{1}{x} \right) y = \frac{1}{x} \quad \dots (I)$$

It is the linear differential equation of the type

$$\frac{dy}{dx} + Py = Q \quad \text{where} \quad P = \cot x + \frac{1}{x},$$

$$Q = \frac{1}{x}$$

Its solution is

$$y \text{ (I.F.)} = \int Q \cdot \text{(I.F.)} dx + c \quad \dots (II)$$

where I.F. = $e^{\int P dx} = e^{\int (\cot x + \frac{1}{x}) dx}$

$$\text{I.F.} = e^{\int \cot x dx + \int \frac{dx}{x}}$$

$$\text{I.F.} = e^{\log |\sin x| + \log x}$$

$$\text{I.F.} = x \sin x$$

eq. (II) becomes,

$$y \cdot x \sin x = \int \frac{1}{x} \times x \sin x \cdot dx + c$$

$$\therefore xy \cdot \sin x = -\cos x + c$$

$$\therefore xy \cdot \sin x + \cos x = c \text{ is the general solution.}$$

(iii) $(1 + y^2) dx = (\tan^{-1} y - x) dy$

$$\therefore \frac{dx}{dy} = \frac{(\tan^{-1} y - x)}{(1 + y^2)}$$

$$\therefore \frac{dx}{dy} + \left(\frac{1}{1 + y^2} \right) x = \frac{\tan^{-1} y}{1 + y^2}$$

This is linear differential equation of the type

$$\frac{dx}{dy} + Px = Q \quad \text{where} \quad P = \frac{1}{1 + y^2}, \quad Q = \frac{\tan^{-1} y}{1 + y^2}$$

Its solution is

$$x \text{ (I.F.)} = \int Q \cdot \text{(I.F.)} dy + c \quad \dots (II)$$

where I.F. = $e^{\int P dy} = e^{\int \frac{1}{1+y^2} dy}$

$$\text{I.F.} = e^{\tan^{-1} y}$$

eq. (II) becomes,

$$x \cdot e^{\tan^{-1} y} = \int \frac{\tan^{-1} y}{1 + y^2} \cdot e^{\tan^{-1} y} \cdot dy + c \dots (III)$$

in R.H.S. Put $\tan^{-1} y = t$

differentiate w. r. t. x, we get

$$\therefore \frac{dy}{1 + y^2} = dt$$

eq. (III) becomes

$$x \cdot e^{\tan^{-1} y} = \int t \cdot e^t \cdot dt + c$$

$$= t \int e^t \cdot dt - \int [1 \times e^t] dt + c$$

$$= t \cdot e^t - \int e^t \cdot dt + c$$

$$= t \cdot e^t - e^t + c$$

$$x \cdot e^{\tan^{-1} y} = \tan^{-1} y \cdot e^{\tan^{-1} y} - e^{\tan^{-1} y} + c$$

$$\therefore x = \tan^{-1} y - 1 + \frac{c}{e^{\tan^{-1} y}}$$

$$\therefore x + 1 - \tan^{-1} y = c \cdot e^{-\tan^{-1} y} \text{ is the solution.}$$

Ex. 2: The slope of the tangent to the curve at any point is equal to $y + 2x$. Find the equation of the curve passing through the origin.

Solution : Let P (x, y) be any point on the curve $y = f(x)$

The slope of the tangent at point P (x, y) is $\frac{dy}{dx}$.

$$\therefore \frac{dy}{dx} = y + 2x \quad \therefore \frac{dy}{dx} - y = 2x$$

This is linear differential equation of the type

$$\frac{dy}{dx} + Py = Q \text{ where } P = -1, Q = 2x$$

Its solution is

$$y \text{ (I.F.)} = \int Q \cdot \text{(I.F.)} dx + c \quad \dots \text{(I)}$$

where I.F. = $e^{\int P dx} = e^{\int -1 dx}$

$$\text{I.F.} = e^{-x}$$

eq. (I) becomes,

$$y \cdot e^{-x} = \int 2x \times e^{-x} \cdot dx + c$$

$$y \cdot e^{-x} = 2 \int x \cdot e^{-x} \cdot dx + c \quad \dots \text{(II)}$$

Consider, $\int x \cdot e^{-x} \cdot dx$

$$= x \int e^{-x} \cdot dx - \int \left(1 \times \frac{e^{-x}}{-1} \right) dx$$

$$= \frac{x \cdot e^{-x}}{-1} + \int e^{-x} \cdot dx$$

$$= -x e^{-x} \cdot dx + \int e^{-x} \cdot dx$$

$$= -x e^{-x} - e^{-x}$$

(II) becomes

$$y \cdot e^{-x} = 2 [-x e^{-x} - e^{-x}] + c$$

$$\therefore y = -2x - 2 + c e^{-x} \quad \dots \text{(III)}$$

The curve passes through the origin (0, 0)

$$\therefore 0 = -2(0) - 2 + c e^{-0}$$

$$\therefore 0 = -2 + c$$

$$\therefore 2 = c \text{ Put in (III)}$$

$$\therefore y = -2x - 2 + 2e^{-x}$$

$$\therefore 2x + y + 2 = 2e^{-x}$$

is the equation of the curve.

EXERCISE 6.5

(1) Solve the following differential equations :

(i) $\frac{dy}{dx} + \frac{y}{x} = x^3 - 3$

(ii) $\cos^2 x \frac{dy}{dx} + y = \tan x$

(iii) $(x + 2y^3) \frac{dy}{dx} = y$

(iv) $\frac{dy}{dx} + y \sec x = \tan x$

(v) $x \frac{dy}{dx} + 2y = x^2 \log x$

(vi) $(x + y) \frac{dy}{dx} = 1$

(vii) $(x + a) \frac{dy}{dx} - 3y = (x + a)^5$

(viii) $dr + (2r \cot \theta + \sin 2\theta) d\theta = 0$

$$(ix) \quad ydx + (x - y^2) dy = 0$$

$$(x) \quad (1 - x^2) \frac{dy}{dx} + 2xy = x(1 - x^2)^{\frac{1}{2}}$$

$$(xi) \quad (1 + x^2) \frac{dy}{dx} + y = e^{\tan^{-1}x}$$

- (2) Find the equation of the curve which passes through the origin and has slope $x + 3y - 1$ at any point (x, y) on it.
- (3) Find the equation of the curve passing through the point $\left(\frac{3}{\sqrt{2}}, \sqrt{2}\right)$ having slope of the tangent to the curve at any point (x, y) is $-\frac{4x}{9y}$.
- (4) The curve passes through the point $(0, 2)$. The sum of the co-ordinates of any point on the curve exceeds the slope of the tangent to the curve at that point by 5. Find the equation of the curve.
- (5) If the slope of the tangent to the curve at each of its point is equal to the sum of abscissa and the product of the abscissa and ordinate of the point. Also the curve passes through the point $(0, 1)$. Find the equation of the curve.

6.5 Application of differential Equations :

There are many situations where the relation in the rate of change of a function is known. This gives a differential equation of the function and we may be able to solve it.

6.5.1 Population Growth and Growth of Bacteria :

It is known that a number of bacteria in a culture increase with time. It means there is growth in the number of bacteria. If the population P increases at time t then the rate of change of P is proportional to the population present at that time.

$$\therefore \frac{dP}{dt} \propto P$$

$$\therefore \frac{dP}{dt} = k \cdot P, \quad (k > 0)$$

$$\therefore \frac{dP}{P} = k dt$$

on integrating


$$\therefore \int \frac{dP}{P} = \int k dt$$

$$\therefore \log P = kt + c_1$$

$$\therefore P = c \cdot e^{kt} \quad \text{where } c = e^{c_1}$$

which gives the population at any time t .





SOLVED EXAMPLES

Ex. 1 : The population of a town increasing at a rate proportional to the population at that time. If the population increases from 40 thousands to 60 thousands in 40 years, what will be the population in another 20 years.

$$\left(\text{Given } \sqrt{\frac{3}{2}} = 1.2247 \right).$$

Solution : Let P be the population at time t . Since rate of increase of P is a proportional to P itself, we have,

$$\frac{dP}{dt} = k \cdot P \quad \dots (1)$$

where k is constant of proportionality.

Solving this differential equation, we get

$$P = a \cdot e^{kt}, \text{ where } a = e^c \quad \dots (2)$$

Initially $P = 40,000$ when $t = 0$

\therefore From equation (2), we have

$$40,000 = a \cdot 1 \quad \therefore a = 40,000$$

eq. (2) becomes

$$\therefore P = 40,000 \cdot e^{kt} \quad \dots (3)$$

Again given that $P = 60,000$ when $t = 40$

\therefore From equation (3),

$$60,000 = 40,000 \cdot e^{40k}$$

$$e^{40k} = \frac{3}{2} \quad \dots (4)$$

Now we have to find P when $t = 40 + 20 = 60$ years

\therefore From equation (3), we have

$$\begin{aligned} P &= 40,000 \cdot e^{60k} = 40,000 (e^{40k})^{\frac{3}{2}} \\ &= 40,000 \left(\frac{3}{2} \right)^{\frac{3}{2}} = 73482 \end{aligned}$$

\therefore Required population will be 73482.

Ex. 2 : Bacteria increase at the rate proportional to the number of bacteria present. If the original number N doubles in 3 hours, find in how many hours the number of bacteria will be $4N$?

Solution : Let x be the number of bacteria at time t .

Since the rate of increase of x is proportional to x , the differential equation can be written as :

$$\frac{dx}{dt} = kx$$

where k is constant of proportionality.

Solving this differential equation we have

$$x = c_1 \cdot e^{kt}, \text{ where } c_1 = e^c \quad \dots (1)$$

Given that $x = N$ when $t = 0$

\therefore From equation (1) we get

$$N = c_1 \cdot 1$$

$$\therefore c_1 = N$$

$$\therefore x = N \cdot e^{kt} \quad \dots (2)$$

Again given that $x = 2N$ when $t = 3$

\therefore From equation (2), we have

$$2N = N \cdot e^{3k} \quad \dots (3)$$

$$e^{3k} = 2$$

Now we have to find t , when $x = 4N$

\therefore From equation (2), we have

$$4N = N \cdot e^{kt}$$

$$\text{i.e. } 4 = e^{kt} = (e^{3k})^{\frac{t}{3}}$$

$$\therefore 2^2 = 2^{\frac{t}{3}} \quad \dots \text{ by eq. (3)}$$

$$\therefore \frac{t}{3} = 2$$

$$\therefore t = 6$$

Therefore, the number of bacteria will be $4N$ in 6 hours.

6.5.2 Radio Active Decay :

We know that the radio active substances (elements) like radium, cesium etc. disintegrate with time. It means the mass of the substance decreases with time.

The rate of disintegration of such elements is always proportional to the amount present at that time.

If x is the amount of any material present at time t then

$$\frac{dx}{dt} = -k \cdot x$$

where k is the constant of proportionality and $k > 0$. The negative sign appears because x decreases as t increases.

Solving this differential equation we get

$$x = a \cdot e^{kt} \quad \text{where } a = e^c \text{ (check!) } \dots (1)$$

If x_0 is the initial amount of radio active substance at time $t = 0$, then from equation (1)

$$\begin{aligned} x_0 &= a \cdot 1 \\ \therefore a &= x_0 \\ \therefore x &= x_0 e^{-kt} \quad \dots (2) \end{aligned}$$

This expression gives the amount of radio active substance at any time t .

Half Life Period :

Half life period of a radio active substance is defined as the time it takes for half the amount/mass of the substance to disintegrate.

Ex. 3 : Bismuth has half life of 5 days. A sample originally has a mass of 800 mg. Find the mass remaining after 30 days.

Solution : Let x be the mass of the Bismuth present at time t .

$$\text{Then } \frac{dx}{dt} = -k \cdot x \quad \text{where } k > 0$$

Solving the differential equation, we get

$$x = c \cdot e^{-kt} \quad \dots (1)$$

where c is constant of proportionality.

Given that $x = 800$, when $t = 0$

using these values in equation (1), we get

$$800 = c \cdot 1 = c$$

$$\therefore x = 800 e^{-kt} \quad \dots (2)$$

Since half life is 5 days, we have $x = 400$ when $t = 5$,

Now let us discuss another application of differential equation.

\therefore From equation (2), we have

$$400 = 800 e^{-5k}$$

$$\therefore e^{-5k} = \frac{400}{800} = \frac{1}{2} \quad \dots (3)$$

Now we have determine x , when $t = 30$,

\therefore From equation (2), we have

$$x = 800 e^{-30k} = 12 \cdot 5 \text{ (verify !)}$$

\therefore The mass after 30 days will be 12.5 mg.

6.5.3 Newton's Law of Cooling :

Newton's law of cooling states that the rate of change of cooling heated body at any time is proportional to the difference between the temperature of a body and that of its surrounding medium.

Let θ be the temperature of a body at time t and θ_0 be the temperature of the medium.

Then $\frac{d\theta}{dt}$ is the rate of change of temperature with respect to time t and $\theta - \theta_0$ is the difference of temperature at time t . According to Newton's law of cooling.

$$\begin{aligned} \therefore \quad & \frac{d\theta}{dt} \propto (\theta - \theta_0) \\ \therefore \quad & \frac{d\theta}{dt} = -k (\theta - \theta_0) \qquad \dots (1) \end{aligned}$$

where k is constant of proportionality and negative sign indicates that difference of temperature is decreasing.

$$\begin{aligned} \text{Now} \quad & \frac{d\theta}{dt} = -k (\theta - \theta_0) \\ \therefore \quad & \frac{d\theta}{(\theta - \theta_0)} = -k dt \end{aligned}$$

\therefore Integrating and using the initial condition viz.

$$\begin{aligned} \therefore \quad & \theta = \theta_1 \quad \text{when } t = 0, \text{ we get} \\ \therefore \quad & \theta = \theta_0 + (\theta_1 - \theta_0)e^{-kt} \text{ (verify)} \qquad \dots (2) \end{aligned}$$

Thus equation (2) gives the temperature of a body at any time t .

Ex. 4 : Water at 100°c cools in 10 minutes to 88°c in a room temperature of 25°c . Find the temperature of water after 20 minutes.

Solution : Let θ be the temperature of water at time t . Room temperature is given to be 25°c . Then according to Newton's law of cooling. we have

$$\begin{aligned} \frac{d\theta}{dt} & \propto (\theta - 25) \\ \frac{d\theta}{dt} & = -k (\theta - 25), \quad \text{where } k > 0 \end{aligned}$$

After integrating and using initial condition.

$$\text{We get } \theta = 25 + 75 \cdot e^{-kt} \qquad \dots (1)$$

But given that $\theta = 88^\circ\text{c}$ when $t = 10$

\therefore From equation (1) we get

$$88 = 25 + 75 \cdot e^{-10k}$$

$$\therefore 63 = 75 \cdot e^{-10k} \therefore e^{-10k} = \frac{63}{75} = \frac{21}{25} \qquad \dots (2)$$

Now we have to find θ , when $t = 20$,

\therefore From equation (1) we have

$$\begin{aligned} \theta & = 25 + 75 \cdot e^{-20k} \\ & = 25 + 75 (e^{-20k})^2 \\ & = 25 + 75 \left(\frac{dy}{dx}\right)^2 \qquad \dots \text{by (2)} \\ & = 25 + \frac{75 \times 21 \times 21}{25 \times 25} \\ & = 25 + \frac{1323}{25} = 77.92 \end{aligned}$$

Therefore temperature of water after 20 minutes will be 77.92°c .




6.5.4 Surface Area :

Knowledge of a differential equation is also used to solve problems related to the surface area. We consider the following examples :

Ex. 5 : Water is being poured into a vessel in the form of an inverted right circular cone of semi vertical angle 45° in such a way that the rate of change of volume at any moment is proportional to the area of the curved surfaces which is wet at that moment. Initially, the vessel is full to a height of 2 cms. And after 2 seconds the height becomes 10 cm. Show that after 3.5 seconds from that start, the height of water will be 16 cms.

Solution : Let the height of water at time t seconds be h cms.



We are given that initial height is 2 cms. and after 2 seconds, the height is 10 cms.

$$\therefore h = 2 \text{ when } t = 0 \quad \dots (1)$$

$$\text{and } h = 10 \text{ when } t = 2 \quad \dots (2)$$

Let v be the volume, r be the radius of the water surface and l be that slant height at time t seconds.

\therefore Area of the curved surface at this moment is $\pi r l$.

But the semi vertical angle is 45° .

$$\therefore \tan 45^\circ = \frac{r}{h} = 1$$

$$\therefore r = h$$

$$\text{and } l^2 = r^2 + h^2 = 2h^2$$

$$\therefore l = \sqrt{2} h$$

$$\begin{aligned} \therefore \text{Area of the curved surface} &= \pi r l = \pi \cdot h \cdot \sqrt{2} h \\ &= \sqrt{2} \pi h^2 \end{aligned}$$

Since rate of change of volume is proportional to this area, we get

$$\frac{dv}{dt} \propto \sqrt{2} \pi h^2$$

$$\therefore \frac{dv}{dt} = c \cdot \sqrt{2} \pi h^2$$

where c is constant of proportionality.

Let $c\sqrt{2} \pi = k$

$$\therefore \frac{dv}{dt} = kh^2 \quad \dots (3)$$

\therefore where k is constant

Now $v = \frac{1}{3} \pi r^2 h$

$$= \frac{1}{3} \pi r^2 \cdot h = \frac{1}{3} \pi h^3, \text{ (since } r = h\text{)}$$

Differentiating with respect to t , we get

$$\therefore \frac{dv}{dt} = \pi h^2 \frac{dh}{dt} \quad \dots (4)$$

Equating $\frac{dv}{dt}$ from (3) and (4) we get

$$\pi h^2 \frac{dh}{dt} = kh^2$$

$$\therefore \frac{dh}{dt} = \frac{k}{\pi} = a \text{ (say)}$$

where a is constant.

integrating we get

$$h = at + b \quad \dots (5)$$

using (1) we have

$$2 = a \cdot 0 + b \quad \therefore b = 2$$

\therefore Equation (5) becomes

$$h = at + 2$$

Now using (2) we get

$$10 = 2a + 2 \quad \therefore a = 4$$

using the values of a and b in equation (5), we have

$$\therefore h = 4t + 2$$

Now put $t = 3.5$

$$\therefore h = 4 \times 3.5 + 2$$

$$= 14 + 2 = 16 \text{ cm}$$

Therefore, height of water after 3.5 seconds will be 16 cms.



EXERCISE 6.6

1. In a certain culture of bacteria the rate of increase is proportional to the number present. If it is found that the number doubles in 4 hours, find the number of times the bacteria are increased in 12 hours.
2. If the population of a country doubles in 60 years, in how many years will it be triple (treble) under the assumption that the rate of increase is proportional to the number of inhabitants?
[Given $\log 2 = 0.6912$, $\log 3 = 1.0986$]
3. If a body cools from 80°C to 50°C at room temperature of 25°C in 30 minutes, find the temperature of the body after 1 hour.
4. The rate of growth of bacteria is proportional to the number present. If initially, there were 1000 bacteria and the number double in 1 hour, find the number of bacteria after $2\frac{1}{2}$ hours.
[Take $\sqrt{2} = 1.414$]
5. The rate of disintegration of a radio active element at any time t is proportional to its mass at that time. Find the time during which the original mass of 1.5 gm. will disintegrate into its mass of 0.5 gm.
6. The rate of decay of certain substance is directly proportional to the amount present at that instant. Initially, there are 25 gms of certain substance and two hours later it is found that 9 gms are left. Find the amount left after one more hour.
7. Find the population of a city at any time t , given that the rate of increase of population is proportional to the population at the instant and that in a period of 40 years the population increased from 30,000 to 40,000.
8. A body cools according to Newton's law from 100°C to 60°C in 20 minutes. The temperature of the surrounding being 20°C how long will it take to cool down to 30°C ?
9. A right circular cone has height 9 cms and radius of the base 5 cms. It is inverted and water is poured into it. If at any instant the water level rises at the rate of $\frac{\pi}{A}$ cms/ sec. where A is the area of water surface at that instant, show that the vessel will be full in 75 seconds.
10. Assume that a spherical raindrop evaporates at a rate proportional to its surface area. If its radius originally is 3mm and 1 hour later has been reduced to 2mm, find an expression for the radius of the raindrop at any time t .
11. The rate of growth of the population of a city at any time t is proportional to the size of the population. For a certain city it is found that the constant of proportionality is 0.04. Find the population of the city after 25 years if the initial population is 10,000. [Take $e = 2.7182$]
12. Radium decomposes at the rate proportional to the amount present at any time. If p percent of amount disappears in one year, what percent of amount of radium will be left after 2 years ?



Let us Remember

- ✿ Equation which contains the derivative of a function is called a **differential equation**.
- ✿ The order of a differential equation is the highest order of the derivative appearing in the equation.
- ✿ The degree of the differential equation is the power of the highest ordered derivative present in the equation.
- ✿ Order and degree of a differential equation are always positive integers.
- ✿ Solution of a differential equation in which number of arbitrary constants is equal to the order of a differential equation is the **general solution** of the differential equation.
- ✿ Solution obtained from the general solution by giving particular values to the arbitrary constants is the particular solution of the differential equation.
- ✿ The most general form of a **linear differential equation** of the first order is : $\frac{dy}{dx} + Py = Q$ where P and Q are functions of x or constant.
Its solution is given by : $y \text{ (I.F.)} = \int Q \cdot \text{(I.F.)} dx + c$, where I.F. (integrating factor) = $e^{\int P dx}$
- ✿ Solution of a differential equation $\frac{dx}{dt} = kx$ is in the form $x = a \cdot e^{kt}$ where a is initial value of x . Further, $k > 0$ represents growth and $k < 0$, represents decay.
- ✿ Newton's law of cooling is $\theta = \theta_0 + (\theta_1 - \theta_0)e^{-kt}$.

MISCELLANEOUS EXERCISE 6

(I) Choose the correct option from the given alternatives :

- (1) The order and degree of the differential equation $\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \left(\frac{d^2y}{dx^2}\right)^{\frac{3}{2}}$ are respectively . . .
- (A) 2, 1 (B) 1, 2 (C) 3, 2 (D) 2, 3
- (2) The differential equation of $y = c^2 + \frac{c}{x}$ is . . .
- (A) $x^4 \left(\frac{dy}{dx}\right)^2 - x \frac{dy}{dx} = y$ (B) $\frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = 0$
- (C) $x^3 \left(\frac{dy}{dx}\right)^2 + x \frac{dy}{dx} = y$ (D) $\frac{d^2y}{dx^2} + \frac{dy}{dx} - y = 0$

(3) $x^2 + y^2 = a^2$ is a solution of ...

(A) $\frac{d^2y}{dx^2} + \frac{dy}{dx} - y = 0$

(B) $y = x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} + a^2y$

(C) $y = x \frac{dy}{dx} + a \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$

(D) $\frac{d^2y}{dx^2} = (x + 1) \frac{dy}{dx}$

(4) The differential equation of all circles having their centers on the line $y = 5$ and touching the X-axis is

(A) $y^2 \left(1 + \frac{dy}{dx}\right) = 25$

(B) $(y - 5)^2 \left[1 + \left(\frac{dy}{dx}\right)^2\right] = 25$

(C) $(y - 5)^2 + \left[1 + \left(\frac{dy}{dx}\right)^2\right] = 25$

(D) $(y - 5)^2 \left[1 - \left(\frac{dy}{dx}\right)^2\right] = 25$

(5) The differential equation $y \frac{dy}{dx} + x = 0$ represents family of ...

(A) circles

(B) parabolas

(C) ellipses

(D) hyper bolas

(6) The solution of $\frac{1}{x} \cdot \frac{dy}{dx} = \tan^{-1} x$ is ...

(A) $\frac{x^2 \tan^{-1} x}{2} + c = 0$

(B) $x \tan^{-1} x + c = 0$

(C) $x - \tan^{-1} x = c$

(D) $y = \frac{x^2 \tan^{-1} x}{2} - \frac{1}{2} (x - \tan^{-1} x) + c$

(7) The solution of $(x + y)^2 \frac{dy}{dx} = 1$ is ...

(A) $x = \tan^{-1} (x + y) + c$

(B) $y \tan^{-1} \left(\frac{x}{y}\right) = c$

(C) $y = \tan^{-1} (x + y) + c$

(D) $y + \tan^{-1} (x + y) = c$

(8) The solution of $\frac{dy}{dx} = \frac{y + \sqrt{x^2 - y^2}}{2}$ is ...

(A) $\sin^{-1} \left(\frac{y}{x}\right) = 2 \log |x| + c$

(B) $\sin^{-1} \left(\frac{y}{x}\right) = \log |x| + c$

(C) $\sin \left(\frac{y}{x}\right) = \log |x| + c$

(D) $\sin \left(\frac{y}{x}\right) = \log |x| + c$

(9) The solution of $\frac{dy}{dx} + y = \cos x - \sin x$ is ...

(A) $y e^x = \cos x + c$

(B) $y e^x + e^x \cos x = c$

(C) $y e^x = e^x \cos x + c$

(D) $y^2 e^x = e^x \cos x + c$

- (10) The integrating factor of linear differential equation $x \frac{dy}{dx} + 2y = x^2 \log x$ is . . .
 (A) $\frac{1}{x}$ (B) k (C) $\frac{1}{n^2}$ (D) x^2
- (11) The solution of the differential equation $\frac{dy}{dx} = \sec x - y \tan x$ is
 (A) $y \sec x + \tan x = c$ (B) $y \sec x = \tan x + c$
 (C) $\sec x + y \tan x = c$ (D) $\sec x = y \tan x + c$
- (12) The particular solution of $\frac{dy}{dx} = xe^{y-x}$, when $x = y = 0$ is . . .
 (A) $e^{x-y} = x + 1$ (B) $e^{x+y} = x + 1$ (C) $e^x + e^y = x + 1$ (D) $e^{y-x} = x - 1$
- (13) $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ is a solution of . . .
 (A) $\frac{d^2y}{dx^2} + yx + \left(\frac{dy}{dx}\right)^2 = 0$ (B) $xy \frac{d^2y}{dx^2} + 2 \left(\frac{dy}{dx}\right)^2 - y \frac{dy}{dx} = 0$
 (C) $y \frac{d^2y}{dx^2} + 2 \left(\frac{dy}{dx}\right)^2 + y = 0$ (D) $xy \frac{dy}{dx} + y \frac{d^2y}{dx^2} = 0$
- (14) The decay rate of certain substance is directly proportional to the amount present at that instant. Initially there are 27 grams of substance and 3 hours later it is found that 8 grams left. The amount left after one more hour is...
 (A) $5 \frac{2}{3}$ grams (B) $5 \frac{1}{3}$ grams (C) 5.1 grams (D) 5 grams
- (15) If the surrounding air is kept at 20°C and a body cools from 80°C to 70°C in 5 minutes, the temperature of the body after 15 minutes will be...
 (A) 51.7°C (B) 54.7°C (C) 52.7°C (D) 50.7°C

(II) Solve the following :

- (1) Determine the order and degree of the following differential equations :

(i) $\frac{d^2y}{dx^2} + 5 \frac{dy}{dx} + y = x^3$

(ii) $\left(\frac{d^3y}{dx^3}\right)^2 = \sqrt[5]{1 + \frac{dy}{dx}}$

(iii) $\sqrt[3]{1 + \left(\frac{dy}{dx}\right)^2} = \frac{d^2y}{dx^2}$

(iv) $\frac{dy}{dx} = 3y + \sqrt[4]{1 + 5 \left(\frac{dy}{dx}\right)^2}$

(v) $\frac{d^4y}{dx^4} + \sin\left(\frac{dy}{dx}\right) = 0$

(6) Find the particular solution of the following differential equations :

(1) $y(1 + \log x) = (\log x^x) \frac{dy}{dx}$, when $y(e) = e^2$

(2) $(x + 2y^2) \frac{dy}{dx} = y$, when $x = 2, y = 1$

(3) $\frac{dy}{dx} - 3y \cot x = \sin 2x$, when $y\left(\frac{\pi}{2}\right) = 2$

(4) $(x + y) dy + (x - y) dx = 0$, when $x = 1 = y$

(5) $2e^{\frac{x}{y}} dx + \left(y - 2xe^{\frac{x}{y}}\right) dy = 0$, when $y(0) = 1$

(7) Show that the general solution of the differential equation $\frac{dy}{dx} = \frac{y^2 + y + 1}{x^2 + x + 1}$ is given by $(x + y + 1) = c(1 - x - y - 2xy)$

(8) The normal lines to a given curve at each point (x, y) on the curve pass through $(2, 0)$. The curve passes through $(2, 3)$. Find the equation of the curve.

(9) The volume of spherical balloon being inflated changes at a constant rate. If initially its radius is 3 units and after 3 seconds it is 6 units. Find the radius of the balloon after t second.

(10) A person's assets start reducing in such a way that the rate of reduction of assets is proportional to the square root of the assets existing at that moment. If the assets at the beginning are ₹ 10 lakhs and they dwindle down to ₹ 10,000 after 2 years, show that the person will be bankrupt in $2\frac{2}{9}$ years from the start.



7. PROBABILITY DISTRIBUTIONS




Let us Learn

- Random variables
- Types of random variables
- Probability distribution of random variable.
 - Discrete random variable
 - Probability mass function
 - Expected values and variance
 - Continuous random variable
 - Probability density function
 - Cumulative distribution function



Let us Recall

- A random experiment and all possible outcomes of an experiment
- The sample space of a random experiment



Let us Study

7.1 Random variables :

We have already studied random experiments and sample spaces corresponding to random experiments. As an example, consider the experiment of tossing two fair coins. The sample space corresponding to this experiment contains four elements, namely $\{HH, HT, TH, TT\}$. We have already learnt to construct the sample space of any random experiment. However, the interest is not always in a random experiment and its sample space. We are often not interested in the outcomes of a random experiment, but only in some number obtained from the outcome. For example, in case of the experiment of tossing two fair coins, our interest may be only in the number of heads when two coins are tossed. In general, it is possible to associate a unique real number to every possible outcome of a random experiment. The number obtained from an outcome of a random experiment can take different values for different outcomes. This is why such a number is a variable. The value of this variable depends on the outcome of the random experiment, therefore it is called a random variable.

A random variable is usually denoted by capital letters like X, Y, Z, \dots

Consider the following examples to understand the concept of random variables.

- (i) When we throw two dice, there are 36 possible outcomes, but if we are interested in the sum of the numbers on the two dice, then there are only 11 different possible values, from 2 to 12.
- (ii) If we toss a coin 10 times, then there are $2^{10} = 1024$ possible outcomes, but if we are interested in the number of heads among the 10 tosses of the coin, then there are only 11 different possible values, from 0 to 10.
- (iii) In the experiment of randomly selecting four items from a lot of 20 items that contains 6 defective items, the interest is in the number of defective items among the selected four items. In this case, there are only 5 different possible outcomes, from 0 to 4.

In all the above examples, there is a rule to assign a unique value to every possible outcome of the random experiment. Since this number can change from one outcome to another, it is a variable. Also, since this number is obtained from outcomes of a random experiment, it is called a random variable.

A random variable is formally defined as follows.

Definition :

A random variable is a real-valued function defined on the sample space of a random experiment. In other words, the domain of a random variable is the sample space of a random experiment, while its co-domain is the set of real numbers.

Thus $X : S \rightarrow R$ is a random variable.

We often use the abbreviation r.v. to denote a random variable. Consider an experiment where three seeds are sown in order to find how many of them germinate. Every seed will either germinate or will not germinate. Let us use the letter Y when a seed germinates and the letter N when a seed does not germinate. The sample space of this experiment can then be written as

$$S = \{YYY, YYN, YNY, NYY, YNN, NYN, NNY, NNN\}, \text{ and } n(S) = 2^3 = 8.$$

None of these outcomes is a number. We shall try to represent every outcome by a number. Consider the number of times the letter Y appears in a possible outcome and denote it by X . Then, we have $X(YYY) = 3, X(YYN) = X(YNY) = X(NYY) = 2, X(YNN) = X(NYN) = X(NNY) = 1, X(NNN) = 0$.

The variable X has four possible values, namely 0, 1, 2, and 3. The set of possible values of X is called the range of X . Thus, in this example, the range of X is the set $\{0, 1, 2, 3\}$.

A random variable is usually denoted by a capital letter, like X or Y . A particular value taken by the random variable is usually denoted by the small letter x . Note that x is always a real number and the set of all possible outcomes corresponding to a particular value x of X is denoted by the event $[X = x]$.



For example, in the experiment of three seeds, the random variable X takes four possible values, namely 0, 1, 2, 3. The four events are then defined as follows.

$$[X = 0] = \{NNN\},$$

$$[X = 1] = \{YNN, NYN, NNY\},$$

$$[X = 2] = \{YYN, YNY, NYY\},$$

$$[X = 3] = \{YYY\}.$$

Note that the sample space in this experiment is finite and so is the random variable defined on it.

A sample space need not always be finite. For example, the experiment of tossing a coin until a head is obtained. The sample space for this experiment is $S = \{H, TH, TTH, TTTH, \dots\}$.

Note that S contains an unending sequence of tosses required to get a head. Here, S is countably infinite. The random variable $X : S \rightarrow R$, denoting the number of tosses required to get a head, has the range $\{1, 2, 3, \dots\}$ which is also countably infinite.

7.2 Types of Random Variables :

There are two types of random variables, namely discrete and continuous.

7.2.1 Discrete Random Variables :

Definition : A random variable is said to be a discrete random variable if the number of its possible values is finite or countably infinite.

The values of a discrete random variable are usually denoted by non-negative integers, that is, $\{0, 1, 2, \dots\}$.

Examples of discrete random variables include the number of children in a family, the number of patients in a hospital ward, the number of cars sold by a dealer, number of stars in the sky and so on.

Note : The values of a discrete random variable are obtained by counting.

7.2.2 Continuous Random Variable :

Definition : A random variable is said to be a continuous random variable if the possible values of this random variable form an interval of real numbers.

A continuous random variable has uncountably infinite possible values and these values form an interval of real numbers.

Examples of continuous random variables include heights of trees in a forest, weights of students in a class, daily temperature of a city, speed of a vehicle, and so on.



The value of a continuous random variable is obtained by measurement. This value can be measured to any degree of accuracy, depending on the unit of measurement. This measurement can be represented by a point in an interval of real numbers.

The purpose of defining a random variable is to study its properties. The most important property of a random variable is its probability distribution. Many other properties of a random variable are obtained with help of its probability distribution. We shall now learn about the probability distribution of a random variable. We shall first learn the probability of a discrete random variable, and then learn the probability distribution of a continuous random variable.

7.3 Probability Distribution of Discrete Random Variables :

Let us consider the experiment of throwing two dice and noting the numbers on the upper-most faces of the two dice. The sample space of this experiment is

$$S = \{(1, 1), (1, 2), \dots, (6, 6)\} \text{ and } n(S) = 36.$$

Let X denote the sum of the two numbers in any single throw.

Then $\{2, 3, \dots, 12\}$ is the set of possible values of X . Further,

$$[X = 2] = \{(1, 1)\},$$

$$[X = 3] = \{(1, 2), (2, 1)\},$$

...

...

$$[X = 12] = \{(6, 6)\}.$$

Next, all of these 36 possible outcomes are equally likely if the two dice are fair, that is, if each of the six faces have the same probability of being uppermost when the die is thrown.

As the result, each of these 36 possible outcomes has the same probability = $\frac{1}{36}$

This leads to the following results.

$$P[X = 2] = P\{(1, 1)\} = \frac{1}{36},$$

$$P[X = 3] = P\{(1, 2), (2, 1)\} = \frac{2}{36},$$

$$P[X = 4] = P\{(1, 3), (2, 2), (3, 1)\} = \frac{3}{36},$$

and so on.



The following table shows the probabilities of all possible values of X .

x	2	3	4	5	6	7	8	9	10	11	12
$P(x)$	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$

Table 7.1

Such a description giving the values of the random variable X along with the corresponding probabilities is called the probability distribution of the random variable X .

In general, the probability distribution of a discrete random variable X is defined as follows.

Definition : The probability distribution of a discrete random variable X is defined by the following system of numbers. Let the possible values of X be denoted by x_1, x_2, x_3, \dots , and the corresponding probabilities be denoted by p_1, p_2, p_3, \dots , where $p_i = P[X = x_i]$ for $i = 1, 2, 3, \dots$.

Note : A discrete random variable can have finite or infinite number of possible values, but they are countable.

Sometimes, the probability distribution of a discrete random variable is presented in the form of ordered pairs of the form $(x_1, p_1), (x_2, p_2), (x_3, p_3), \dots$. A common practice is to present the probability distribution of a discrete random variable in a tabular form as shown below.

x_i	x_1	x_2	x_3	\dots
$P[X = x_i]$	p_1	p_2	p_3	\dots

Table 7.2

Note : If x_i is a possible value of X and $p_i = P[X = x_i]$, then there is an event $[E_i]$ in the sample space S such that $p_i = P[E_i]$. Since x_i is a possible value of X , $p_i = P[X = x_i] > 0$. Also, all possible values of X cover all sample points in the sample space S , and hence the sum of their probabilities is 1. That is, $p_i \geq 0$, for all i and $\sum_i p_i = 1$.

7.3.1 Probability Mass Function (p. m. f.) :

Sometimes the probability p_i of X taking the value x_i is a function of x_i for every possible value of X . Such a function is called the **probability mass function (p. m. f.)** of the discrete random variable X .

For example, consider the coin-tossing experiment where the random variable X is defined as the number of tosses required to get a head. Let probability of getting head be ' t ' and that of not getting head be $1 - t$. The possible values of X are given by the set of natural numbers, $1, 2, 3, \dots$ and

$$P[X = i] = (1 - t)^{i-1} t, \text{ for } i = 1, 2, 3, \dots$$

This result can be verified by noting that if head is obtained for the first time on the i^{th} toss, then the first $i - 1$ tosses have resulted in tail. In other words, $X = i$ represents the event of having $i - 1$ tails followed by the first head on the toss.

We now define the probability mass function (p. m. f.) of a discrete random variable.

Definition : Let the possible values of a discrete random variable X be denoted by x_1, x_2, x_3, \dots , with the corresponding probabilities $p_i = P [X = x_i]$, $i = 1, 2, \dots$. If there is a function f such that $p_i = P [X = x_i] = f(x_i)$ for all possible values of X , then f is called the probability mass function (p. m. f.) of X .

For example, consider the experiment of tossing a coin 4 times and defining the random variable X as the number of heads in 4 tosses. The possible values of X are 0, 1, 2, 3, 4, and the probability distribution of X is given by the following table.

x	0	1	2	3	4
$P [X = x]$	$\frac{1}{16}$	$\frac{1}{4}$	$\frac{3}{8}$	$\frac{1}{4}$	$\frac{1}{16}$

Table 7.3

Note that :
$$P [X = x] = \binom{4}{x} \left(\frac{1}{2}\right)^4, x = 0, 1, 2, 3, 4, \dots$$

where $\binom{4}{x}$ is the number of ways of getting x heads in 4 tosses.

7.3.2 Cumulative Distribution Function (c. d. f.) :


The probability distribution of a discrete random variable can be specified with help of the p. m. f. It is sometimes more convenient to use the cumulative distribution function (c.d.f.) of the random variable. The cumulative distribution function (c. d. f.) of a discrete random variable is defined as follows.

Definition : The cumulative distribution function (c. d. f.) of a discrete random variable X is denoted by F and is defined as follows.

$$\begin{aligned} F(x) = P [X \leq x] &= \sum_{x_i \leq x} P [X = x_i] \\ &= \sum_{x_i \leq x} p_i \\ &= \sum_{x_i \leq x} f(x_i) \end{aligned}$$

where f is the probability mass function (p. m. f.) of the discrete random variable X .





SOLVED EXAMPLES

Ex. 1 : Two persons A and B play a game of tossing a coin thrice. If the result of a toss is head, A gets ₹ 2 from B. If the result of a toss is tail, B gets ₹ 1.50 from A. Let X denote the amount gained or lost by A. Show that X is a discrete random variable and show how it can be defined as a function on the sample space of the experiment.

Solution : X is a number whose value depends on the outcome of a random experiment.

Therefore, X is a random variable. Since the sample space of the experiment has only 8 possible outcomes, X is a discrete random variable. Now, the sample space of the experiment is

$$S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}.$$

The values of X in rupees corresponding to these outcomes of the experiment are as follows.

$$X(HHH) = 2 \times 3 = ₹ 6$$

$$X(HHT) = X(HTH) = X(THH) = 2 \times 2 - 1.50 \times 1 = ₹ 2.50$$

$$X(HTT) = X(THT) = X(TTH) = 2 \times 1 - 1.50 \times 2 = ₹ - 1.00$$

$$X(TTT) = - 1.50 \times 3 = ₹ - 4.50$$

Here, a negative amount shows a loss to player A. This example shows that X takes a unique value for every element of the sample space and therefore X is a function on the sample space. Further, possible values of X are 4.50, 1, 2.50, 6.

Ex. 2 : A bag contains 1 red and 2 green balls. One ball is drawn from the bag at random, its colour is noted, and then ball is put back in the bag. One more ball is drawn from the bag at random and its colour is also noted. Let X denote the number of red balls drawn from the bag as described above. Derive the probability distribution of X .

Solution : Let the balls in the bag be denoted by r, g_1, g_2 . The sample space of the experiment is then given by

$$S = \{r r, r g_1, r g_2, g_1 r, g_2 r, g_1 g_1, g_1 g_2, g_2 g_1, g_2 g_2\}.$$

Since X is defined as the number of red balls, we have

$$X(\{r r\}) = 2,$$

$$X(\{r g_1\}) = X(r g_2) = X(g_1 r) = X(g_2 r) = 1,$$

$$X(\{g_1 g_1\}) = X(g_1 g_2) = X(g_2 g_1) = X(g_2 g_2) = 0.$$

Thus, X is a discrete random variable that can take values 0, 1, and 2.

The probability distribution of X is then obtained as follows :

x	0	1	2
$P[X = x]$	$\frac{4}{9}$	$\frac{4}{9}$	$\frac{1}{9}$

Ex. 3 : Two cards are randomly drawn, with replacement, from a well shuffled deck of 52 playing cards. Find the probability distribution of the number of aces drawn.

Solution : Let X denote the number of aces among the two cards drawn with replacement.

Clearly, 0, 1, and 2 are the possible values of X . Since the draws are with replacement, the outcomes of the two draws are independent of each other. Also, since there are 4 aces in the deck of 52 cards,

$$P[\text{ace}] = \frac{4}{52} = \frac{1}{13} \text{ and } P[\text{non-ace}] = \frac{12}{13}. \text{ Then}$$

$$P[X = 0] = P[\text{non-ace and non-ace}] = \frac{12}{13} \times \frac{12}{13} = \frac{144}{169},$$

$$\begin{aligned} P[X = 1] &= P[\text{ace and non-ace}] + P[\text{non-ace and ace}] \\ &= \frac{1}{13} \times \frac{12}{13} + \frac{12}{13} \times \frac{1}{13} = \frac{24}{169}, \end{aligned}$$

$$\text{and } P[X = 2] = P[\text{ace and ace}] = \frac{1}{13} \times \frac{1}{13} = \frac{1}{169}.$$

The required probability distribution is then as follows.

x	0	1	2
$P[X = x]$	$\frac{144}{169}$	$\frac{24}{169}$	$\frac{1}{169}$

Ex. 4 : A fair die is thrown. Let X denote the number of factors of the number on the upper face. Find the probability distribution of X .

Solution : The sample space of the experiment is $S = \{1, 2, 3, 4, 5, 6\}$. The values of X for the possible outcomes of the experiment are as follows.

$X(1) = 1, X(2) = 2, X(3) = 2, X(4) = 3, X(5) = 2, X(6) = 4$. Therefore,

$$p_1 = P[X = 1] = P[\{1\}] = \frac{1}{6}$$

$$p_2 = P[X = 2] = P[\{2, 3, 5\}] = \frac{3}{6}$$

$$p_3 = P[X = 3] = P[\{4\}] = \frac{1}{6}$$

$$p_4 = P[X = 4] = P[\{6\}] = \frac{1}{6}$$

The probability distribution of X is then as follows.

x	1	2	3	4
$P[X = x]$	$\frac{1}{6}$	$\frac{3}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

Ex. 5 : Find the probability distribution of the number of doublets in three throws of a pair of dice.

Solution : Let X denote the number of doublets. Possible doublets in a pair of dice are (1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6).

Since the dice are thrown thrice, 0, 1, 2, and 3 are possible values of X . Probability of getting a doublet in a single throw of a pair of dice is $p = \frac{1}{6}$ and $q = 1 - \frac{1}{6} = \frac{5}{6}$.

$$P[X = 0] = P[\text{no doublet}] = qqq = \frac{5}{6} \times \frac{5}{6} \times \frac{5}{6} = \frac{125}{216}.$$

$$P[X = 1] = P[\text{one doublet}] = pqq + qpq + qqp = 3pq^2 = \frac{75}{216}.$$

$$P[X = 2] = P[\text{two doublets}] = ppq + pqp + qpp = 3p^2q = \frac{15}{216}.$$

$$P[X = 3] = P[\text{three doublets}] = ppp = \frac{1}{216}.$$

Ex. 6 : The probability distribution of X is as follows :

x	0	1	2	3	4
$P[X = x]$	0.1	k	$2k$	$2k$	k

Find (i) k , (ii) $P[X < 2]$, (iii) $P[X \geq 3]$, (iv) $P[1 \leq X < 4]$, (v) $F(2)$.

Solution : The table gives a probability distribution and therefore

$$P[X = 0] + P[X = 1] + P[X = 2] + P[X = 3] + P[X = 4] = 1.$$

$$\text{That is, } 0.1 + k + 2k + 2k + k = 1.$$

$$\text{That is, } 6k = 0.9. \text{ Therefore } k = 0.15.$$

(i) $k = 0.15.$

(ii) $P[X < 2] = P[X = 0] + P[X = 1] = 0.1 + k = 0.1 + 0.15 = 0.25$

(iii) $P[X \geq 3] = P[X = 3] + P[X = 4] = 2k + k = 3(0.15) = 0.45$

(iv) $P[1 \leq X < 4] = P[X = 1] + P[X = 2] + P[X = 3] = k + 2k + 2k = 5k = 5(0.15) = 0.75.$

(v) $F(2) = P[X \leq 2] = P[X = 0] + P[X = 1] + P[X = 2] = 0.1 + k + 2k = 0.1 + 3k = 0.1 + 3(0.15) = 0.1 + 0.45 = 0.55.$

7.3.3 Expected value and Variance of a random variable :

In many problems, it is desirable to describe some feature of the random variable by means of a single number that can be computed from its probability distribution. Few such numbers are mean, median, mode and variance and standard deviation. In this section, we shall discuss mean and variance only. Mean is a measure of location or central tendency in the sense that it roughly locates a **middle** or **average value** of the random variable.



Definition : Let X be a random variable whose possible values $x_1, x_2, x_3, \dots, x_n$ occur with probabilities $p_1, p_2, p_3, \dots, p_n$ respectively. The expected value or arithmetic mean of X , denoted by $E(X)$ or μ is defined by

$$E(X) = \mu = \sum_{i=1}^n x_i p_i = x_1 p_1 + x_2 p_2 + x_3 p_3 + \dots + x_n p_n$$

In other words, the mean or expectation of a random variable X is the sum of the products of all possible values of X by their respective probabilities.

Definition : Let X be a random variable whose possible values $x_1, x_2, x_3, \dots, x_n$ occur with probabilities $p_1, p_2, p_3, \dots, p_n$ respectively. The variance of X , denoted by $Var(X)$ or σ_x^2 is defined as

$$\sigma_x^2 = Var(X) = \sum_{i=1}^n (x_i - \mu)^2 p_i$$

The non-negative number $\sigma_x = \sqrt{Var(X)}$ is called the **standard deviation** of the random variable X .

We can also use the simplified form of

$$Var(X) = \left(\sum_{i=1}^n x_i^2 p_i \right) - \left(\sum_{i=1}^n x_i p_i \right)^2$$

$$Var(X) = E(X^2) - [E(X)]^2 \quad \text{where } \sum_{i=1}^n x_i^2 p_i = E(X^2)$$



SOLVED EXAMPLES

Ex. 1 : Three coins are tossed simultaneously, X is the number of heads. Find expected value and variance of X .

Solution : $S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$ and $X = \{0, 1, 2, 3\}$

$X = x_i$	$P = p_i$	$x_i p_i$	$x_i^2 p_i$
0	$\frac{1}{8}$	0	0
1	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{3}{8}$
2	$\frac{3}{8}$	$\frac{6}{8}$	$\frac{12}{8}$
3	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{9}{8}$
		$\sum_{i=1}^n x_i p_i = \frac{12}{8}$	$\sum_{i=1}^n x_i^2 p_i = \frac{24}{8}$

$$\text{Then } E(X) = \sum_{i=1}^n x_i p_i = \frac{12}{8} = 1.5$$

$$Var(X) = \left(\sum_{i=1}^n x_i^2 p_i \right) - \left(\sum_{i=1}^n x_i p_i \right)^2 = \frac{24}{8} - (1.5)^2 = 3 - 2.25 = 0.75$$

Ex. 2 : Let a pair of dice be thrown and the random variable X be the sum of the numbers that appear on the two dice. Find the mean or expectation of X and variance of X .

Solution : The sample space of the experiment consists of 36 elementary events in the form of ordered pairs (x_i, y_i) , where $x_i = 1, 2, 3, 4, 5, 6$ and $y_i = 1, 2, 3, 4, 5, 6$.

The random variable X i.e. the sum of the numbers on the two dice takes the values 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12.

$X = x_i$	$P = p_i$	$x_i p_i$	$x_i^2 p_i$
2	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{4}{36}$
3	$\frac{2}{36}$	$\frac{6}{36}$	$\frac{18}{36}$
4	$\frac{3}{36}$	$\frac{12}{36}$	$\frac{48}{36}$
5	$\frac{4}{36}$	$\frac{20}{36}$	$\frac{100}{36}$
6	$\frac{5}{36}$	$\frac{30}{36}$	$\frac{180}{36}$
7	$\frac{6}{36}$	$\frac{42}{36}$	$\frac{294}{36}$
8	$\frac{5}{36}$	$\frac{40}{36}$	$\frac{320}{36}$
9	$\frac{4}{36}$	$\frac{36}{36}$	$\frac{324}{36}$
10	$\frac{3}{36}$	$\frac{30}{36}$	$\frac{300}{36}$
11	$\frac{2}{36}$	$\frac{22}{36}$	$\frac{242}{36}$
12	$\frac{1}{36}$	$\frac{12}{36}$	$\frac{144}{36}$
		$\sum_{i=1}^n x_i p_i = \frac{252}{36} = 7$	$\sum_{i=1}^n x_i^2 p_i = \frac{1974}{36} = 54.83$

$$\text{Then } E(X) = \sum_{i=1}^n x_i p_i = 7$$

$$\begin{aligned} \text{Var}(X) &= \left(\sum_{i=1}^n x_i^2 p_i \right) - \left(\sum_{i=1}^n x_i p_i \right)^2 = 54.83 - (7)^2 \\ &= 54.83 - 49 \\ &= 5.83 \end{aligned}$$

Ex. 3 : Find the mean and variance of the number randomly selected from 1 to 15.

Solution : The sample space of the experiment is $S = \{ 1, 2, 3, \dots, 15 \}$.

Let X denote the number selected.

Then X is a random variable which can take values 1, 2, 3, ..., 15. Each number selected is equiprobable therefore

$$P(1) = P(2) = P(3) = \dots = P(15) = \frac{1}{15}$$

$$\begin{aligned} \mu = E(X) &= \sum_{i=1}^n x_i p_i = 1 \times \frac{1}{15} + 2 \times \frac{1}{15} + 3 \times \frac{1}{15} + \dots + 15 \times \frac{1}{15} \\ &= (1 + 2 + 3 + \dots + 15) \times \frac{1}{15} = \left(\frac{15 \times 16}{2} \right) \times \frac{1}{15} = 8 \end{aligned}$$

$$\begin{aligned} Var(X) &= \left(\sum_{i=1}^n x_i^2 p_i \right) - \left(\sum_{i=1}^n x_i p_i \right)^2 = 1^2 \times \frac{1}{15} + 2^2 \times \frac{1}{15} + 3^2 \times \frac{1}{15} + \dots + 15^2 \times \frac{1}{15} - (8)^2 \\ &= (1^2 + 2^2 + 3^2 + \dots + 15^2) \times \frac{1}{15} - (8)^2 \\ &= \left(\frac{15 \times 16 \times 31}{6} \right) \times \frac{1}{15} - (8)^2 \\ &= 82.67 - 64 = 18.67 \end{aligned}$$

Ex. 4 : Two cards are drawn simultaneously (or successively without replacement) from a well shuffled pack of 52 cards. Find the mean, variance and standard deviation of the number of kings drawn.

Solution : Let X denote the number of kings in a draw of two cards. X is a random variable which can assume the values 0, 1 or 2.

$$\text{Then } P(X=0) = P(\text{no card is king}) = \frac{{}^{48}C_2}{{}^{52}C_2} = \frac{48 \times 47}{52 \times 51} = \frac{188}{221}$$

$$\text{Then } P(X=1) = P(\text{exactly one card is king}) = \frac{{}^4C_1 \times {}^{48}C_1}{{}^{52}C_2} = \frac{4 \times 48 \times 27}{52 \times 51} = \frac{32}{221}$$

$$\text{Then } P(X=2) = P(\text{both cards are king}) = \frac{{}^4C_2}{{}^{52}C_2} = \frac{4 \times 3}{52 \times 51} = \frac{1}{221}$$

$$\mu = E(X) = \sum_{i=1}^n x_i p_i = 0 \times \frac{188}{221} + 1 \times \frac{32}{221} + 2 \times \frac{1}{221} = \frac{34}{221}$$

$$\begin{aligned} Var(X) &= \left(\sum_{i=1}^n x_i^2 p_i \right) - \left(\sum_{i=1}^n x_i p_i \right)^2 = \left(0^2 \times \frac{188}{221} + 1^2 \times \frac{32}{221} + 2^2 \times \frac{1}{221} \right) - \left(\frac{34}{221} \right)^2 \\ &= \frac{36}{221} - \frac{1156}{48841} = \frac{6800}{48841} = 0.1392 \end{aligned}$$

$$\sigma = \sqrt{Var(X)} = \sqrt{0.1392}$$



EXERCISE 7.1

- Let X represent the difference between number of heads and number of tails obtained when a coin is tossed 6 times. What are possible values of X ?
- An urn contains 5 red and 2 black balls. Two balls are drawn at random. X denotes number of black balls drawn. What are possible values of X ?
- State which of the following are not the probability mass function of a random variable. Give reasons for your answer.

(i)

X	0	1	2
$P(X)$	0.4	0.4	0.2

(ii)

X	0	1	2	3	4
$P(X)$	0.1	0.5	0.2	-0.1	0.2

(iii)

X	0	1	2
$P(X)$	0.1	0.6	0.3

(iv)

Z	3	2	1	0	-1
$P(Z)$	0.3	0.2	0.4	0	0.05

(v)

Y	-1	0	1
$P(Y)$	0.6	0.1	0.2

(vi)

X	0	-1	-2
$P(X)$	0.3	0.4	0.3

- Find the probability distribution of (i) number of heads in two tosses of a coin. (ii) Number of tails in the simultaneous tosses of three coins. (iii) Number of heads in four tosses of a coin.
- Find the probability distribution of the number of successes in two tosses of a die, where a success is defined as number greater than 4 appears on at least one die.
- From a lot of 30 bulbs which include 6 defectives, a sample of 4 bulbs is drawn at random with replacement. Find the probability distribution of the number of defective bulbs.
- A coin is biased so that the head is 3 times as likely to occur as tail. If the coin is tossed twice, find the probability distribution of number of tails.
- A random variable X has the following probability distribution :

X	0	1	2	3	4	5	6	7
$P(X)$	0	k	$2k$	$2k$	$3k$	k^2	$2k^2$	$7k^2 + k$

Determine : (i) k

(ii) $P(X < 3)$

(iii) $P(X > 4)$

- Find expected value and variance of X for the following p.m.f.

X	-2	-1	0	1	2
$P(X)$	0.2	0.3	0.1	0.15	0.25

10. Find expected value and variance of X , where X is number obtained on uppermost face when a fair die is thrown.
11. Find the mean number of heads in three tosses of a fair coin.
12. Two dice are thrown simultaneously. If X denotes the number of sixes, find the expectation of X .
13. Two numbers are selected at random (without replacement) from the first six positive integers. Let X denote the larger of the two numbers obtained. Find $E(X)$.
14. Let X denote the sum of the numbers obtained when two fair dice are rolled. Find the standard deviation of X .
15. A class has 15 students whose ages are 14, 17, 15, 14, 21, 17, 19, 20, 16, 18, 20, 17, 16, 19 and 20 years. One student is selected in such a manner that each has the same chance of being chosen and the age X of the selected student is recorded. What is the probability distribution of the random variable X ? Find mean, variance and standard deviation of X .
16. In a meeting, 70% of the members favour and 30% oppose a certain proposal. A member is selected at random and we take $X = 0$ if he opposed, and $X = 1$ if he is in favour. Find $E(X)$ and $Var(X)$.

7.4 Probability Distribution of a Continuous Random Variable :

A continuous random variable differs from a discrete random variable in the sense that the possible values of a continuous random variable form an interval of real numbers. In other words, a continuous random variable has uncountably infinite possible values.

For example, the time an athlete takes to complete a thousand-meter race is a continuous random variable.

We shall extend what we learnt about a discrete random variable to a continuous random variable. More specifically, we shall study the probability distribution of a continuous random variable with help of its probability density function (p. d. f.) and its cumulative distribution function (c. d. f.). If the possible values of a continuous random variable X form the interval $[a, b]$, where a and b are real numbers and $a < b$, then the interval $[a, b]$ is called the support of the continuous random variable X . The support of a continuous random variable is often denoted by S .

In case of a discrete random variable X that takes finite or countably infinite distinct values, the probability $P[X = x]$ is determined for every possible value x of the random variable X . The probability distribution of a continuous random variable is not defined in terms of probabilities of possible values of the random variable since the number of possible values are unaccountably infinite. Instead, the probability distribution of a continuous random variable is characterized by probabilities of intervals of the form $[c, d]$, where $c < d$. That is, for a continuous random variable, the interest is in probabilities of the form $P[c < X < d]$, where $a \leq c < d \leq b$.



This probability is obtained by integrating a function of X over the interval $[c, d]$. Let us first define the probability density function (p.d.f.) of a continuous random variable.

7.4.1 Probability Density Function (p. d. f.) :

Let X be a continuous random variable defined on interval $S = (a, b)$. A non-negative integrable function $f(x)$ is called the probability density function (p. d. f.) of X if it satisfies the following conditions.

1. $f(x)$ is positive or zero every where in S , that is, $f(x) \geq 0$, for all $x \in S$.
2. The area under the curve $y = f(x)$ over S is 1. That is, $\int_S f(x) dx = 1$
3. The probability that X takes a value in A , where A is some interval, is given by the integral of $f(x)$ over that interval. That is

$$P [X \in A] = \int_A f(x) dx$$

7.4.2 Cumulative Distribution Functions (c. d. f.) :

The cumulative distribution function for continuous random variables is just a straightforward extension of that of the discrete case. All we need to do is replace the summation with an integral.

Definition : The **cumulative distribution function** (c. d. f.) of a continuous random variable X is defined as :

$$F(x) = \int_a^x f(t) dt \quad \text{for } a < x < b.$$

You might recall, for discrete random variables, that $F(x)$ is, in general, a non-decreasing step function. For continuous random variables, $F(x)$ is a non-decreasing **continuous function**.

SOLVED EXAMPLES

Ex. 1 : Let X be a continuous random variable whose probability density function is $f(x) = 3x^2$, for $0 < x < 1$. note that $f(x)$ is not $P[X = x]$.

For example, $f(0.9) = 3(0.9)^2 = 2.43 > 1$, which is clearly not a probability. In the continuous case, $f(x)$ is the height of the curve at $X = x$, so that the total area under the curve is 1. Here it is areas under the curve that define the probabilities.

Solution : Now, let's start by verifying that $f(x)$ is a valid probability density function.

For this, note the following results.

1. $f(x) = 3x^2 \geq 0$ for all $x \in [0, 1]$.
2. $\int_0^1 f(x) dx = \int_0^1 3x^2 dx = 1$

Therefore, the function $f(x) = 3x^2$, for $0 < x < 1$ is a proper probability density function.

Also, for real numbers c and d such that $0 \leq c < d \leq 1$, note that

$$P [c < X < d] = \int_c^d f(x) dx = \int_c^d 3x^2 dx = [x^3]_c^d = d^3 - c^3 > 0$$

- What is the probability that X falls between $\frac{1}{2}$ and 1? That is, what is $P \left[\frac{1}{2} < X < 1 \right]$?

Substitute $c = \frac{1}{2}$ and $d = 1$ in the above integral to obtain

$$P \left[\frac{1}{2} < X < 1 \right] = 1^3 - \left(\frac{1}{2} \right)^3 = 1 - \frac{1}{8} = \frac{7}{8}.$$

- What is $P \left(X = \frac{1}{2} \right)$?

See that the probability is 0 . This is so because

$$\int_c^d f(x) dx = \int_{\frac{1}{2}}^{\frac{1}{2}} x^3 dx = 1 - 1 = 0.$$

$\left[\text{The ordinate AB, with A} \left(\frac{1}{2}, 0 \right) \text{ and B} \left(\frac{1}{2}, \frac{1}{8} \right) \text{ is degenerate case of rectangle and has area } 0 \right]$

As a matter of fact, in general, if X is a continuous random variable, then the probability that X takes any specific value x is 0. That is, when X is a continuous random variable, then

$$P [X = x] = 0 \text{ for every } x \text{ in the support.}$$

- Ex. 2 :** Let X be a continuous random variable whose probability density function is $f(x) = \frac{x^3}{4}$ for an interval $0 < x < c$. What is the value of the constant c that makes $f(x)$ a valid probability density function?

Solution : Note that the integral of the p. d. f. over the support of the random variable must be

$$\text{That is, } \int_0^c f(x) dx = 1.$$

$$\text{That is, } \int_0^c \left(\frac{x^3}{4} \right) dx = 1. \text{ But, } \int_0^c \left(\frac{x^3}{4} \right) dx = \left[\frac{x^4}{16} \right]_0^c = \frac{c^4}{16}. \text{ Since this integral must be equal to 1,}$$

we have $\frac{c^4}{16} = 1$, or equivalently $c^4 = 16$, so that $c = 2$ since c must be positive.

- Ex. 3 :** Let's return to the example in which X has the following probability density function :

$$f(x) = 3x^2$$

for $0 < x < 1$. What is the cumulative distribution function $F(x)$?

Solution : $F(x) = \int_{-0}^x f(x) dx = \int_0^x 3x^2 dx = [x^3]_0^x = x^3$



Ex. 4 : Let's return to the example in which X has the following probability density function :

$$f(x) = \frac{x^3}{4} \text{ for } 0 < x < 4. \text{ What is the cumulative distribution function } X?$$


Solution : $F(x) = \int_0^x f(x) dx = \int_0^x \frac{x^3}{4} dx = \frac{1}{4} \left[\frac{x^4}{4} \right]_0^x = \frac{1}{16} [x^4 - 0] = \frac{x^4}{16}$

Ex. 5 : Suppose the p.d.f. of a continuous random variable X is defined as:

$$f(x) = x + 1, \text{ for } -1 < x < 0, \text{ and } f(x) = 1 - x, \text{ for } 0 \leq x < 1.$$

Find the c.d.f. $F(x)$.

Solution : If we look the p.d.f. it is defined in two steps



Now for the other two intervals :

For $-1 < x < 0$ and $0 < x < 1$.

$$\begin{aligned} F(x) &= \int_{-1}^x (x + 1) dx \\ &= \left[\frac{x^2}{2} + x \right]_{-1}^x = \left(\frac{x^2}{2} + x \right) - \left(\frac{1}{2} - 1 \right) \\ &= \frac{x^2}{2} + x + \frac{1}{2} = \frac{x^2 + 2x + 2}{2} \\ &= \frac{(x + 1)^2}{2} \end{aligned}$$

$$F(0) = \frac{1}{2}$$

For $0 < x < 1$

$$\begin{aligned} F(x) = P(0 < x < 1) &= P(-1 < x < 0) + P(0 < x < 1) = \int_{-1}^0 (x + 1) dx + \int_0^x (1 - x) dx \\ &= \frac{1}{2} + \int_0^x (1 - x) dx \\ &= \frac{1}{2} + x - \frac{x^2}{2} \end{aligned}$$

$$F(x) = 0, \text{ for } x \leq -1$$

$$\begin{aligned} F(x) &= \frac{1}{2} (x + 1)^2, \text{ for } -1 < x \leq 0 \\ &= \frac{1}{2} + x - \frac{x^2}{2}, \text{ for } 0 < x < 1 \end{aligned}$$

If probability function $f(x)$ is defined on (a, b) with $f(x) \geq 0$ and $\int_a^b f(x) dx = 1$, then we can extend this function to the whole of \mathbb{R} as follows.

For $x \leq a$ and $x \geq b$, define $f(x) = 0$.

Then note that $\int_{-\infty}^t f(x) dx = 0$, for $t \leq a$ and for $x \geq b$

$$\int_{-\infty}^t f(x) dx = \int_{-\infty}^a f(x) dx + \int_a^b f(x) dx + \int_b^t f(x) dx = 0 + 1 + 0$$

Thus $F(t) = 0$, for $t \leq a$ and $F(t) = 1$, for $t \geq b$

Ex. 6 : Verify if the following functions are p.d.f. of a continuous r.v. X .

(i) $f(x) = e^{-x}$, for $0 < x < \infty$ and $= 0$, otherwise.

(ii) $f(x) = \frac{x}{2}$, for $-2 < x < 2$ and $= 0$, otherwise.

Solution : (i) e^{-x} is ≥ 0 for any value of x since $e > 0$,

$$\therefore e^{-x} > 0, \text{ for } 0 < x < \infty$$

$$\int_0^{\infty} f(x) dx = \int_0^{\infty} e^{-x} dx = \left[-e^{-x} \right]_0^{\infty} = \left[\frac{1}{e^{\infty}} - e^0 \right] = -(0 - 1) = 1$$

Both the conditions of p.d.f. are satisfied $f(x)$ is p.d.f. of r.v.

(ii) $f(x) < 0$ i.e. negative for $-2 < x < 0$ therefore $f(x)$ is not p.d.f.

Ex. 7 : Find k if the following function is the p.d.f. of r.v. X .

$$f(x) = kx^2(1-x), \quad \text{for } 0 < x < 1 \text{ and } = 0, \text{ otherwise.}$$

Solution : Since $f(x)$ is the p.d.f. of r.v. X

$$\int_0^1 kx^2(1-x) dx = 1$$

$$\therefore \int_0^1 k(x^2 - x^3) dx = 1$$

$$\therefore k \left[\frac{x^3}{3} - \frac{x^4}{4} \right]_0^1 = 1$$

$$\therefore k \left\{ \left[\frac{1}{3} - \frac{1}{4} \right] - (0) \right\} = 1$$

$$\therefore k \times \frac{1}{12} = 1 \qquad \therefore k = 12$$



Ex. 8 : For each of the following p.d.f. of r.v. X , find (a) $P(X < 1)$ and (b) $P(|X| < 1)$

(i) $f(x) = \frac{x^2}{18}$, for $-3 < x < 3$ and $= 0$, otherwise.

(ii) $f(x) = \frac{x+2}{18}$, for $-2 < x < 4$ and $= 0$, otherwise.

Solution :

(i) (a) $P(X < 1) = \int_{-3}^1 \frac{x^2}{18} dx = \left[\frac{\frac{1}{18}(x^3)}{3} \right]_{-3}^1 = \frac{1}{54} [1 - (-3)^3] = \frac{1}{54} (1 + 27) = \frac{28}{54} = \frac{14}{27}$

(b) $P(|X| < 1) = P(-1 < x < 1) = \int_{-1}^1 \frac{x^2}{18} dx = \left[\frac{\frac{1}{18}(x^3)}{3} \right]_{-1}^1$
 $= \frac{1}{54} [1 - (-1)^3] = \frac{1}{54} (1 + 1) = \frac{2}{54} = \frac{1}{27}$

(ii) (a) $P(X < 1) = \int_{-2}^1 \frac{x+2}{18} dx = \frac{1}{18} \left[\frac{x^2}{2} + 2x \right]_{-2}^1$
 $= \frac{1}{18} \left\{ \left(\frac{1}{2} + 2 \right) - \left(\frac{(-2)^2}{2} + 2(-2) \right) \right\} = \frac{1}{18} \left\{ \frac{5}{2} + 2 \right\} = \frac{1}{18} \times \frac{9}{2} = \frac{1}{4}$

(b) $P(|X| < 1) = P(-1 < x < 1) = \int_{-1}^1 \frac{x+2}{18} dx = \frac{1}{18} \left[\frac{x^2}{2} + 2x \right]_{-1}^1$
 $= \frac{1}{18} \left\{ \left(\frac{1}{2} + 2 \right) - \left(\frac{1}{2} - 2 \right) \right\} = \frac{1}{18} \left\{ \frac{5}{2} + \frac{3}{2} \right\} = \frac{1}{18} \times 4 = \frac{2}{9}$

Ex. 9 : Find the c.d.f. $F(x)$ associated with p.d.f. $f(x)$ of r.v. X where

$f(x) = 3(1 - 2x^2)$ for $0 < x < 1$ and $= 0$, otherwise.

Solution : Since $f(x)$ is p.d.f. of r.v. therefore c.d.f. is

$$F(x) = \int_0^x 3(1 - 2x^2) dx = \left[3 \left(x - \frac{2x^3}{3} \right) \right]_0^x = [3x - 2x^3] = 3x - 2x^3$$

EXERCISE 7.2

1. Verify which of the following is p.d.f. of r.v. X :

(i) $f(x) = \sin x$, for $0 \leq x \leq \frac{\pi}{2}$ (ii) $f(x) = x$, for $0 \leq x \leq 1$ and $= 2 - x$ for $1 < x < 2$

(iii) $f(x) = 2$, for $0 \leq x \leq 1$

2. The following is the p.d.f. of r.v. $X : f(x) = \frac{x}{8}$, for $0 < x < 4$ and $= 0$ otherwise
Find (a) $P(x < 1.5)$ (b) $P(1 < x < 2)$ (c) $P(x > 2)$
3. It is known that error in measurement of reaction temperature (in 0°C) in a certain experiment is continuous r.v. given by
 $f(x) = \frac{x^2}{3}$, for $-1 < x < 2$ and $= 0$ otherwise
(i) Verify whether $f(x)$ is p.d.f. of r.v. X . (ii) Find $P(0 < x \leq 1)$
(iii) Find probability that X is negative.
4. Find k if the following function represent p.d.f. of r.v. X .
(i) $f(x) = kx$, for $0 < x < 2$ and $= 0$ otherwise, Also find $P\left(\frac{1}{4} < x < \frac{3}{2}\right)$.
(ii) $f(x) = kx(1-x)$, for $0 < x < 1$ and $= 0$ otherwise, Also find $P\left(\frac{1}{4} < x < \frac{1}{2}\right)$, $P\left(x < \frac{1}{2}\right)$.
5. Let X be amount of time for which a book is taken out of library by randomly selected student and suppose X has p.d.f.
 $f(x) = 0.5x$, for $0 \leq x \leq 2$ and $= 0$ otherwise.
Calculate: (i) $P(X \leq 1)$ (ii) $P(0.5 \leq x \leq 1.5)$ (iii) $P(x \geq 1.5)$
6. Suppose that X is waiting time in minutes for a bus and its p.d.f. is given by
 $f(x) = \frac{1}{5}$, for $0 \leq x \leq 5$ and $= 0$ otherwise.
Find the probability that (i) waiting time is between 1 and 3
(ii) waiting time is more than 4 minutes.
7. Suppose error involved in making a certain measurement is continuous r.v. X with p.d.f.
 $f(x) = k(4-x^2)$, for $-2 \leq x \leq 2$ and $= 0$ otherwise.
Compute : (i) $P(x > 0)$ (ii) $P(-1 < x < 1)$ (iii) $P(-0.5 < x \text{ or } x > 0.5)$
8. The following is the p.d.f. of continuous r.v.
 $f(x) = \frac{x}{8}$, for $0 < x < 4$ and $= 0$ otherwise.
(i) Find expression for c.d.f. of X (ii) Find $F(x)$ at $x = 0.5, 1.7$ and 5 .
9. Given the p.d.f. of a continuous r.v. $X, f(x) = \frac{x^2}{3}$, for $-1 < x < 2$ and $= 0$ otherwise
Determine c.d.f. of X hence find $P(x < 1), P(x < -2), P(X > 0), P(1 < x < 2)$
10. If a r.v. X has p.d.f.,
 $f(x) = \frac{c}{x}$, for $1 < x < 3, c > 0$, Find $c, E(X)$ and $\text{Var}(X)$.



Let us Remember

- ✿ A random variable (r.v.) is a real-valued function defined on the sample space of a random experiment.

The domain of a random variable is the sample space of a random experiment, while its co-domain is the real line.

Thus $X: S \rightarrow R$ is a random variable.

There are two types of random variables, namely discrete and continuous.

- ✿ **Discrete random variable :** Let the possible values of discrete random variable X be denoted by x_1, x_2, x_3, \dots , and the corresponding probabilities be denoted by p_1, p_2, p_3, \dots , where $p_i = P[X = x_i]$ for $i = 1, 2, 3, \dots$. If there is a function f such that $p_i = P[X = x_i] = f(x_i)$ for all possible values of X , then f is called the probability mass function (p. m. f.) of X .

Note : If x_i is a possible value of X and $p_i = P[X = x_i]$, then there is an event E_i in the sample space S such that $p_i = P[E_i]$. Since x_i is a possible value of X , $p_i = P[X = x_i] > 0$. Also, all possible values of X cover all sample points in the sample space S , and hence the sum of their probabilities is 1. That is, $p_i > 0$ for all i and $\sum p_i = 1$.

- ✿ **c.d.f ($F(x)$) :** The cumulative distribution function (c. d. f.) of a discrete random variable X is denoted by F and is defined as follows.

$$\begin{aligned} F(x) = P[X \leq x] &= \sum_{x_i < x} P[X = x_i] \\ &= \sum_{x_i < x} p_i \\ &= \sum_{x_i < x} f(x_i) \end{aligned}$$

- ✿ **Expected Value or Mean of Discrete r. v. :** Let X be a random variable whose possible values $x_1, x_2, x_3, \dots, x_n$ occur with probabilities $p_1, p_2, p_3, \dots, p_n$ respectively. The expected value or arithmetic mean of X , denoted by $E(X)$ or μ is defined by

$$E(X) = \mu = \sum_{i=1}^n x_i p_i = x_1 p_1 + x_2 p_2 + x_3 p_3 + \dots + x_n p_n$$

In other words, the mean or expectation of a random variable X is the sum of the products of all possible values of X by their respective probabilities.

✿ **Variance of Discrete r. v. :** Let X be a random variable whose possible values $x_1, x_2, x_3, \dots, x_n$ occur with probabilities $p_1, p_2, p_3, \dots, p_n$ respectively. The variance of X , denoted by $Var(X)$ or σ_x^2 is defined as

$$\sigma_x^2 = Var(X) = \sum_{i=1}^n (x_i - \mu)^2 p_i$$

The non-negative number $\sigma_x = \sqrt{Var(X)}$

is called the **standard deviation** of the random variable X .

Another formula to find the variance of a random variable. We can also use the simplified form of

$$Var(X) = \left(\sum_{i=1}^n x_i^2 p_i \right) - \left(\sum_{i=1}^n x_i p_i \right)^2$$

$$Var(X) = E(X^2) - [E(X)]^2 \quad \text{where } \sum_{i=1}^n x_i^2 p_i = E(X^2)$$

✿ **Probability Density Function (p. d. f.) :** Let X be a continuous random variable defined on interval $S = (a, b)$. A non-negative integrable function $f(x)$ is called the probability density function (p. d. f.) of X if it satisfies the following conditions.

1. $f(x)$ is positive every where in S , that is, $f(x) > 0$, for all $x \in S$.
2. The area under the curve $f(x)$ over S is 1. That is, $\int_S f(x) dx = 1$
3. The probability that X takes a value in A , where A is some interval, is given by the integral of $f(x)$ over that interval. That is

$$P[X \in A] = \int_A f(x) dx$$

✿ The **cumulative distribution function** (c. d. f.) of a continuous random variable X is defined as :

$$F(x) = \int_a^x f(t) dt \quad \text{for } a < x < b.$$

MISCELLANEOUS EXERCISE 7

(I) Choose the correct option from the given alternatives :

(1) P.d.f. of a.c.r.v X is $f(x) = 6x(1-x)$, for $0 \leq x \leq 1$ and $= 0$, otherwise (elsewhere)

If $P(X < a) = P(X > a)$, then $a =$

- (A) 1 (B) $\frac{1}{2}$ (C) $\frac{1}{3}$ (D) $\frac{1}{4}$

- (2) If the p.d.f of a.c.r.v. X is $f(x) = 3(1 - 2x^2)$, for $0 < x < 1$ and $= 0$, otherwise (elsewhere) then the c.d.f of X is $F(x) =$
- (A) $2x - 3x^2$ (B) $3x - 4x^3$ (C) $3x - 2x^3$ (D) $2x^3 - 3x$
- (3) If the p.d.f of a.c.r.v. X is $f(x) = \frac{x^2}{18}$, for $-3 < x < 3$ and $= 0$, otherwise then $P(|X| < 1) =$
- (A) $\frac{1}{27}$ (B) $\frac{1}{28}$ (C) $\frac{1}{29}$ (D) $\frac{1}{26}$
- (4) If a d.r.v. X takes values $0, 1, 2, 3, \dots$ which probability $P(X=x) = k(x+1) \cdot 5^{-x}$, where k is a constant, then $P(X=0) =$
- (A) $\frac{7}{25}$ (B) $\frac{16}{25}$ (C) $\frac{18}{25}$ (D) $\frac{19}{25}$
- (5) If p.m.f. of a d.r.v. X is $P(X=x) = \frac{{}^5C_x}{2^5}$, for $x = 0, 1, 2, 3, 4, 5$ and $= 0$, otherwise If $a = P(X \leq 2)$ and $b = P(X \geq 3)$, then $E(X) =$
- (A) $a < b$ (B) $a > b$ (C) $a = b$ (D) $a + b$
- (6) If p.m.f. of a d.r.v. X is $P(X=x) = \frac{x}{n(n+1)}$, for $x = 1, 2, 3, \dots, n$ and $= 0$, otherwise then $E(X) =$
- (A) $\frac{n}{1} + \frac{1}{2}$ (B) $\frac{n}{3} + \frac{1}{6}$ (C) $\frac{n}{2} + \frac{1}{5}$ (D) $\frac{n}{1} + \frac{1}{3}$
- (7) If p.m.f. of a d.r.v. X is $P(x) = \frac{c}{x^3}$, for $x = 1, 2, 3$ and $= 0$, otherwise (elsewhere) then $E(X) =$
- (A) $\frac{343}{297}$ (B) $\frac{294}{251}$ (C) $\frac{297}{294}$ (D) $\frac{294}{297}$
- (8) If the a d.r.v. X has the following probability distribution :
- | | | | | | | |
|----------|-----|-----|-----|------|-----|-----|
| X | -2 | -1 | 0 | 1 | 2 | 3 |
| $P(X=x)$ | 0.1 | k | 0.2 | $2k$ | 0.3 | k |
- then $P(X = -1) =$
- (A) $\frac{1}{10}$ (B) $\frac{2}{10}$ (C) $\frac{3}{10}$ (D) $\frac{4}{10}$
- (9) If the a d.r.v. X has the following probability distribution :
- | | | | | | | | |
|----------|-----|------|------|------|-------|--------|------------|
| X | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| $P(X=x)$ | k | $2k$ | $2k$ | $3k$ | k^2 | $2k^2$ | $7k^2 + k$ |
- then $k =$
- (A) $\frac{1}{7}$ (B) $\frac{1}{8}$ (C) $\frac{1}{9}$ (D) $\frac{1}{10}$

- (10) Find expected value of and variance of
- X
- for the following p.m.f.

X	-2	-1	0	1	2
$P(X)$	0.3	0.4	0.2	0.15	0.25

- (A) 0.85 (B) -0.85 (C) 0.15 (D) -0.15

(II) Solve the following :

- (1) Identify the random variable as either discrete or continuous in each of the following. Write down the range of it.
- (i) An economist is interested the number of unemployed graduate in the town of population 1 lakh.
- (ii) Amount of syrup prescribed by physician.
- (iii) The person on the high protein diet is interested gain of weight in a week.
- (iv) 20 white rats are available for an experiment. Twelve rats are male. Scientist randomly selects 5 rats number of female rats selected on a specific day.
- (v) A highway safety group is interested in studying the speed (km/hrs) of a car at a check point.

- (2) The probability distribution of discrete r.v.
- X
- is as follows

$X = x$	1	2	3	4	5	6
$P(X = x)$	k	$2k$	$3k$	$4k$	$5k$	$6k$

- (i) Determine the value of
- k
- . (ii) Find
- $P(X \leq 4)$
- ,
- $P(2 < X < 4)$
- ,
- $P(X \geq 3)$
- .

- (3) The following probability distribution of r.v.
- X

$X = x$	-3	-2	-1	0	1	2	3
$P(X = x)$	0.05	0.1	0.15	0.20	0.25	0.15	0.1

Find the probability that

- (i)
- X
- is positive. (ii)
- X
- is non negative. (iii)
- X
- is odd. (iv)
- X
- is even.

- (4) The p.m.f. of a r.v.
- X
- is given by
- $P(X = x) = \frac{{}^5C_x}{2^5}$
- , for
- $x = 0, 1, 2, 3, 4, 5$
- and
- $= 0$
- , otherwise. Then show that
- $P(X \leq 2) = P(X \geq 3)$
- .

- (5) In the p.m.f. of r.v.
- X

x	1	2	3	4	5
$P(X)$	$\frac{1}{20}$	$\frac{3}{20}$	a	$2a$	$\frac{1}{20}$

Find a and obtain c.d.f. of X .

- (6) A fair coin is tossed 4 times. Let
- X
- denotes the number of heads obtained write down the probability distribution of
- X
- . Also find the formula for p.m.f. of
- X
- .

- (7) Find the probability distribution of the number of successes in two tosses of a die, where a success is defined as (i) number greater than 4 (ii) six appears on at least one die.
- (8) A random variable X has the following probability distribution.

X	0	1	2	3	4	5	6	7
$P(X)$	0	k	$2k$	$2k$	$3k$	k^2	$2k^2$	$7k^2 + k$

Determine (i) k (ii) $P(X > 6)$ (iii) $P(0 < X < 3)$

- (9) The following is the c.d.f. of r.v. X

X	-3	-2	-1	0	1	2	3	4
$F(X)$	0.1	0.3	0.5	0.65	0.75	0.85	0.9	1

Find p.m.f. of X . (i) $P(-1 \leq X \leq 2)$ (ii) $P(X \leq 3 / X > 0)$

- (10) Find the expected value, variance and standard deviation of r.v. X whose p.m.f. are given below.

(i)

$X = x$	1	2	3
$P(X = x)$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{2}{5}$

(ii)

$X = x$	-1	0	1
$P(X = x)$	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{2}{5}$

(iii)

$X = x$	1	2	3	...	n
$P(X = x)$	$\frac{1}{n}$	$\frac{1}{n}$	$\frac{1}{n}$...	$\frac{1}{n}$

(iv)

$X = x$	0	1	2	3	4	5
$P(X = x)$	$\frac{1}{32}$	$\frac{5}{32}$	$\frac{10}{32}$	$\frac{10}{32}$	$\frac{5}{32}$	$\frac{1}{32}$

- (11) A player tosses two coins he wins ₹ 10 if 2 heads appears, ₹ 5 if 1 head appears and ₹ 2 if no head appears. Find the expected winning amount and variance of winning amount.

- (12) Let the p.m.f. of r.v. X be $P(x) = \frac{3-x}{10}$, for $x = -1, 0, 1, 2$ and $= 0$, otherwise. Calculate $E(X)$ and $Var(X)$.

- (13) Suppose error involved in making a certain measurement is continuous r.v. X with p.d.f. $f(x) = k(4 - x^2)$, for $-2 \leq x \leq 2$ and $= 0$ otherwise. Compute (i) $P(X > 0)$ (ii) $P(-1 < x < 1)$ (iii) $P(X < 0.5 \text{ or } X > 0.5)$

- (14) The p.d.f. of r.v. X is given by $f(x) = \frac{1}{2a}$, for $0 < x < 2a$ and $= 0$, otherwise. Show that $P\left(X < \frac{a}{2}\right) = P\left(X > \frac{3a}{2}\right)$.

- (15) The p.d.f. of r.v. of X is given by $f(x) = \frac{k}{\sqrt{x}}$, for $0 < x < 4$ and $= 0$, otherwise. Determine k . Determine c.d.f. of X and hence $P(X \leq 2)$ and $P(X \leq 1)$.



8. BINOMIAL DISTRIBUTION



Let us Study

- Bernoulli Trial
- Binomial distribution
- Mean and variance of Binomial Distribution.



Let us Recall

- Many experiments are dichotomous in nature. For example, a tossed coin shows a 'head' or 'tail', A result of student 'pass' or 'fail', a manufactured item can be 'defective' or 'non-defective', the response to a question might be 'yes' or 'no', an egg has 'hatched' or 'not hatched', the decision is 'yes' or 'no' etc. In such cases, it is customary to call one of the outcomes a 'success' and the other 'not success' or 'failure'. For example, in tossing a coin, if the occurrence of the head is considered a success, then occurrence of tail is a failure.



Let us Learn

8.1.1 Bernoulli Trial :

Each time we toss a coin or roll a die or perform any other experiment, we call it a trial. If a coin is tossed, say, 4 times, the number of trials is 4, each having exactly two outcomes, namely, success or failure. The outcome of any trial is independent of the outcome of any other trial. In each of such trials, the probability of success or failure remains constant. Such independent trials which have only two outcomes usually referred to as 'success' or 'failure' are called Bernoulli trials.

Definition:

Trials of a random experiment are called Bernoulli trials, if they satisfy the following conditions :

- Each trial has exactly two outcomes : success or failure.
- The probability of success remains the same in each trial.

Throwing a die 50 times is a case of 50 Bernoulli trials, in which each trial results in success (say an even number) or failure (an odd number) and the probability of success (p) is same for all 50 throws. Obviously, the successive throws of the die are independent trials. If the die is fair and has six numbers 1 to 6 written on six faces, then

$$p = \frac{1}{2} \text{ and } q = 1 - p \quad \therefore q = \frac{1}{2}$$

For example :

Consider a die to be thrown 20 times. if the result is an even number, consider it a success, else it is a failure. Then $p = \frac{1}{2}$ as there are 3 even numbers in the possible outcomes.

If in the same experiment, we consider the result a success if it is a multiple of 3, then $p = \frac{1}{3}$ as there are 2 multiples of 3 among the six possible outcomes. Both above trials are Bernoulli trials.

**SOLVED EXAMPLE**

Ex. 1 : Six balls are drawn successively from an urn containing 7 red and 9 black balls. Tell whether or not the trials of drawing balls are Bernoulli trials when after each draw the ball drawn is

- (i) replaced (ii) not replaced in the urn.

Solution :

- (i) The number of trials is finite. When the drawing is done with replacement, the probability of success (say, red ball) is $p = \frac{7}{16}$ which is same for all six trials (draws). Hence, the drawing of balls with replacements are Bernoulli trials.
- (ii) When the drawing is done without replacement, the probability of success (i.e. red ball) in first trial is $\frac{7}{16}$ in second trial is $\frac{6}{15}$ if first ball drawn is red and is $\frac{7}{15}$ if first ball drawn is black and so on. Clearly probability of success is not same for all trials, hence the trials are not Bernoulli trials.

8.2 Binomial distribution:

Consider the experiment of tossing a coin in which each trial results in success (say, heads) or failure (tails). Let S and F denote respectively success and failure in each trial. Suppose we are interested in finding the ways in which we have one success in six trials. Clearly, six different cases are there as listed below:

SFFFFF, FSFFFF, FFSFFF, FFFSFF, FFFFSF, FFFFFS.

Similarly, two successes and four failures can have $\frac{6!}{4! \times 2!} = 15$ combinations.

But as n grows large, the calculation can be lengthy. To avoid this the number for certain probabilities can be obtained with Bernoulli's formula. For this purpose, let us take the experiment made up of three Bernoulli trials with probabilities p and $q = 1 - p$ for success and failure respectively in each trial. The sample space of the experiment is the set

$$S = \{SSS, SSF, SFS, FSS, SFF, FSF, FFS, FFF\}$$



The number of successes is a random variable X and can take values 0, 1, 2, or 3. The probability distribution of the number of successes is as below :

$$\begin{aligned} P(X=0) &= P(\text{no success}) \\ &= P(\{\text{FFF}\}) = P(F) \cdot P(F) \cdot P(F), \text{ since trials are independent.} \\ &= q \cdot q \cdot q = q^3 \end{aligned}$$

$$\begin{aligned} P(X=1) &= P(\text{one success}) \\ &= P(\{\text{SFF, FSF, FFS}\}) \\ &= P(\{\text{SFF}\}) + P(\{\text{FSF}\}) + P(\{\text{FFS}\}) \\ &= P(S) \cdot P(F) \cdot P(F) + P(F) \cdot P(S) \cdot P(F) + P(F) \cdot P(F) \cdot P(S) \\ &= p \cdot q \cdot q + q \cdot p \cdot q + q \cdot q \cdot p = 3pq^2 \end{aligned}$$

$$\begin{aligned} P(X=2) &= P(\text{two success}) \\ &= P(\{\text{SSF, SFS, FSS}\}) \\ &= P(\{\text{SSF}\}) + P(\{\text{SFS}\}) + P(\{\text{FSS}\}) \\ &= P(S) \cdot P(S) \cdot P(F) + P(S) \cdot P(F) \cdot P(S) + P(F) \cdot P(S) \cdot P(S) \\ &= p \cdot p \cdot q + p \cdot q \cdot p + q \cdot p \cdot p = 3p^2q \end{aligned}$$

and

$$\begin{aligned} P(X=3) &= P(\text{three successes}) \\ &= P(\{\text{SSS}\}) \\ &= P(S) \cdot P(S) \cdot P(S) \\ &= p^3 \end{aligned}$$

Thus, the probability distribution of X is

X	0	1	2	3
$P(X)$	q^3	$3q^2p$	$3qp^2$	p^3

Also, the binominal expansion of

$$(q + p)^3 \text{ is } q^3 + 3q^2p + 3qp^2 + p^3$$

Note that the probabilities of 0, 1, 2 or 3 successes are respectively the 1st, 2nd, 3rd and 4th term in the expansion of $(q + p)^3$.

Also, since $q + p = 1$, it follows that the sum of these probabilities, as expected, is 1. Thus, we may conclude that in an experiment of n -Bernoulli trials, the probabilities of 0, 1, 2, ..., n successes can be



obtained as 1st, 2nd, 3rd, . . . , (n + 1)th terms in the expansion of (q + p)ⁿ. To prove this assertion (result), let us find the probability of x successes in an experiment of n-Bernoulli trials.

Clearly, in case of x successes (S), there will be (n - x) failures (F). Now x successes (S) and (n - x) failures (F) can be obtained in $\frac{n!}{x!(n-x)!}$ ways.

In each of these ways the probability of x successes and (n - x) failures

$$\begin{aligned} &= P(x \text{ successes}) \cdot P((n-x) \text{ failures}) \\ &= (P(S) \cdot P(S) \dots P(S) \text{ } x \text{ times}) \cdot (P(F) \cdot P(F)) \dots (P(F) \cdot (n-x) \text{ times}) \\ &= (p \cdot p \cdot p \dots p \text{ } x \text{ times}) (q \cdot q \cdot q \dots q \text{ } (n-x) \text{ times}) \\ &= p^x \cdot q^{n-x} \end{aligned}$$

Thus probability of getting x successes in n-Bernoulli trial is

$$P(x \text{ successes out of } n \text{ trials}) = \frac{n!}{x!(n-x)!} \times p^x \times q^{n-x} = {}^n C_x p^x \times q^{n-x}$$

Clearly, P(x successes), i.e. ${}^n C_x p^x q^{n-x}$ is the (x + 1)th term in the binomial expansion of (q + p)ⁿ.

Thus, the probability distribution of number of successes in an experiment consisting of n-Bernoulli trials may be obtained by the binomial expansion of (q + p)ⁿ. Hence, this distribution of number of successes X can be written as

X	0	1	2	...	x	...	n
P(X)	${}^n C_0 p^0 \times q^n$	${}^n C_1 p^1 \times q^{n-1}$	${}^n C_2 p^2 \times q^{n-2}$...	${}^n C_x p^x \times q^{n-x}$...	${}^n C_n p^n \times q^0$

The above probability distribution is known as binomial distribution with parameters n and p, because for given values of n and p, we can find the complete probability distribution. It is represented $X \sim B(n, p)$ as read as X follows binomial distribution with parameters n, p

The probability of x successes P(X = x) is also denoted by P(x) and is given by

$$P(x) = {}^n C_x \cdot q^{n-x} \times p^x, x = 0, 1, \dots, n, (q = 1 - p)$$

This P(x) is called the **probability function** of the binomial distribution.

A binomial distribution with n-Bernoulli trials and probability of success in each trial as p, is denoted by B(n, p) or $X \sim B(n, p)$.

- Lets Note :**
- (i) The number of trials should be fixed.
 - (ii) The trials should be independent.





SOLVED EXAMPLES

Ex. 1 : If a fair coin is tossed 10 times, find the probability of getting

(i) exactly six heads

(ii) at least six heads

(iii) at most six heads

Solution : The repeated tosses of a coin are Bernoulli trials. Let X denote the number of heads in an experiment of 10 trials.

Clearly, $X \sim B(n, p)$ with $n = 10$ and $p = \frac{1}{2}$, $q = 1 - p = 1 - \frac{1}{2} \quad \therefore q = \frac{1}{2}$

$$P(X = x) = {}^n C_x p^x \times q^{n-x}$$

$$= {}^{10} C_x \left(\frac{1}{2}\right)^x \times \left(\frac{1}{2}\right)^{n-x}$$

(i) Exactly six successes means $x = 6$

$$\begin{aligned} P(X = 6) &= {}^{10} C_6 \left(\frac{1}{2}\right)^6 \times \left(\frac{1}{2}\right)^{10-6} = \frac{10!}{6!(10-6)!} \times \left(\frac{1}{2}\right)^6 \times \left(\frac{1}{2}\right)^4 = \frac{10 \times 9 \times 8 \times 7}{4 \times 3 \times 2 \times 1} \times \left(\frac{1}{2}\right)^{10} \\ &= \frac{105}{512} \end{aligned}$$

(ii) At least six successes means $x \geq 6$

$$\begin{aligned} P(X \geq 6) &= [P(X = 6) + P(X = 7) + P(X = 8) + P(X = 9) + P(X = 10)] \\ &= {}^{10} C_6 \left(\frac{1}{2}\right)^6 \times \left(\frac{1}{2}\right)^4 + {}^{10} C_7 \left(\frac{1}{2}\right)^7 \times \left(\frac{1}{2}\right)^3 + {}^{10} C_8 \left(\frac{1}{2}\right)^8 \times \left(\frac{1}{2}\right)^2 + {}^{10} C_9 \left(\frac{1}{2}\right)^9 \times \left(\frac{1}{2}\right)^1 + {}^{10} C_{10} \left(\frac{1}{2}\right)^{10} \times \left(\frac{1}{2}\right)^0 \\ &= \frac{10 \times 9 \times 8 \times 7}{4 \times 3 \times 2 \times 1} \times \left(\frac{1}{2}\right)^{10} + \frac{10 \times 9 \times 8}{3 \times 2 \times 1} \times \left(\frac{1}{2}\right)^{10} + \frac{10 \times 9}{2 \times 1} \times \left(\frac{1}{2}\right)^{10} + 10 \left(\frac{1}{2}\right)^{10} + \left(\frac{1}{2}\right)^{10} \\ &= (210 + 120 + 45 + 10 + 1) \times \frac{1}{1024} \\ &= \frac{386}{1024} = \frac{193}{512} \end{aligned}$$

(iii) At most six successes means $x \leq 6$

$$\begin{aligned} P(X \leq 6) &= 1 - (P(X > 6)) \\ &= 1 - [P(X = 7) + P(X = 8) + P(X = 9) + P(X = 10)] \\ &= 1 - \left[{}^{10} C_7 \left(\frac{1}{2}\right)^7 \times \left(\frac{1}{2}\right)^3 + {}^{10} C_8 \left(\frac{1}{2}\right)^8 \times \left(\frac{1}{2}\right)^2 + {}^{10} C_9 \left(\frac{1}{2}\right)^9 \times \left(\frac{1}{2}\right)^1 + {}^{10} C_{10} \left(\frac{1}{2}\right)^{10} \times \left(\frac{1}{2}\right)^0 \right] \\ &= 1 - \left[\frac{10 \times 9 \times 8}{3 \times 2 \times 1} \times \left(\frac{1}{2}\right)^{10} + \frac{10 \times 9}{2 \times 1} \times \left(\frac{1}{2}\right)^{10} + 10 \left(\frac{1}{2}\right)^{10} \times \left(\frac{1}{2}\right)^0 \right] \\ &= 1 - \left[(120 + 45 + 10 + 1) \times \frac{1}{1024} \right] = 1 - \frac{176}{1024} = 1 - \frac{88}{512} = \frac{512 - 88}{512} = \frac{424}{512} = \frac{53}{64} \end{aligned}$$

Ex. 2 : Ten eggs are drawn successively with replacement from a lot containing 10% defective eggs. Find the probability that there is at least one defective egg.

Solution : Let X denote the number of defective eggs in the 10 eggs drawn.

Since the drawing is done with replacement, the trials are Bernoulli trials.

$$\text{Probability of success} = \frac{1}{10}$$

$$p = \frac{1}{10}, \quad q = 1 - p = 1 - \frac{1}{10} \quad \therefore q = \frac{9}{10}$$

$$n = 10$$

$$X \sim B \left(10, \frac{1}{10} \right)$$

$$P(X=x) = {}^{10}C_x \left(\frac{1}{10} \right)^x \times \left(\frac{9}{10} \right)^{10-x}$$

Here $X \geq 1$

$$\begin{aligned} P(X \geq 1) &= 1 - {}^{10}C_0 \left(\frac{1}{10} \right)^0 \times \left(\frac{9}{10} \right)^{10} \\ &= 1 - 1 \times 1 \times \left(\frac{9}{10} \right)^{10} \\ &= 1 - \left(\frac{9}{10} \right)^{10} \end{aligned}$$

8.3 Mean and Variance of Binomial Distribution (Formulae without proof) :

Let $X \sim B(n, p)$ then mean or expected value of r.v. X is denoted by μ or $E(X)$ and given by $\mu = E(X) = np$.

The variance is denoted by $Var(X)$ and given by $Var(X) = npq$.

Standard deviation of X is denoted by $SD(X)$ or σ and given by $SD(X) = \sigma_x = \sqrt{Var(X)}$

For example : If $X \sim B(10, 0.4)$ then find $E(X)$ and $Var(X)$.

Solution : Here $n = 10, p = 0.4,$

$$q = 1 - p$$

$$q = 1 - 0.4 = 0.6$$

$$E(X) = np$$

$$= 10 \times 0.4 = 4$$

$$Var(X) = npq$$

$$= 10 \times 0.4 \times 0.6$$

$$= 2.4$$





SOLVED EXAMPLES

Ex. 1 : Let the p.m.f. of r.v. X be

$$P(X=x) = {}^4C_x \left(\frac{5}{9}\right)^x \times \left(\frac{4}{9}\right)^{4-x}, \text{ for } x = 0, 1, 2, 3, 4.$$

then find $E(X)$ and $Var(X)$.

Solution : $P(X=x)$ is binomial distribution with $n = 4$, $p = \frac{5}{9}$ and $q = \frac{4}{9}$

$$E(X) = np$$

$$= 4 \times \left(\frac{5}{9}\right) = \frac{20}{9}$$

$$Var(X) = npq$$

$$= 4 \times \left(\frac{5}{9}\right) \times \left(\frac{4}{9}\right) = \frac{80}{81}$$

Ex. 2 : If $E(X) = 6$ and $Var(X) = 4.2$, find n and p .

Solution : $E(X) = 6$ therefore $np = 6$ and $Var(X) = 4.2$ therefore $npq = 4.2$

$$\frac{npq}{np} = \frac{4.2}{6} \quad \therefore q = 0.7$$

$$\therefore p = 1 - q = 1 - 0.7 \quad \therefore p = 0.3$$

$$np = 6$$

$$\therefore n \times 0.3 = 6 \quad \therefore n = \frac{6}{0.3} = 20$$

EXERCISE 8.1

- (1) A die is thrown 6 times. If 'getting an odd number' is a success, find the probability of
 - (i) 5 successes
 - (ii) at least 5 successes
 - (iii) at most 5 successes.
- (2) A pair of dice is thrown 4 times. If getting a doublet is considered a success, find the probability of two successes.
- (3) There are 5% defective items in a large bulk of items. What is the probability that a sample of 10 items will include not more than one defective item?
- (4) Five cards are drawn successively with replacement from a well-shuffled deck of 52 cards. find the probability that
 - (i) all the five cards are spades
 - (ii) only 3 cards are spades
 - (iii) none is a spade.
- (5) The probability that a bulb produced by a factory will fuse after 150 days of use is 0.05. Find the probability that out of 5 such bulbs
 - (i) none
 - (ii) not more than one
 - (iii) more than one
 - (iv) at least one will fuse after 150 days of use.

- (6) A bag consists of 10 balls each marked with one of the digits 0 to 9. If four balls are drawn successively with replacement from the bag, what is the probability that none is marked with the digit 0?
- (7) On a multiple choice examination with three possible answers for each of the five questions, what is the probability that a candidate would get four or more correct answers just by guessing?
- (8) A person buys a lottery ticket in 50 lotteries, in each of which his chance of winning a prize is $1/100$. find the probability that he will win a prize
 (i) at least once (ii) exactly once (iii) at least twice
- (9) In a box of floppy discs it is known that 95% will work. A sample of three of the discs is selected at random. Find the probability that
 (i) none (ii) 1 (iii) 2 (iv) all 3 of the sample will work.
- (10) Find the probability of throwing at most 2 sixes in 6 throws of a single die.
- (11) It is known that 10% of certain articles manufactured are defective. What is the probability that in a random sample of 12 such articles, 9 are defective?
- (12) Given that $X \sim B(n, p)$
 (i) If $n = 10$ and $p = 0.4$, find $E(X)$ and $Var(X)$ (ii) If $p = 0.6$ and $E(X) = 6$, find n and $Var(X)$.
 (iii) If $n = 25$, $E(X) = 10$ find p and $SD(X)$. (iv) If $n = 10$, $E(X) = 8$, find $Var(X)$.



Let us Remember

✿ Trials of a random experiment are called **Bernoulli trials**, if they satisfy the following conditions :

- (i) Each trial has exactly two outcomes : success or failure.
- (ii) The probability of success remains the same in each trial.

Thus probability of getting x successes in n -Bernoulli trial is

$$P(x \text{ successes out of } n \text{ trials}) = \frac{n!}{x!(n-x)!} \times p^x \times q^{n-x} = {}^n C_x p^x \times q^{n-x}$$

Clearly, $P(x \text{ successes})$, i.e. ${}^n C_x p^x q^{n-x}$ is the $(x + 1)^{\text{th}}$ term in the binomial expansion of $(q + p)^n$.

✿ Let $X \sim B(n, p)$ then mean of expected value of r.v. X is denoted by μ .
 $E(X)$ and given by $\mu = E(X) = np$.

The variance is denoted by $Var(X)$ and given by $Var(X) = npq$.

Standard deviation of X is denoted by $SD(X)$ or σ and given by $SD(X) = \sigma_x = \sqrt{Var(X)}$

MISCELLANEOUS EXERCISE 8

(I) Choose the correct option from the given alternatives :

- (1) A die is thrown 100 times. If getting an even number is considered a success, then the standard deviation of the number of successes is
 (A) $\sqrt{50}$ (B) 5 (C) 25 (D) 10
- (2) The mean and the variance of a binomial distribution are 4 and 2 respectively. Then the probability of 2 successes is
 (A) $\frac{128}{256}$ (B) $\frac{219}{256}$ (C) $\frac{37}{256}$ (D) $\frac{28}{256}$
- (3) For a binomial distribution, $n = 5$. If $P(X = 4) = P(X = 3)$ then $p = \dots$
 (A) $\frac{1}{3}$ (B) $\frac{3}{4}$ (C) 1 (D) $\frac{2}{3}$
- (4) In a binomial distribution, $n = 4$. If $2P(X = 3) = 3P(X = 2)$ then $p = \dots$
 (A) $\frac{4}{13}$ (B) $\frac{5}{13}$ (C) $\frac{9}{13}$ (D) $\frac{6}{13}$
- (5) If $X \sim B(4, p)$ and $P(X = 0) = \frac{16}{81}$, then $P(X = 4) = \dots$
 (A) $\frac{1}{16}$ (B) $\frac{1}{81}$ (C) $\frac{1}{27}$ (D) $\frac{1}{8}$
- (6) The probability of a shooter hitting a target is $\frac{3}{4}$.
 How many minimum number of times must he fire so that the probability of hitting the target at least once is more than 0.99 ?
 (A) 2 (B) 3 (C) 4 (D) 5
- (7) If the mean and variance of a binomial distribution are 18 and 12 respectively, then $n = \dots$
 (A) 36 (B) 54 (C) 18 (D) 27

(II) Solve the following :

- (1) Let $X \sim B(10, 0.2)$, Find (i) $P(X = 1)$ (ii) $P(X \geq 1)$ (iii) $P(X \leq 8)$.
- (2) Let $X \sim B(n, p)$
 (i) If $n = 10$, $E(X) = 5$, find p and $Var(X)$.
 (ii) If $E(X) = 5$ and $Var(X) = 2.5$, find n and p .
- (3) If fair coin is tossed 10 times find the probability that it shows heads
 (i) 5 times. (ii) in the first four tosses and tail in last six tosses.

- (4) Probability that bomb will hit target is 0.8 . Find the probability that out of 10 bombs dropped exactly 2 will miss the target.
- (5) The probability that a mountain-bike rider travelling along a certain track will have a tyre burst is 0.05 . Find the probability that among 17 riders : (i) exactly one has a burst tyre
(ii) at most three have a burst tyre (iii) two or more have burst tyres.
- (6) Probability that a lamp in a classroom will burnt out will be 0.3 . Six lamps are fitted in the classroom. If it is known that the classroom is unusable if the number of lamps burning in it is less than four, find the probability that classroom can not used at random occasion.
- (7) Lot of 100 items contains 10 defective items. Five items are selected at random from the lot and sent to the retail store. What is the probability that the store will receive at most one defective item?
- (8) A large chain retailer purchases certain kind of electric device from manufacturer. The manufacturer indicates that the defective rate of the device is 3% . The inspector of the retailer picks 20 items from a shipment. What is the probability that the store will receive at most one defective item?
- (9) The probability that the certain kind of component will survive a check test is 0.6 . Find the probability that exactly 2 of the next 4 components tested survive.
- (10) An examination consists of 10 multiple-choice questions, in each of which a candidate has to deduce which one of five suggested answers is correct. A completely unprepared student guesses each answer completely randomly. What is the probability that this student gets 8 or more questions correct? Draw the appropriate moral !
- (11) The probability that a machine will produce all bolts in a production run within specification is 0.998 . A sample of 8 machines is taken at random. Calculate the probability that
(i) all 8 machines (ii) 7 or 8 machines
(iii) at least 6 machines will produce all bolts within specification
- (12) The probability that a machine develops a fault within the first 3 years of use is 0.003 . If 40 machines are selected at random, calculate the probability that 38 or more will not develop any faults within the first 3 years of use.
- (13) A computer installation has 10 terminals. Independently, the probability that any one terminal will require attention during a week is 0.1 . Find the probabilities that
(i) 0 (ii) 1 (iii) 2
(iv) 3 or more, terminals will require attention during the next week.

- (14) In a large school, 80% of the pupils like mathematics. A visitor to the school asks each of 4 pupils, chosen at random, whether they like mathematics.
- (i) Calculate the probabilities of obtaining an answer yes from 0, 1, 2, 3, 4 of the pupils
 - (ii) Find the probability that the visitor obtains the answer yes from at least 2 pupils:
 - (a) when the number of pupils questioned remains at 4
 - (b) when the number of pupils questioned is increased to 8.
- (15) It is observed that, it rains on 12 days out of 30 days. Find the probability that
- (i) it rains exactly 3 days of week.
 - (ii) it will rain on at least 2 days of given week.
- (16) If probability of success in a single trial is 0.01. How many trials are required in order to have probability greater than 0.5 of getting at least one success?
- (17) In binomial distribution with five Bernoulli's trials, probability of one and two success are 0.4096 and 0.2048 respectively. Find probability of success.



ANSWERS

1. DIFFERENTIATION

EXERCISE 1.1

- (1) (i) $5(3x^2 - 2)(x^3 - 2x - 1)^4$
 (ii) $\frac{5}{2}(3\sqrt{x} - 4\sqrt[3]{x})(2x^{\frac{3}{2}} - 3x^{\frac{4}{3}} - 5)^{\frac{3}{2}}$
 (iii) $\frac{x+2}{\sqrt{x^2+4x-7}}$
 (iv) $\frac{x(2\sqrt{x^2+1}+1)}{2\sqrt{x^2+1} \cdot \sqrt{x^2+\sqrt{x^2+1}}}$
 (v) $-\frac{4x-7}{(2x^2-7x-5)^{\frac{8}{3}}}$
 (vi) $\frac{15(3x-4)}{2(3x-5)^{\frac{3}{2}}}\left(\sqrt{3x-5}-\frac{1}{\sqrt{3x-5}}\right)^4$
- (2) (i) $-2x \sin(x^2+a^2)$ (ii) $\frac{3e^{(3x+2)}}{2\sqrt{e^{(3x+2)}+5}}$
 (iii) $\operatorname{cosec} x$ (iv) $\frac{\sec^2 \sqrt{x}}{4\sqrt{x} \cdot \sqrt{\tan \sqrt{x}}}$
 (v) $\frac{-9 \operatorname{cosec}^2[\log(x^3)] \cdot \cot^2[\log(x^3)]}{x}$
 (vi) $3 \sin^2 x \cdot \cos x \cdot 5^{\sin^3 x + 3} \cdot \log 5$
 (vii) $\frac{\sin x \operatorname{cosec} \sqrt{\cos x} \cdot \cot(\sqrt{\cos x})}{2\sqrt{\cos x}}$
 (viii) $-3x^2 \tan(x^3-5)$
 (ix) $5 \sin 2x \cdot e^{3 \sin^2 x - 2 \cos^2 x}$
 (x) $\frac{-2x \cdot \sin[2 \log(x^2+7)]}{x^2+7}$
 (xi) $-\sec^2[\cos(\sin x)] \cdot \sin(\sin x) \cdot \cos x$
 (xii) $4x^3 \cdot \sec^2(x^4+4) \cdot \sec[\tan(x^4+4)] \cdot \tan[\tan(x^4+4)]$

- (xiii) $\frac{2 \log x}{x} - \frac{2}{x}$
 (xiv) $\frac{\cos \sqrt{\sin \sqrt{x}} \cdot \cos \sqrt{x}}{4\sqrt{x} \cdot \sqrt{\sin \sqrt{x}}}$
 (xv) $2x \cdot e^{x^2} [\tan(e^{x^2})]$ (xvi) $\frac{1}{2x \log x}$
 (xvii) $\frac{2[\log[\log(\log x)]]}{x \log x \cdot \log(\log x)}$
 (xviii) $4x \sin(2x^2)$
- (3) (i) $6(x+2)(x^2+4x+1)^2 + 4(3x^2-5)(x^3-5x-2)^3$
 (ii) $8(1-2x)(1+4x)^5(3+x-x^2)^7 + 20(1+4x)^4(3+x-x^2)^8$
 (iii) $\frac{14-3x}{2(7-3x)^{\frac{3}{2}}}$ (iv) $\frac{6x^2(x^3+15)(x^3-5)^4}{(x^3+3)^6}$
 (v) $\sin 2x(1+\sin^2 x)(1+\cos^2 x)^2(1-5\sin^2 x)$
 (vi) $-\frac{\sin x}{2\sqrt{\cos x}} - \frac{\sin \sqrt{x}}{4\sqrt{x} \cdot \sqrt{\cos \sqrt{x}}}$
 (vii) $3 \sec 3x$ (viii) $\frac{\pi \cos x^\circ}{90(1-\sin x^\circ)^2}$
 (ix) $-\frac{\operatorname{cosec}^2\left(\frac{\log x}{2}\right)}{2x} + \tan x \cdot \operatorname{cosec}^2 x$
 (x) $\frac{8e^{4x}}{(e^{4x}+1)^2}$ (xi) $-\frac{e^{\sqrt{x}}}{\sqrt{x}(e^{\sqrt{x}}-1)^2}$
 (xii) $6 \operatorname{cosec} 2x + 4 \cot x + \frac{14x}{x^2+7}$
 (xiii) $3 \operatorname{cosec} 3x$ (xiv) $-\frac{5}{2} \operatorname{cosec}\left(\frac{5x}{2}\right)$

- (xv) $-\sec x$
 (xvi) $2 \log 4 + \frac{3x}{x^2+5} - \frac{9x^2}{2(2x^3-4)}$
 (xvii) $2x - \frac{6}{5-4x} + \frac{2}{7-6x}$
 (xviii) $-\sin x \log a - \frac{6x}{x^2-3} - \frac{1}{x \log x}$
 (xix) 0 (xx) $\frac{x(x^2+2)^3(7x^2+38)}{(x^2+5)^{\frac{3}{2}}}$
- (4) (i) -16 (ii) 35 (iii) -20 (iv) 28
 (5) -5 (6) $\frac{12}{5}$ (7) $x = 0$ or $\frac{2\pi}{3}$ or 2π
 (8) $e^{2x} + 6e^x + 14, e^{x^2+5}, 2x, e^x, f'[g(x)] \cdot g'(x),$
 $2e^{2x} + 6e^x, 8, g'[f(x)] \cdot f'(x), 2xe^{x^2+5}, -2e^6.$

EXERCISE 1.2

- (1) (i) $\frac{1}{2\sqrt{x}}$ (ii) $-\frac{1}{4\sqrt{x}\sqrt{2-\sqrt{x}}}$
 (iii) $\frac{1}{3\sqrt[3]{(x-2)^2}}, \text{ for } x > 2$ (iv) $\frac{2}{2x-1}$
 (v) 2 (vi) e^x
 (vii) $2e^{2x-3}$ (viii) $\frac{1}{x \log 2}$
- (2) (i) $\frac{1}{x \cdot e^x(x+2)}$ (ii) $\frac{1}{\cos x - x \sin x}$
 (iii) $\frac{1}{7^x(x \log 7 + 1)}$ (iv) $\frac{x}{2x^2+1}$
 (v) $\frac{1}{1+\log x}$
- (3) (i) $\frac{1}{14}$ (ii) $\frac{1}{4}$ (iii) $\frac{1}{12}$ (iv) $\frac{1}{5}$
 (4) 1
 (5) (i) and (ii) derivative proved.

- (6) (i) $\frac{1}{x[1+(\log x)^2]}$ (ii) $\frac{e^x}{\sqrt{1-e^{2x}}}$
 (iii) $-\frac{3x^2}{1+x^6}$ (iv) $-\frac{4^x \log 4}{1+4^{2x}}$
 (v) $\frac{1}{2\sqrt{x}(1+x)}$ (vi) $\frac{x}{\sqrt{1-x^4}}$
 (vii) $\frac{2}{\sqrt{2-x^2}}$ (viii) $\frac{3\sqrt{x}}{2\sqrt{1-x^3}}$
 (ix) $9x^8$ (x) $2x$
- (7) (i) $2xe^{x^2}$ (ii) $-5^x \log 5$ (iii) $\frac{1}{2}$
 (iv) $-x$ (v) $-\frac{1}{2}$ (vi) -6
 (vii) $-\frac{1}{6}$ (viii) $-\frac{3}{2}$ (ix) $-\frac{7}{2}$
 (x) $-\frac{1}{2}$ (xi) $-\frac{1}{2}$ (xii) $\frac{2}{3}$
- (8) (i) 1 (ii) 1 (iii) $\frac{1}{2\sqrt{x}}$
 (iv) 3 (v) e^x (vi) $2^x \log 2$
- (9) (i) $\frac{2}{1+x^2}$ (ii) $\frac{2}{1+x^2}$
 (iii) $-\frac{2}{1+x^2}$ (iv) $\pm \frac{2}{\sqrt{1-x^2}}$
 (v) $-\frac{3}{\sqrt{1-x^2}}$ (vi) $-\frac{2e^x}{1+e^{2x}}$
 (vii) $\frac{2 \cdot 3^x \log 3}{1+3^{2x}}$
 (viii) $\frac{2 \cdot 4^x \log 4}{1+4^{2x}}$ or $\left(\frac{4^{x+\frac{1}{2}} \log 4}{1+4^{2x}} \right)$
 (ix) $-\frac{10}{1+25x^2}$ (x) $-\frac{3\sqrt{x}}{1+x^3}$
 (xi) $\frac{5x\sqrt{x}}{1+x^5}$ (xii) $\frac{1}{2\sqrt{x}(1+x)}$
- (10) (i) $\frac{3}{1+9x^2} + \frac{5}{1+25x^2}$

(ii) $\frac{7}{1+49x^2} - \frac{5}{1+25x^2}$

(vi) $\frac{3a}{a^2+9x^2} + \frac{2a}{a^2+4x^2}$ (vii) 1

(iii) $\frac{1}{2\sqrt{x}} \left(\frac{3}{1+9x} - \frac{1}{1+x} \right)$

(viii) $\frac{2}{1+(2x+1)^2} - \frac{3}{1+(3x-4)^2}$

(iv) $2^x \log 2 \left(\frac{3}{1+9(2^{2x})} + \frac{1}{1+2^{2x}} \right)$

(ix) $\frac{2}{1+(2x+3)^2} + \frac{1}{1+(x-1)^2}$

(v) $2^x \log 2 \left(\frac{2}{1+4(2^{2x})} - \frac{1}{1+2^{2x}} \right)$

EXERCISE 1.3

(1) (i) $\frac{(x+1)^2}{(x+3)^3(x+3)^4} \left[\frac{2}{x+1} - \frac{3}{x+2} - \frac{4}{x+3} \right]$

(ii) $\frac{1}{3} \sqrt[3]{\frac{4x-1}{(2x+3)(5-2x)^2}} \left(\frac{4}{4x-1} - \frac{2}{2x+3} + \frac{4}{5-2x} \right)$

(iii) $(x^2+3)^{\frac{3}{2}} \cdot \sin^3 2x \cdot 2^{x^2} \left[\frac{3x}{x^2+3} + 6 \cot 2x + 2x \log 2 \right]$

(iv) $\frac{(x^2+2x+2)^{\frac{3}{2}}}{(\sqrt{x}+3)^3(\cos x)^x} \left[\frac{3(x+1)}{x^2+2x+2} - \frac{3}{2\sqrt{x}(\sqrt{x}+3)} + x \tan x - \log(\cos x) \right]$

(v) $\frac{x^5 \cdot \tan^3 4x}{\sin^2 3x} \left[\frac{5}{x} + 24 \operatorname{cosec} 8x - 6 \cot 3x \right]$

(vi) $x^{\tan^{-1}x} \left[\frac{\tan^{-1}x}{x} + \frac{\log x}{1+x^2} \right]$

(vii) $\sin^x x [x \cot x + \log(\sin x)]$

(viii) $\cos(x^x) \cdot x^x(1 + \log x)$

(2) (i) $ex^{e-1} + e^x + x^x(1 + \log x)$ (ii) $x^{x^x} \cdot x^x \cdot \log x \left[1 + \log x + \frac{1}{x \log x} \right] + e^{x^x} \cdot x^x(1 + \log x)$

(iii) $(\log x)^x \left[\frac{1}{\log x} + \log(\log x) \right] + (\cos x)^{\cot x} [1 + \operatorname{cosec}^2 x \log(\cos x)]$

(iv) $x^{e^x} \cdot e^x \left[\frac{1}{x} + \log x \right] + (\log x)^{\sin x} \left[\frac{\sin x}{x \log x} + \cos x \log(\log x) \right]$

(v) $\sec^2 x \cdot e^{\tan x} + (\log x)^{\tan x} \left[\frac{\tan x}{x \log x} + \sec^2 x \log(\log x) \right]$

(vi) $(\sin x)^{\tan x} [1 + \sec^2 x \log(\sin x)] + (\cos x)^{\cot x} [1 + \operatorname{cosec}^2 x \log(\cos x)]$

(vii) $10^{x^x} x^x \log 10 (1 + \log x) + x^{x^{10}} \cdot x^9 (1 + 10 \log x) + x^{10^x} \cdot 10^x \left(\frac{1}{x} + \log x \cdot \log 10 \right)$

(viii) 2

- (3) (i) $-\frac{\sqrt{y}}{\sqrt{x}}$ (ii) $-\frac{\sqrt{x}}{\sqrt{y}}$
 (iii) $-\frac{\sqrt{y}(2\sqrt{x}+\sqrt{y})}{\sqrt{x}(2\sqrt{y}+\sqrt{x})}$ (iv) $-\frac{3x^2+2xy+y^2}{x^2+2xy+3y^2}$
 (v) $-\frac{y}{x}$ (vi) $-\frac{e^y+ye^x}{e^x+xe^y}$
 (vii) $\frac{\sin(x-y)+e^{x+y}}{\sin(x-y)-e^{x+y}}$ (viii) $-\frac{1+y\sin(xy)}{1+x\sin(xy)}$
 (ix) $\frac{y(1-xe^{x-y})}{x(1-ye^{x-y})}$
 (x) $\frac{\sin(x-y)-\cos(x+y)-1}{\sin(x-y)+\cos(x+y)-1}$

EXERCISE 1.4

- (1) (i) $\frac{1}{t}$ (ii) $\frac{b}{a}\cos\theta$ (iii) $\frac{2}{\sqrt{a^2+m^2}}$
 (iv) $\sec^3\theta$ (v) $\frac{b}{a}\tan\left(\frac{\theta}{2}\right)$
 (vi) $\frac{y(t^2+1)\log a}{axt}$ (vii) $-\frac{1}{2}$ (viii) $\frac{1}{3}$
 (2) (i) $\frac{3\sqrt{3}}{2}$ (ii) $-\sqrt{3}$ (iii) $-\frac{\pi}{6}$
 (iv) $1-\sqrt{2}$ (v) $3+\pi$
 (4) (i) $\frac{x\cos x+\sin x}{\sec^2 x}$ (ii) 1
 (iii) $-\frac{1}{2}$ (iv) 2 (v) $-x(\log x)^2 \cdot 3^x$
 (vi) $-\frac{x\sqrt{x^2-1}}{2}$
 (vii) $\frac{(1+\log x) \cdot x^{x+1-\sin x}}{\sin x+x\cos x \cdot \log x}$
 (viii) $\frac{\sqrt{1-x^2}}{4(1+x^2)}$

EXERCISE 1.5

- (1) (i) $40x^3-24x-\frac{12}{x^4}$
 (ii) $2e^{2x}(1+\tan x) \cdot (2+\tan x+\tan^2 x)$
 (iii) $-e^{4x}(9\cos 5x+40\sin 5x)$
 (iv) $x(5+6\log x)$ (v) $-\frac{1+\log x}{(x\log x)^2}$
 (iv) $x^{x-1}+x^x(1+\log x)^2$
 (2) (i) $-\frac{1}{4a}\operatorname{cosec}^4\left(\frac{\theta}{2}\right)$ (ii) $-\frac{1}{4at^3}$
 (iii) 6 (iv) $-\frac{2\sqrt{2}b}{a^2}$
 (4) (i) $\frac{d^ny}{dx^n} = \frac{m! a^n (ax+b)^{m-n}}{(m-n)!}$ if $m > 0, m > n$,
 $\frac{d^ny}{dx^n} = 0$ if $m > 0, m < n$
 $\frac{d^ny}{dx^n} = n! a^n$ if $m > 0, m = n$
 (ii) $\frac{(-1)^n n!}{x^{n+1}}$ (iii) $a^n e^{ax+b}$
 (iv) $p^n a^{px+q} (\log a)^n$
 (v) $\frac{(-1)^{n-1} (n-1)! a^n}{(ax+b)^n}$ (vi) $\cos\left(\frac{n\pi}{2}+x\right)$
 (vii) $a^n \sin\left(\frac{n\pi}{2}+ax+b\right)$
 (viii) $(-2)^n \cos\left(\frac{n\pi}{2}+3-2x\right)$
 (ix) $\frac{(-1)^{n-1} (n-1)! 2^n}{(2x+3)^n}$
 (x) $\frac{(-1)^n \cdot n! \cdot 3^n}{(3x-5)^{n+1}}$
 (xi) $e^{ax} (a^2+b^2)^{\frac{n}{2}} \cdot \cos\left[bx+c+n\tan^{-1}\left(\frac{b}{a}\right)\right]$
 (xii) $e^{8x} \cdot (10)^n \cos\left[6x+7+n\tan^{-1}\left(\frac{3}{4}\right)\right]$

MISCELLANEOUS EXERCISE 1

(I)

1	2	3	4	5	6	7	8	9	10	11	12
D	C	C	B	A	C	D	C	B	C	A	B

- (II) (1) $\frac{3}{4}$ (ii) Does not exist (iii) -2 (iv) $-\frac{1}{2\sqrt{1-x^2}}$ [Hint : $x = \cos 2\theta$]
- (2) (A) 3, (B) 5, (C) 4, (D) 1. (v) $\frac{3}{1+9x^2} + \frac{5}{1+25x^2}$
- (3) (i) $-\frac{1}{9}$ (ii) $-\frac{40}{3}$ (iii) $-\frac{29}{96}$ (vi) $\frac{1}{2(1+x^2)}$ [Hint : $x = \tan \theta$]
- (iv) $-\frac{4}{9}$ (6) (i) $\frac{\sqrt{1-x^2}}{4(1+x^2)}$
- (4) (i) $-\frac{x}{\sqrt{1-x^2}}$ [Hint : $x = \cos 2\theta$] (ii) $-\frac{2x}{\sqrt{1+x^2} \cdot \sin(\log x)}$ (iii) 1
- (ii) $-\frac{1}{2}$ [Hint : $x = \cos 2\theta$]
- (iii) $\frac{3}{2\sqrt{x}(1+x)}$ [Hint : $\sqrt{x} = \tan \theta$]

2. APPLICATIONS OF DERIVATIVES

EXERCISE 2.1

- (1) (i) $2x - y + 4 = 0, x + 2y - 8 = 0$ (3) $(2, -2) \left(-\frac{2}{3}, -\frac{14}{27}\right)$ (4) $y = 0$ and $y = 4$
- (ii) $4x - 5y + 12 = 0, 5x + 4y - 26 = 0,$ (5) $x + 3y - 8 = 0, x + 3y + 8 = 0$
- (iii) $y = 2, x = \sqrt{3}$ (6) $a = 2, b = -7$ (7) $(4, 11)$ and $\left(-4, -\frac{31}{3}\right)$
- (iv) $\pi x + 2y - 2\pi = 0,$ (8) $0.8 \pi \text{ cm}^2/\text{sec}.$ (9) $6 \text{ cm}^3/\text{sec}.$
- $4x - 2\pi y + \pi^2 - 4 = 0$ (10) $\frac{3\sqrt{6}}{2} \text{ cm}^2/\text{sec}.$ (11) $8 \text{ cm}^2/\text{sec}$
- (v) $2x - y = 0, 4x + 8y - 5\pi = 0$ (12) $7.2 \text{ cm}^3/\text{sec}$ (13) 3 km/hr
- (vi) $4x + 2y - 3 = 0, 2x - 4y + 1 = 0$ (14) (i) $\left(\frac{3}{8}\right) \text{ meter/sec}.$ (ii) $\frac{9}{8} \text{ meter/sec}.$
- (vii) $17x - 4y - 20 = 0, 8x + 34y - 135 = 0$ (15) $0.9 \text{ meter/sec}.$ (16) $\left(\frac{4\pi}{3}\right) \text{ cm}^3/\text{sec}$
- (2) (4, 1)

EXERCISE 2.2

- (1) (i) 2.9168 (ii) 3.03704 (iii) 1.9997
 (iv) 248.32 (v) 64.48
- (2) (i) 0.953 (ii) 0.42423 (iii) 0.4924
 (iv) 1.02334
- (3) (i) 0.7845 (ii) 0.7859 (iii) 0.7859
- (4) (i) 2.70471 (ii) 8.1279 (iii) 9.09887
- (5) (i) 4.6152 (ii) 2.1983 (iii) 3.006049
- (6) (i) 6.91 (ii) 9.72

EXERCISE 2.3

- (1) (i) Valid (ii) Valid
 (iii) Invalid (iv) Valid
 (v) Invalid (vi) Invalid
- (2) $b = 1$
- (3) (i) $\frac{\pi}{4}$ or $\frac{5\pi}{4}$ (ii) $c = \pi$ (iii) $c = \frac{5}{2}$
- (4) $p = -6, q = 11$ (6) $c = -2$
- (7) (i) $e - 1$ (ii) $2 \pm \frac{2}{\sqrt{3}}$ (iii) $\frac{1}{7}$
 (iv) $\frac{1}{2}$ (v) $3 + \sqrt{2}$

EXERCISE 2.4

- (1) (i) Increasing $\forall x \in R$
 (ii) Decreasing $\forall x \in R$
 (iii) Increasing $\forall x \in R$
- (2) (i) $x < -1$ and $x > 2$ (ii) $R - \{1\}$
 (iii) $x < -2$ and $x > 6$
- (3) (i) $-1 < x < 2$ (ii) $(-5, 5) - \{0\}$
 (iii) $x \in (2, 4)$
- (4) (a) $(-\infty, -4] \cup [12, \infty)$
 (b) $-4 \leq x \leq 12$ i.e. $[-4, 12]$
- (5) (a) $x < -3$ and $x > 8$ (b) $-3 < x < 8$
- (6) (a) $-1 < x < 1$ (b) $(-\infty, -1) \cup (1, \infty)$
- (9) (i) $\text{Max} = \frac{36}{25}, \text{Min} = -\frac{16}{27}$
 (ii) $\text{Max} = -3, \text{Min} = -128$
 (iii) $\text{Max} = 20, \text{Min} = 16$ (iv) $\text{Min} = 8$
 (v) $\text{Min} = -\frac{1}{e}$ (vi) $\text{Max} = \frac{1}{e}$
- (10) 15, 15 (11) 10, 10 (12) 9 (13) 12.8
- (14) $l = \sqrt{2}$ and $b = \frac{1}{\sqrt{2}}$
- (15) Radius = Height = a (16) 3, 3
- (17) Side of square base = 8 cm, Height = 4 cm
- (18) $x = 75, P = 4000$ (19) 6, 9
- (22) $\frac{4\pi r^3}{3\sqrt{3}} \text{ cm}^3$

MISCELLANEOUS EXERCISE 2

(I)

1	2	3	4	5	6	7	8	9	10
A	C	B	B	D	C	D	A	D	D

(II) (2) 4

(3) $14x - 13y + 12 = 0, 13x + 14y - 41 = 0$

(4) $\frac{2}{9\pi}$ ft/sec (5) $\left(\frac{16}{3}, 3\right), \left(-\frac{16}{3}, -3\right)$

(6) $c = 0$ (7) $c = 2$ (8) 2.025

(9) 1.03565

(10) Decreasing in $\left(0, \frac{1}{e}\right]$ and

Increasing in $\left[\frac{1}{e}, \infty\right)$

(11) Increasing in $[e, \infty)$, Decreasing in $(1, e]$

(15) $l = \frac{60}{\pi + 4}, b = \frac{30}{\pi + 4}, r = \frac{30}{\pi + 4}$

(17) Side = $\frac{l}{\pi + 4}$, Radius = $\frac{l}{2(\pi + 4)} = \frac{x}{2}$

(18) 24, 45 (21) Max = $\frac{5}{4}$, Min = 1

3. INDEFINITE INTEGRATION

EXERCISE 3.1

(1) (i) $\frac{x^4}{4} + \frac{x^3}{3} - \frac{x^2}{2} + x + c$ (ii) $\frac{x^3}{3} - 2x^2 + 4x + c$

(iii) $3 \tan x - 4 \log x - \frac{2}{\sqrt{x}} - 7x + c$

(iv) $\frac{x^2}{4} - \frac{5x^2}{2} + 3 \log x - \frac{1}{x^4} + c$

(v) $\frac{6}{5} x^2 \sqrt{x} - 4\sqrt{x} - \frac{10}{\sqrt{x}} + c$

(2) (i) $\tan x - x + c$ (ii) $-2 \cos x + c$

(iii) $\sec x + c$ (iv) $-\cot x - 2x + c$

(v) $-\cot x - \tan x + x + c$

(vi) $\sec x - \tan x + x + c$

(vii) $\sec x - \tan x + x + c$

(viii) $\sin x - \cos x + c$ (ix) $-\sqrt{2} \cos x + c$

(x) $-\frac{1}{14} \cos 7x - \frac{1}{2} \cos x + c$

(3) (i) $x - 2 \log(x + 2) + c$

(ii) $2x + \frac{1}{2} \log(2x + 1) + c$

(iii) $\frac{5}{3} x - \frac{26}{9} \log(3x - 4) + c$

(iv) $\frac{2(x+5)^{\frac{3}{2}}}{3} - 14\sqrt{x+5} + c$

(v) $\frac{1}{12} (4x-1)^{\frac{3}{2}} - \frac{13}{4} \sqrt{4x-1} + c$

(vi) $-\cos 2x + c$

(vii) $\frac{2}{5} \left(\sin \frac{5x}{2} - \cos \frac{5x}{2} \right) + c$

(viii) $\frac{1}{4} (2x + \sin 2x) + c$

(ix) $-\frac{4}{9} \left[x^{\frac{3}{2}} + (x+3)^{\frac{3}{2}} \right] + c$

(x) $\frac{2}{21} \left[(7x-2)^{\frac{3}{2}} + (7x-5)^{\frac{3}{2}} \right] + c$

(4) $f(x) = \frac{x^2}{2} + \frac{3}{2x^2} + \frac{7}{2}$

EXERCISE 3.2 (A)

I. 1. $\frac{(\log x)^{n+1}}{n+1} + c$ 2. $\frac{2}{5} (\sin^{-1} x)^{\frac{5}{2}} + c$

3. $\log(\operatorname{cosec}(x + \log x) - \cot(x + \log x)) + c$

4. $\frac{-1}{\sqrt{\tan(x^2)}} + c$ 5. $\frac{1}{3} (e^{3x} + 1) + c$

6. $\frac{1}{\log a} \cdot a^{x+\tan^{-1}x} + c$
7. $\frac{1}{2} [\log (\sin e^x)]^2 + c$
8. $\log (e^x - e^{-x}) + c$
9. $\frac{1}{5} \sin^5 x - \frac{1}{7} \sin^7 x + c$
10. $\frac{1}{48} \log (4x^{12} + 5) + c$
11. $\frac{1}{10} \tan x^{10} + c$ 12. $\frac{1}{4} \log (x^4 + 1) + c$
13. $2 \sqrt{\tan x} + c$ 14. $\tan^{-1} x + \frac{1}{x^2 + 1} + c$
15. $\log (3 \cos^2 x + 4 \sin^2 x) + c$
16. $2 \tan^{-1} \sqrt{x} + c$ 17. $\log (10^x + x^{10}) + c$
18. $\frac{\sqrt{1+4x^n}}{2n} + c$
19. $\frac{4}{5} (x+2)^{\frac{5}{2}} - 2 (x+2)^{\frac{3}{2}} + c$
20. $\frac{1}{7} (a^2+x^2)^{\frac{7}{2}} - \frac{4a^2}{5} (a^2+x^2)^{\frac{5}{2}} + \frac{2a^4}{3} (a^2+x^2)^{\frac{3}{2}} + c$
21. $-2 \sqrt{2-3x} - \frac{2}{9} (2-3x)^{\frac{3}{2}} + c$
22. $\frac{5}{12} (2x+3)^{\frac{3}{2}} - \frac{11}{2} (2x+3)^{\frac{1}{2}} - \frac{49}{4 \sqrt{2x+3}} + c$
23. $\frac{1}{3} \sin^{-1} \left(\frac{x^3}{3} \right) + c$ 25. $\frac{1}{3} \log \left(\frac{x^3-1}{x^3} \right) + c$

24. $\log (\log (\log x)) + c$

II. 1. $2 \cdot \log \left(\sec \frac{x}{2} \right) + c$

2. $\cos a \cdot \log (\sin (x-a)) - (\sin a) x + c$

3. $\cos (a+b) \cdot \log (\sec (x+b)) - (\sin (a+b)) \cdot x + c$

4. $\log (\tan x + 2) + c$

5. $\frac{11}{75} x + \frac{2}{25} \log (3 \sin x + 4 \cos x) + c$

6. $\frac{2x}{13} + \frac{3}{13} \log (2 \cos x + 3 \sin x) + c$

7. $5x - 3 \log |2e^x - 5| + c$

8. $-5x - \log |3e^x - 4| + c$

9. $-x + \frac{7}{8} \log |4e^{2x} - 5| + c$

10. $\frac{\cos^8 x}{8} + \frac{\cos^6 x}{6} + \frac{\cos^4 x}{4} + \frac{\cos^2 x}{2} + \frac{1}{2} \log (\cos^2 x - 1) + c$

11. $\frac{\tan^4 x}{4} - \frac{\tan^2 x}{2} + \log (\sec x) + c$

12. $\sin x - \sin^3 x + \frac{3}{5} \sin^5 x - \frac{1}{7} \sin^7 x + c$

13. $\frac{1}{6} \log \left[\frac{(\sec 3x)^2}{(\sec 2x)^3 (\sec x)^6} \right] + c$

14. $\frac{1}{6} \cos^{11} x - \frac{1}{9} \cos^9 x + \frac{1}{13} \cos^{13} x + c$

15. $-\frac{1}{\log 3} \cdot 3^{\cos^2 x} + c$

16. $\frac{1}{20} \log \left[\frac{\sin^5 4x}{\sin^2 10x} \right] + c$

17. $\frac{1}{2} \log [(1 + \cos^2 x) - \cos^2 x] + c$

EXERCISE 3.2 (B)

I. 1. $\frac{1}{4\sqrt{3}} \log \left(\frac{2x - \sqrt{3}}{2x + \sqrt{3}} \right) + c$

2. $\frac{1}{30} \log \left(\frac{5+3x}{5-3x} \right) + c$

3. $\frac{1}{\sqrt{14}} \tan^{-1} \left(\frac{\sqrt{2}x}{\sqrt{7}} \right) + c$

4. $\frac{1}{\sqrt{3}} \log \left(x + \sqrt{x^2 + \frac{8}{3}} \right) + c$

$$5. \frac{1}{2} \sin^{-1} \left(\frac{2x}{\sqrt{11}} \right) + c$$

$$6. \frac{1}{\sqrt{2}} \log \left(x + \sqrt{x^2 - \frac{5}{2}} \right) + c$$

$$7. 9 \sin^{-1} \left(\frac{x}{9} \right) - \sqrt{9 - x^2} + c$$

$$8. 2 \sin^{-1} \left(\frac{x}{2} \right) - \sqrt{4 - x^2} + c$$

$$9. 2 \sin^{-1} \left(\frac{x}{10} \right) - \frac{1}{2} (\sqrt{100 - x^2}) + c$$

$$10. \frac{1}{4} \log \left| \frac{x+2}{x+6} \right| + c$$

$$11. \frac{1}{\sqrt{5}} \log \left(\frac{\sqrt{5} - 1 + 2x}{\sqrt{5} + 1 - 2x} \right) + c$$

$$12. \frac{1}{8\sqrt{2}} \log \left(\frac{2x - 5 - 2\sqrt{2}}{2x - 5 + 2\sqrt{2}} \right) + c$$

$$13. \frac{1}{2\sqrt{19}} \log \left(\frac{3x + 2 + \sqrt{19}}{3x + 2 - \sqrt{19}} \right) + c$$

$$14. \frac{1}{\sqrt{3}} \log \left(x + \frac{5}{6} + \sqrt{x^2 + \frac{5}{3}x + \frac{7}{3}} \right) + c$$

$$15. \log (x + 4 + \sqrt{x^2 - 8x - 20}) + c$$

$$16. \frac{1}{\sqrt{2}} \log \left(x - \frac{3}{4} + \sqrt{x^2 - \frac{3}{2}x + 4} \right) + c$$

$$17. \log \left(x - \frac{1}{2} + \sqrt{x^2 - x - 6} \right) + c$$

$$18. \frac{1}{2\sqrt{7}} \tan^{-1} \left(\frac{2 \tan x}{\sqrt{7}} \right) + c$$

$$19. \frac{1}{\sqrt{2}} \tan^{-1} (\sqrt{2} \tan x) + c$$

$$20. \frac{1}{2\sqrt{3}} \log \left| \frac{\sqrt{3} + \tan x}{\sqrt{3} - \tan x} \right| + c$$

$$\text{II. } 1. \frac{2}{\sqrt{5}} \tan^{-1} \left(\frac{2 \tan \frac{x}{2} + 2}{\sqrt{5}} \right) + c$$

$$2. \frac{1}{3} \log \left[\frac{3 \tan \left(\frac{x}{2} \right) - 1}{3 \tan \left(\frac{x}{2} \right) + 1} \right] + c$$

$$3. \sqrt{2} \tan^{-1} \left(\frac{\tan \frac{x}{2} - 1}{\sqrt{2}} \right) + c$$

$$4. \tan^{-1} \left[2 \tan \left(\frac{x}{2} \right) + 1 \right] + c$$

$$5. \frac{1}{\sqrt{5}} \tan^{-1} (\sqrt{5} \tan x) + c$$

$$6. -\frac{1}{\sqrt{5}} \tan^{-1} \left(\frac{3 \tan x - 2}{\sqrt{5}} \right) + c$$

$$7. \frac{1}{2\sqrt{11}} \log \left(\frac{\sqrt{11} - 2 + \tan x}{\sqrt{11} + 2 - \tan x} \right) + c$$

$$8. \frac{1}{\sqrt{2}} \log \left[\sec \left(x + \frac{\pi}{4} \right) + \tan \left(x + \frac{\pi}{4} \right) \right] + c$$

$$9. \frac{1}{2} \log \left[\sec \left(x + \frac{\pi}{4} \right) + \tan \left(x + \frac{\pi}{4} \right) \right] + c$$

EXERCISE 3.2 (C)

$$\text{I. } 1. \frac{3}{2} \log (x^2 + 6x + 5) - \frac{5}{4} \log \left(\frac{x+1}{x+5} \right) + c$$

$$2. \log (x^2 + 4x - 5) - \frac{1}{2} \log \left(\frac{x-1}{x+5} \right) + c$$

$$3. \frac{1}{2} \log (2x^2 + 3x - 1) + \frac{3}{2\sqrt{17}}$$

$$\log \left(\frac{4x + 3 - \sqrt{17}}{4x + 3 + \sqrt{17}} \right) + c$$

$$4. \frac{3}{2} \sqrt{2x^2 + 2x + 1} + \frac{5}{2\sqrt{2}}$$

$$\log \left(x + \frac{1}{2} + \sqrt{x^2 + x + \frac{1}{2}} \right) + c$$

$$5. -7 \sqrt{3 + 2x - x^2} + 10 \cdot \sin^{-1} \left(\frac{x-1}{2} \right) + c$$

$$6. \sqrt{x^2 - 16x + 63} + \log \left\{ (x-8) + \sqrt{x^2 - 16x + 63} \right\} + c$$

$$7. \sqrt{9x - x^2} + \frac{9}{2} \sin^{-1} \left(\frac{2x-9}{9} \right) + c$$

$$8. \frac{3}{4\sqrt{2}} \log \left(\frac{2\sqrt{2} \sin x + \sqrt{2} - 2}{2\sqrt{2} \sin x + \sqrt{2} + 2} \right) + c$$

$$9. \sqrt{e^{2x} - 1} - \log (e^x + \sqrt{e^{2x} - 1}) + c$$

EXERCISE 3.3

I. 1. $\frac{x^3}{9} (3 \cdot \log x - 1) + c$

2. $-\frac{x^2}{3} \cos 3x + \frac{2}{9} x \sin 3x + \frac{2}{27} \cos 3x + c$

3. $\frac{x^2}{2} \tan^{-1} x - \frac{1}{2} (x - \tan^{-1} x) + c$

4. $\frac{x^3}{3} \tan^{-1} x - \frac{x^2}{6} + \frac{1}{6} \log (1 + x^2) + c$

5. $\frac{1}{4} (\tan^{-1} x) (x^4 - 1) - \frac{x}{12} (x^3 - 3x) + c$

6. $x [(\log x)^2 - 2 (\log x) + 2] + c$

7. $\frac{1}{2} \log (\sec x + \tan x) + \frac{1}{2} \sec x \cdot \tan x + c$

8. $\frac{1}{4} \left[x^2 - x \cdot \sin 2x - \frac{1}{2} \cos 2x \right] + c$

9. $\frac{x^4}{4} \log x - \frac{x^4}{16} + c$

10. $\frac{e^{2x}}{13} [2 \cos 3x + 3 \sin 3x] + c$

11. $\frac{x^2}{2} \sin^{-1} x + \frac{1}{4} x \sqrt{1-x^2} - \frac{1}{4} \sin^{-1} x + c$

12. $\frac{x^3}{3} \cos^{-1} x - \frac{1}{3} \sqrt{1-x^2} + \frac{1}{9} (1-x^2)^{\frac{3}{2}} + c$

13. $(\log x) [\log (\log x) - 1] + c$

14. $-(\sin^{-1} t) \sqrt{1-t^2} + t + c$

15. $2 [\sqrt{x} \cdot \sin \sqrt{x} + \cos \sqrt{x}] + c$

16. $(\cos \theta) \cdot [1 - \log (\cos \theta)] + c$

17. $\frac{1}{4} \left[\frac{x}{3} \sin 3x + \frac{1}{9} \cos 3x + 3x \sin x + 3 \cos x \right] + c$

18. $-\frac{1}{2} \cos (\log x)^2 + c$

19. $-\frac{1}{2} (\log x)^2 + c$

20. $\frac{x}{6} \sin 3x + \frac{1}{18} \cos x - \frac{1}{14} x \sin 7x - \frac{1}{98} \cos 7x + c$

21. $(3x^{\frac{2}{3}} - 6) \sin \sqrt[3]{x} + 6 \sqrt[3]{x} \cos \sqrt[3]{x} + c$

II. 1. $\frac{e^{2x}}{13} [2 \sin 3x - 3 \cos 3x] + c$

2. $\frac{e^{-x}}{5} [-\cos x + 2 \sin 2x] + c$

3. $\frac{x}{2} [\sin (\log x) - \cos (\log x)] + c$

4. $\sqrt{5} \left[\frac{x}{2} \sqrt{x^2 + \frac{3}{5}} + \frac{3}{10} \log \left(x + \sqrt{x^2 + \frac{3}{5}} \right) \right] + c$

5. $\frac{x^3}{6} \cdot \sqrt{a^2 - x^6} + \frac{a^2}{2} \sin^{-1} \left(\frac{x^3}{a} \right) + c$

6. $\frac{x-5}{2} \sqrt{(x-3)(7-x)} + 2 \sin^{-1} \left(\frac{x-5}{2} \right) + c$

7. $\frac{1}{\log 2} \left\{ \frac{2^x}{2} \sqrt{4^x + 4} + 2 \log (2^x + \sqrt{4^x + 4}) \right\} + c$

8. $\frac{1}{6} (2x^2 + 3)^{\frac{3}{2}} +$

$\sqrt{2} \left[\frac{x}{2} \sqrt{x^2 + \frac{3}{2}} + \frac{3}{4} \log \left(x + \sqrt{x^2 + \frac{3}{2}} \right) \right] + c$

9. $-\frac{1}{3}(5-4x-x^2)^{\frac{3}{2}} - (x+2)\sqrt{5-4x-x^2} - 9\sin^{-1}\left(\frac{x+2}{3}\right) + c$

10. $\frac{(1+2\tan x)}{4}\sqrt{\tan^2 x + \tan x - 7} - \frac{29}{8}\log\left\{\frac{1}{2} + \tan x + \sqrt{\tan^2 x + \tan x - 7}\right\} + c$

11. $\left(\frac{x+1}{2}\right)\sqrt{x^2+2x+5} + 2\log\left\{x+1 + \sqrt{x^2+2x+5}\right\} + c$

12. $\sqrt{2}\left\{\left(\frac{4x+3}{8}\right)\sqrt{x^2+\frac{3}{2}x+2} + \frac{23}{16\sqrt{2}}\log\left[\left(x+\frac{3}{4}\right) + \sqrt{x^2+\frac{3}{2}x+2}\right]\right\} + c$

III. 1. $e^x(2 + \cot x) + c$ 2. $e^x \cdot \tan \frac{x}{2} + c$ 3. $e^x \cdot \frac{1}{x} + c$ 4. $e^x \cdot \left(\frac{1}{x+1}\right) + c$

5. $e^x \cdot (\log x)^2 + c$ 6. $e^{5x} \cdot \log x + c$ 7. $e^{\sin^{-1} x} \cdot x + c$

8. $\frac{(1+x)^2}{2}\left(\log(1+x) - \frac{1}{2}\right) + c$ 9. $x \cdot \operatorname{cosec}(\log x) + c$

EXERCISE 3.4

I. 1. $\frac{1}{4}\log(x-1) - 2\log(x+2) + \frac{11}{4}(x+3) + c$

2. $\frac{1}{6}\tan^{-1}x + \frac{1}{15\sqrt{2}}\log\left(\frac{x-\sqrt{2}}{x+\sqrt{2}}\right) - \frac{\sqrt{3}}{10}\tan^{-1}\left(\frac{x}{\sqrt{3}}\right) + c$

3. $\frac{51}{41}\log(2x+9) + \frac{31}{41}\log(3x-7) + c$ 4. $-\frac{8}{5}\log(x+4) - \frac{2}{5}\log(x-1) + c$

5. $x - \log(x+3) + \log(x-2) + c$ 6. $x^2 + 3x + \frac{5}{3}\log(3x+1) + \log(x-1) + c$

7. $\frac{1}{2}\log\left|\frac{2x+1}{2x-1}\right| + 3\log(x+3) + c$ 8. $\frac{1}{5}\log\left(\frac{x^5}{x^5+1}\right) + c$ 9. $\frac{11}{\sqrt{5}}\tan^{-1}\left(\frac{x}{2}\right) - \frac{9}{2}\tan^{-1}\left(\frac{x}{2}\right) + c$

10. $2\log\left(\frac{x+1}{x-1}\right) + \frac{5}{2\sqrt{2}}\log\left(\frac{x+\sqrt{2}}{x-\sqrt{2}}\right) + c$ 11. $\log\left(\frac{2+x^2}{3+x^2}\right) + c$

12. $\frac{1}{5 \cdot \log 2}\log\left(\frac{2^x-4}{2^{x+1}}\right) + c$ 13. $\frac{5}{2}\left(\frac{1}{x+1}\right) + \frac{11}{4}\log\left(\frac{x+1}{x+3}\right) + c$

14. $6 \cdot \log x - \log(x+1) - \frac{9}{x+1} + c$ 15. $\frac{1}{8}\log\left(\frac{x^6(x^3+3)}{(3x^3+1)^3}\right) + c$

16. $\frac{1}{3}\log(x-1) - \frac{1}{6}\log(x^2+x+1) - \frac{1}{\sqrt{3}}\tan^{-1}\left(\frac{2x+1}{\sqrt{3}}\right) + c$

17. $3 \cdot \log (\sin x - 2) - \frac{4}{\sin x - 2} + c$

18. $\frac{1}{2} \log (\cos x + 1) + \frac{1}{6} \log (\cos x - 1) - \frac{2}{3} \log (2 \cos x + 1) + c$

19. $\frac{1}{8} \log \left(\frac{\cos x - 1}{\cos x + 1} \right) + \frac{1}{4 \cdot (\cos x + 1)} + c$ 20. $\frac{1}{6} \log \left[\frac{(1 + 2 \sin x)^4}{(1 - \sin x)(1 + \sin x)^3} \right] + c$

21. $\frac{1}{10} \log (1 - \cos x) - \frac{1}{2} \log (1 + \cos x) + \frac{2}{5} \log (3 + 2 \cos x) + c$

22. $\frac{1}{2} \log \left[\frac{e^x + 1}{(e^{2x} + 9)^{\frac{1}{2}}} \right] + \frac{1}{6} \tan^{-1} \left(\frac{e^x}{3} \right) + c$ 23. $\frac{5}{26} \log \left[\frac{(3 \log x + 2)^2}{\sqrt{(\log x)^2 + 1}} \right] + \frac{11}{26} \tan^{-1} (\log x) + c$

MISCELLANEOUS EXERCISE 3

(I)

1	2	3	4	5	6	7	8	9	10
B	A	B	A	D	B	A	A	C	B
11	12	13	14	15	16	17	18	19	20
A	A	D	C	A	D	A	D	C	A

(II) (1) $\frac{2}{7} x^{\frac{7}{2}} - \frac{8}{5} x^{\frac{5}{2}} - \frac{8}{3} x^{\frac{3}{2}} + c$

(2) $\frac{x^7}{7} - \frac{x^6}{6} + \frac{x^5}{5} - \frac{x^4}{4} + \frac{x^3}{3} - \frac{x^2}{2} + x - \log (x + 1) + c$

(3) $\frac{1}{15} (6x + 5)^{\frac{5}{2}} + c$

(4) $\frac{t^2}{2} - 2t + 3 \cdot \log (t + 1) + \frac{1}{t + 1} + c$

(5) $3 \tan x - 2 \sec x + c$

(6) $\tan \theta - \cot \theta - 3 \theta + c$

(7) $\frac{1}{48} (2 \sin 6x + 3 \sin 4x + 6 \sin 2x + 12 x) + c$ (8) $\frac{1}{2} \sin 2x - \frac{1}{3} \sin 3x + c$

(9) $\frac{\pi}{4} x - \frac{1}{4} x^2 + c$

(III) (1) $\frac{1}{4} (1 + \log x)^4 + c$

(2) $(\tan^{-1} x) x - \frac{1}{2} \log (1 + x^2) - (1 - x) \tan^{-1} (1 - x) + \frac{1}{2} \log (x^2 - 2x + 2) + c$

(3) $-\cot (\log x) + c$

(4) $\frac{2}{3} \sec x^{\frac{3}{2}} + c$

- | | |
|--|--|
| <p>(5) $x \log (1 + \cos x) + c$</p> <p>(7) $\frac{1}{4} \log (3 - 2 \cot x) + c$</p> <p>(9) $\frac{2}{\sqrt{13}} \tan^{-1} \left(\frac{2 \tan \left(\frac{x}{2}\right) - 3}{\sqrt{13}} \right) + c$</p> <p>(11) $-\frac{3}{2} \sqrt{-2x^2 + x + 3} + \frac{7}{4\sqrt{2}} \sin^{-1} \left(\frac{2x - 1}{\sqrt{7}} \right) + c$</p> <p>(13) $\frac{1}{4} e^{2x} \cdot [\sin 2x - \cos 2x] + c$</p> <p>(14) $\frac{1}{18} \log (3x - 1) + \frac{1}{2} \log (x - 1) - \frac{4}{9} \log (3x - 2) + c$</p> <p>(15) $\frac{1}{6} \log \left\{ \frac{(\cos x - 1)(\cos x + 1)^3}{(2 \cos x + 1)^4} \right\} + c$</p> <p>(16) $\left(\frac{\tan x - 1}{2} \right) \sqrt{7 + 2 \tan x - \tan^2 x} + 4 \sin^{-1} \left(\frac{\tan x - 1}{2\sqrt{2}} \right) + c$</p> <p>(17) $\frac{1}{4} \log \left\{ \frac{(x - 1)^3 (x + 3)}{(x + 1)^4} \right\} + c$</p> <p>(19) $2 \sqrt{\tan x} + c$</p> | <p>(6) $\frac{1}{3} \sin^{-1} (x^3) + c$</p> <p>(8) $x \cdot \left(\log (\log x) - \frac{1}{\log x} \right) + c$</p> <p>(10) $\frac{1}{4} \left(2 \sec^{-1} x + \frac{2\sqrt{x^2 - 1}}{x^2} \right) + c$</p> <p>(12) $x \cdot \log (x^2 + 1) - 2 [x - \tan^{-1} x] + c$</p> <p>(18) $\frac{1}{5} \log \left(\frac{x^5}{x^5 + 1} \right) + c$</p> <p>(20) $\frac{1}{3 \cot^3 x} + \frac{2}{\cot x} - \cot x + c$</p> |
|--|--|

4. DEFINITE INTEGRATION

EXERCISE 4.1

- I. (1) 4 (2) $\frac{64}{3}$ (3) $e^2 - 1$ (4) 6 (5) 20

EXERCISE 4.2

- I. (1) $\frac{64}{3}$ (2) $\log \left(\frac{25}{24} \right)$ (3) $- \left(1 + \frac{\pi}{4} \right)$ (4) 2
- (5) $\frac{1}{18} [13\sqrt{13} + 7\sqrt{7} - 3\sqrt{3} - 27]$ (6) $1 - \frac{3\pi}{4}$ (7) $\frac{4}{7\sqrt{2}}$ (8) 1 (9) $\frac{3\pi}{16}$
- (10) $\frac{1}{3} \left(\tan^{-1} \frac{4}{3} + \tan^{-1} \frac{2}{3} \right)$ (11) π (12) $\frac{\pi}{6}$ (13) 1 (14) $\frac{\pi}{4} - \frac{1}{2}$ (15) 1

- II. (1) $\frac{\pi}{4} - \frac{1}{2} \log 2$ (2) $\frac{1}{2} \log 2$ (3) $\frac{\pi}{4}$
 (4) 0 (5) $\frac{2}{3} \tan^{-1} \left(\frac{1}{3} \right)$ (6) $\frac{1}{4} \log \left(\frac{2\sqrt{2}+1}{2\sqrt{2}-1} \right)$
 (7) $\log \left(\frac{4}{3} \right)$ (8) $\frac{1}{ab} \left[\tan^{-1} \left(\frac{ae}{b} \right) - \tan^{-1} \left(\frac{a}{be} \right) \right]$
 (9) $\frac{\pi}{4}$ (10) $\frac{4}{3}$ (11) $\frac{\pi}{2} - 1$
 (12) $\frac{8}{3}$ (13) $\frac{\pi}{2} - 1$
 (14) $e^{\frac{\pi}{4}} \left[\frac{\pi}{4} + 1 \right] - \left[\frac{\pi}{2} + 1 \right]$ (15) $\sin(\log 3)$

- III. (1) $\frac{\pi}{4}$ (2) 0 (3) 0 (4) 0 (5) $\frac{16}{77}(3)^{\frac{7}{2}}$
 (6) 0 (7) 0 (8) $\frac{\pi^2}{6\sqrt{3}}$ (9) 0 (10) 0
 (11) $4 \log \left(\frac{1+\sqrt{5}}{2} \right)$ (12) 0 (13) $\frac{16}{105}$ (14) $\frac{\pi}{3}$ (15) $\frac{\pi}{2} \log \left(\frac{1}{2} \right)$

MISCELLANEOUS EXERCISE 4

(I)

1	2	3	4	5	6	7	8	9	10
A	A	C	C	D	C	A	D	B	A

- (II) (1) $\frac{1}{10} (3 - \log 3)$ (2) $2 - \sqrt{2}$ (3) $6 - 4 \log 2$ (4) $\frac{1}{8}$ (5) $\frac{1}{21}$
 (6) $\pi - 2$ (7) $\frac{1}{3} \log 2$ (8) $\frac{\pi}{5}$ (9) 0 (10) $\frac{\pi}{2}$
- (III) (1) $\frac{\pi^2}{16}$ (2) $\frac{2}{\sqrt{35}} \tan^{-1} \sqrt{\frac{7}{5}}$ (3) $\frac{1}{\sqrt{5}a} \log \left(\frac{7+3\sqrt{5}}{2} \right)$ (4) $\frac{\pi}{20}$
 (5) $\frac{\pi}{2} - \log 2$ (6) $\frac{1}{2} \left(\frac{\pi}{4} - \log \sqrt{2} \right)$ (7) $-\frac{\pi}{2} \log 2$ (8) $\frac{\pi^3}{6}$
 (9) $\log \left(\frac{5+3\sqrt{3}}{1+\sqrt{3}} \right)$ (10) $\frac{17}{2}$
- (IV) (1) $\frac{1}{2}$ when $a = 0$; $\frac{9}{2}$ when $a = 4$ (2) $k = \frac{1}{2}$

5. APPLICATION OF DEFINITE INTEGRAL

EXERCISE 5.1

- (1) (i) 25 (ii) 16 (iii) 20
 (iv) 1 (v) $2 \log 4$ (vi) $\frac{32}{3}$
 (vii) $\frac{128}{3}$ sq. units
- (2) (i) $\frac{128}{3}$ (ii) $\frac{16}{3}$
- (3) (i) $\frac{1}{12}$ (ii) $\frac{8}{3}$ (iii) $\frac{32}{3}$
 (iv) $8 \frac{a^2}{3}$ (v) $\frac{1}{6}$

MISCELLANEOUS EXERCISE 5

(I)

1	2	3	4	5	6	7	8	9	10
A	A	C	B	A	D	B	D	A	B

11	12	13	14	15	16	17	18	19	20
A	D	B	B	C	C	A	D	A	C

- (II)
1. (i) 10 (ii) 2 (iii) $\frac{1}{2}$ 5. $\frac{\pi}{3}$ 6. $\frac{1}{6}$ 7. $\frac{\pi}{4} - \frac{1}{2}$
2. 9π 3. 20π 8. $\frac{56}{3}$ 9. $36 \frac{3}{4}$ 10. $\frac{7}{3}$
4. (i) $\frac{16}{3}$ (ii) $\frac{8}{3}$ (iii) $\frac{1}{3}$

6. DIFFERENTIAL EQUATIONS

EXERCISE 6.1

- (1) (i) 2, 1 (ii) 2, 3 (iii) 1, 2 (iii) $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = 0$ (iv) $8 \left(\frac{dy}{dx}\right)^3 - 27y = 0$
- (iv) 3, 1 (v) 2, 1 (vi) 3, 2 (v) $\frac{d^2y}{dx^2} - 25y = 0$ (vi) $2 \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^3 = 0$
- (vii) 2, not defined (viii) 2, 2 (vii) $(x^2 + xy) \frac{dy}{dx} + y = 0$ (viii) $\frac{d^2y}{dx^2} - 7 \frac{dy}{dx} + 10y = 0$
- (ix) 3, 3 (x) 2, 1

EXERCISE 6.2

- (1) (i) $2x^3 + 3xy^2 \frac{dy}{dx} - y^3 = 0$ (ix) $xy \frac{d^2y}{dx^2} + x \left(\frac{dy}{dx}\right)^2 - 2y \frac{dy}{dx} = 0$
- (ii) $xy \frac{d^2y}{dx^2} + x \left(\frac{dy}{dx}\right)^2 - y \frac{dy}{dx} = 0$ (x) $\frac{d^2y}{dx^2} + 4 \frac{dy}{dx} + 5y = 0$

(2) $\frac{d^2y}{dx^2} = 0$ (3) $2a \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^3 = 0$

(4) $x + 4y \frac{dy}{dx} = 0$ (5) $3 \frac{dy}{dx} + 2 = 0$

(6) $81 \left(\frac{d^2y}{dx^2}\right)^2 = \left[\left(\frac{dy}{dx}\right)^2 + 1\right]^3$

(7) $y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 0$

EXERCISE 6.3

(2) (i) $\tan^{-1} y = \tan^{-1} x + c$

(ii) $2e^{-3y} + 3e^{2x} = c$ (iii) $x = cy$

(iv) $\tan x \cdot \tan y = c$ (v) $\sin y \cdot \cos x = c$

(vi) $y = -kx + c$

(vii) $2(x^2 + y^2) + 2(x \sin 2x + y \sin 2y) + \cos 2y + \cos 2x + c = 0$

(viii) $2y^2 \tan^{-1} x + 1 = cy^2$

(ix) $4e^x + 3e^{-2y} = c$

(x) $3e^x + 3e^{-y} + x^3 = c$

(3) (i) $(1 + e^x)^3 \tan y = 0$

(ii) $(1 + x^2)(1 - y^2) = 5$

(iii) $y = ex \log x$ (iv) $(\sin x)(e^y + 1) = \sqrt{2}$

(v) $2(2 + e^y) = 3(x + 1)$

(vi) $\cos\left(\frac{y-2}{x}\right) = a$

(4) (i) $\tan\left(\frac{x+y}{2}\right) = x + c$

(ii) $c + 2y = a \log\left(\frac{x-y-a}{x-y+a}\right)$

(iii) $\sin(x^2 + y^2) + 2x = c$

(iv) $x = \tan(x - 2y) + c$

(v) $(2x - y) - \log(x - y + 2) + 1 = 0$

EXERCISE 6.4

(1) $\cos\left(\frac{y}{x}\right) dy = \log(x) + c$

(2) $x^2 - y^2 = cx$ (3) $x + 2ye^{\frac{x}{y}} = c$

(4) $xy^2 = c^2(x + 2y)$ (5) $x^2 + y^2 = cx$

(6) $y = c(x + y)^3 + x$

(7) $x \left[1 - \cos\left(\frac{y}{x}\right)\right] = \sin\left(\frac{y}{x}\right)$

(8) $x + ye^{\frac{x}{y}} = c$ (9) $\log(y) + \frac{y}{x} = c$

(10) $x^2y = 4$ (11) $x^2 + y^2 = x^4$

(12) $\tan^{-1}\left(\frac{y}{x}\right) = \log(x) + c$

(13) $(3x + y)^3(x + y)^2 = c$

(14) $c = \log(x) + \frac{x}{x+y}$ (15) $x^2 - y^2 = cx$

EXERCISE 6.5

1. (i) $\frac{x^5}{5} - \frac{3x^2}{2} - xy = c$

(ii) $ye^{\tan x} = e^{\tan x}(\tan x - 1) + c$

(iii) $x = y(c + y^2)$

(iv) $y(\sec x + \tan x) = \sec x + \tan x - x + c$

(v) $x^2y = \frac{x^4 \log x}{4} - \frac{x^4}{16} + c$

(vi) $x + y + 1 = ce^y$

(vii) $2y = (x + a)^5 + 2c(x + a)^3$

(viii) $r \sin^2 \theta + \frac{\sin^4 \theta}{2} = c$

(ix) $\frac{y^3}{3} = xy + c$

(x) $y = \sqrt{1 - x^2} + c(1 - x^2)$

(xi) $y = \frac{1}{2} e^{\tan^{-1} x} + c e^{-\tan^{-1} x}$

2. $3(x + 3y) = 2(1 - e^{3x})$

4. $y = 4 - x - 2e^x$

3. $4x^2 + 9y^2 = 36$

5. $1 + y = 2e^{\frac{x^2}{2}}$

EXERCISE 6.6

1. 8 times of original.

2. 95.4 years

3. $36 \cdot 36^\circ\text{C}$

4. 5656

5. $\frac{\log 3}{k}$

6. $\frac{27}{5}$ gms

7. $(3000) \left(\frac{4}{9}\right)^{\frac{t}{40}}$

8. 1 hour

10. $r = 3 - t$

11. 27,182

12. $\left(10 - \frac{P}{10}\right)^2 \%$

MISCELLANEOUS EXERCISE 6

(I)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
D	A	C	B	A	D	C	B	C	D	B	A	B	B	B

(II) (1) (i) 2, 1 (ii) 3, 10 (iii) 2, 3 (iv) 1.4 (v) 4, not defined

(3) (i) $xy \frac{d^2y}{dx^2} + x \left(\frac{dy}{dx}\right)^2 - 2y \frac{dy}{dx} = 0$ (ii) $\frac{d^2y}{dx^2} + y = 0$ (iii) $(y - a) \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = 0$

(iv) $2x^2y \frac{d^2y}{dx^2} + 2x^2 \left(\frac{dy}{dx}\right)^2 + y = 0$ (v) $\frac{d^2y}{dx^2} - 9y = 0$

(4) (i) $2xy \frac{dy}{dx} + x^2 - y^2 = 0$ (ii) $2b \frac{d^2y}{dx^2} - 1 = 0$ (iii) $x + 4y \frac{dy}{dx} = 0$ (iv) $2 \frac{dy}{dx} - 3 = 0$

(5) (i) $2e^{-3y} + 3e^{2x} + 6c = 0$ (ii) $\log(y) = \frac{x^3}{3} + x + c$ (iii) $y = \frac{x}{2} \log(x^2) + 2 + cx$

(iv) $y = 1 + x \log x + cx$ (v) $y = x^2 + c \cdot \operatorname{cosec} x$ (vi) $x \log y = (\log y)^2 + c$

(vii) $4xe^{2y} + 5e^{-y} = c$

(6) (i) $ex \log x - y = 0$ (ii) $x = 2y^2$ (iii) $y \operatorname{cosec}^2 x + 2 = 4 \sin 2x$

(iv) $\log \sqrt{x^2 + y^2} + \tan^{-1} \left(\frac{y}{x}\right) = \frac{\pi}{4}$ (v) $x + 2ye^{\frac{x}{y}} = 2$

(8) $x^2 + y^2 = 4x + 5$ (9) $r = (63t + 27)^{\frac{1}{3}}$ (10) $\frac{20}{9}$ years

7. PROBABILITY DISTRIBUTIONS

EXERCISE 7.1

1. $\{-6, -4, -2, 0, 2, 4, 6\}$

2. $\{0, 1, 2\}$

3. (i) p.m.f. (ii) Not p.m.f

(iii) p.m.f (iv) Not p.m.f

(v) Not p.m.f. (iv) p.m.f

5.

X	0	1	2
$P(X)$	$\frac{2}{3}$	$\frac{2}{9}$	$\frac{1}{9}$

6.

X	0	1	2	3	4
$P(X)$	$\left(\frac{4}{5}\right)^4$	$\left(\frac{4}{5}\right)^3 \frac{1}{5}$	$\left(\frac{4}{5}\right)^2 \left(\frac{1}{5}\right)^2$	$\frac{4}{5} \left(\frac{1}{5}\right)^3$	$\left(\frac{1}{5}\right)^4$

7.

X	0	1	2
$P(X)$	$\frac{9}{16}$	$\frac{3}{8}$	$\frac{1}{16}$

4. (i)

X	0	1	2
$P(X)$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

(ii)

X	0	1	2	3
$P(X)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$

(iii)

X	0	1	2	3	4
$P(X)$	$\frac{1}{16}$	$\frac{1}{4}$	$\frac{3}{8}$	$\frac{1}{4}$	$\frac{1}{16}$

8. (i) $\frac{1}{10}$ (ii) $\frac{3}{10}$ (iii) $\frac{1}{5}$

9. $-0.05, 2.2475$ 10. $\frac{7}{3}, \frac{524}{54}$ 11. 1.5

12. $\frac{1}{3}$ 13. 4.67 14. 2.41

15. $17.53, 4.9, 2.21$ 16. $0.7, 0.21$

EXERCISE 7.2

1. (i) p.d.f. (ii) Not a p.d.f

(iii) Not a p.d.f

2. (a) $\frac{2.25}{16}$, (b) $\frac{3}{16}$, (c) $\frac{3}{4}$

3. (i) p.d.f. (ii) $\frac{1}{9}$ (iii) $\frac{1}{9}$

4. (i) $\frac{1}{2}, \frac{35}{64}$ (ii) $6, \frac{11}{32}, \frac{1}{2}$ 8. (i) $\frac{x^2}{16}$ (ii) $\frac{1}{64}, 0.18, 1$
5. (i) $\frac{1}{4}$ (ii) $\frac{1}{2}$ (iii) $\frac{7}{16}$ 9. $\frac{2}{9}, 0, \frac{8}{9}, \frac{7}{9}$
6. (i) $\frac{2}{5}$ (ii) $\frac{1}{5}$ 10. $\frac{1}{\log 3}, \frac{4}{\log 3}, \frac{4(\log 3 - 1)}{(\log 3)^2}$
7. (i) $\frac{1}{2}$ (ii) $\frac{11}{16}$ (iii) 0.6328

MISCELLANEOUS EXERCISE 7

(I)

1	2	3	4	5	6	7	8	9	10
B	C	A	B	C	B	B	A	D	B

(II) Solve the following :

- (1) (i) Discrete $\{1, 2, 3, \dots, 100000\}$ (ii) Continuous. (iii) Continuous.
 (iv) Discrete $\{0, 1, 2, 3, 4, 5\}$ (v) Continuous
- (2) (i) $\frac{1}{21}$ (ii) $\frac{10}{21}, \frac{1}{7}, \frac{6}{7}$ (3) (i) 0.5 (ii) 0.7 (iii) 0.55 (iv) 0.45
- (5) $\frac{1}{4}$

x	1	2	3	4	5
$P(X)$	$\frac{1}{20}$	$\frac{3}{20}$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{20}$
$F(x)$	$\frac{1}{20}$	$\frac{1}{5}$	$\frac{9}{20}$	$\frac{19}{20}$	1

(6)

X	0	1	2	3	4
$P(X)$	$\frac{1}{16}$	$\frac{1}{4}$	$\frac{3}{8}$	$\frac{1}{4}$	$\frac{1}{16}$

$$\frac{{}^4C_x}{2^4}$$

(7) (i) (ii)

X	0	1	2
$P(X)$	$\frac{4}{9}$	$\frac{4}{9}$	$\frac{1}{9}$

X	0	1	2
$P(X)$	$\frac{25}{36}$	$\frac{10}{36}$	$\frac{1}{36}$

(8) (i) $\frac{1}{10}$ (ii) $\frac{17}{100}$ (iii) $\frac{3}{10}$

(9)

X	-3	-2	-1	0	1	2	3	4
$F(X)$	0.1	0.3	0.5	0.65	0.75	0.85	0.9	1
$P(x)$	0.1	0.2	0.2	0.15	0.10	0.10	0.05	0.10

(i) 0.55 (ii) 0.25

(10) (i) $\frac{11}{5}, \frac{14}{25}, \frac{\sqrt{14}}{5}$ (ii) $\frac{1}{5}, \frac{14}{25}, \frac{\sqrt{14}}{5}$ (iii) $\frac{n+1}{2}, \frac{n^2-1}{12}, \sqrt{\frac{n^2-1}{12}}$ (iv) $\frac{5}{2}, \frac{5}{4}, \frac{\sqrt{5}}{2}$

(11) ₹ 5.5, 8.25 (12) 0, 1 (13) (i) $\frac{1}{2}$ (ii) $\frac{11}{16}$ (iii) $\frac{81}{128}$

(15) $k = \frac{1}{\theta}, \frac{1}{e}$ (16) $k = \frac{1}{4}, F(x) = \frac{\sqrt{x}}{2}, \frac{1}{\sqrt{2}}, \frac{1}{2}$

8. BINOMIAL DISTRIBUTION

EXERCISE 8.1

1. (i) $\frac{3}{2^5}$ (ii) $\frac{7}{2^6}$ (iii) $\frac{63}{64}$
2. $\frac{25}{216}$ 3. $29 \left(\frac{19^9}{20^{10}} \right)$
4. (i) $\frac{1}{1024}$ (ii) $\frac{45}{1024}$
5. (i) $(0.95)^5$ (ii) $(1.2)(0.95)^4$
(iii) $1 - (1.2)(0.95)^4$ (iv) $1 - (0.95)^5$
6. $\left(\frac{9}{10} \right)^4$ 7. $\frac{11}{243}$
8. (i) $1 - \left(\frac{99}{100} \right)^{50}$ (ii) $50 \left(\frac{99^{49}}{100^{50}} \right)$
(iii) $1 - 149 \left(\frac{99^{49}}{100^{50}} \right)$
9. (i) $\frac{1}{20^3}$ (ii) $3 \left(\frac{19}{20^3} \right)$ (iii) $3 \left(\frac{19^2}{20^3} \right)$ (iv) $\left(\frac{19}{20} \right)^3$
10. $\frac{7}{3} \left(\frac{5}{6} \right)^5$ 11. $22 \left(\frac{9^3}{10^{11}} \right)$
12. (i) 4, 2.4 (ii) 10, 2.4 (iii) $\frac{2}{5}; \sqrt{6}$ (iv) $\frac{8}{5}$

MISCELLANEOUS EXERCISE 8

(I)

1	2	3	4	5	6	7
B	D	D	C	B	C	B

(II) Solve the following :

(1) (i) $2 \times (0.8)^9$ (ii) $1 - (0.8)^{10}$

(iii) $1 - (8.2)(0.2)^9$

(2) (i) $p = \frac{1}{2}$, $\text{Var}(X) = 2.5$

(ii) $n = 10$, $p = \frac{1}{2}$

(3) (i) $\frac{63}{256}$ (ii) $\frac{105}{512}$

(4) $45 \left(\frac{2^{26}}{2^{10}} \right)$

(5) (i) $0.65 \times (0.95)^{16}$

(ii) $(2.0325) \times (0.95)^{14}$

(iii) $1 - (1.6) \times (0.95)^{16}$

(6) 0.2114

(7) $1.4 \times (0.9)^4$

(8) $6.97 \times (0.97)^{19}$

(9) 0.3456

(10) $\frac{30.44}{5^8}$

(11) (i) $(0.998)^8$ (ii) $1.014 \times (0.998)^7$

(iii) $1 - 1.014 \times (0.998)^7$

(12) $775.44 \times (0.003)^{38}$

(13) (i) 0.9^{10} (ii) 0.9^9

(iii) $0.45 \times (0.9)^8$

(iv) $1 - 2.16 \times (0.9)^8$

(14) (i) $\frac{1}{5^4}$, $\frac{16}{5^4}$, $\frac{96}{5^4}$, $\frac{256}{5^4}$, $\frac{256}{5^4}$

(ii) (a) $\frac{608}{5^4}$ (b) $1 - \frac{33}{5^8}$

(15) (i) $35 \times 8 \times \frac{81}{5^7}$ (ii) $1 - \frac{12393}{5^7}$

(16) (i) $\frac{\log 0.5}{\log 0.99}$

(17) $\frac{1}{5}$



4!



Maharashtra State Bureau of Textbook
Production and Curriculum Research,
Pune - 411 004.

गणित आणि संख्याशास्त्र (कला व विज्ञान) भाग-२, इ. १२ वी (इंग्रजी माध्यम)

₹ 150.00