

CHAPTER – 6**TISSUES****TISSUES**

The body of plants and animals is made up of different types of cells. These cells originate from a single cell by repeated divisions and get differentiated during development. In unicellular organisms all the body functions are performed by a single cell. But in multicellular organisms, different functions are performed by different groups of cells.

The groups of cells having a common origin and performing similar functions are called **tissues**. Several tissues are organized to form tissue system and the tissue systems form the organs and several organs into organism.



Study of tissues is called **Histology**

Tissue and Division of Labour: In complex organisms, different tasks are carried out by different organs and organ systems. Tissues are the first step towards division of labour in complex organisms.

INTEXT QUESTIONS PAGE NO. 69

Q1. What is a tissue?

Answer: A group of cells that are similar in structure and/or work together to achieve a particular function is called tissue.

Q2. What is the utility of tissues in multi-cellular organisms?

Answer: In multicellular organisms, the body system is based on the division of labour. It means the cells performing a specific function are grouped together to form a particular tissue. The different tissues are organized in a way to provide highest efficiency in functioning of the body.

PLANT TISSUES

Plant tissues are of two main types, viz. meristematic tissue and permanent tissue.

MERISTEMATIC TISSUE


The growth of plants occurs only in certain specific regions. This is because the dividing tissue also known as meristematic tissue (Meristos – divisible) is located only at these points.

The meristematic tissues are made up of group of similar and immature cells, which can divide and form new cells. Meristematic cells divide continuously and thus help in increasing the length and thickness of the plant. Depending upon the position, meristematic tissues are of three types. They are as follows:

i) **Apical meristems:** Apical meristem is present at the growing tips of stems and roots and increases the length of the plant body. They are responsible for growth in length, i.e. primary growth.

ii) **Intercalary meristems:** These meristems occupy base of the leaves and the base of the internodal regions in plants such as grasses (mostly in monocotyledonous plants). These help in elongation of the internodes.

iii) **Lateral meristems:** This includes the meristematic tissues occupying the lateral regions of the stems and roots which bring about increase in the width of the plant body. (e.g. Cork cambium and Vascular cambium).




Characteristic features of Meristematic tissues

- The meristematic cells may be round, oval, polygonal or rectangular in shape.
- Their cell walls are thin, elastic and made up of cellulose.
- They are closely arranged without intercellular spaces.
- They have dense cytoplasm with large nucleus.

PERMANENT TISSUE:

Once the cells of meristematic tissue divide to a certain extent, they become specialized for a particular function. This process is called differentiation. Once differentiation is accomplished, the cells lose their capability to divide and the tissue becomes permanent tissue.

Some cells produced by meristematic tissues stop dividing and form a permanent tissue. They have definite structure and function. They are differentiated into various types to perform different functions.




The permanent tissues are classified as

- i) Simple tissues and
- ii) Complex tissues

SIMPLE TISSUES

A tissue with the cells of similar structure (one type of cells) and function is called simple tissue. It is of three types.

1. Parenchyma
2. Collenchyma
3. Sclerenchyma




PARENCHYMA

The cells of parenchyma have thin cell wall. They are loosely packed; with lot of intercellular spaces between them. They are living cells. They are generally present in all organs of a plant. They are oval or spherical or rectangular or cylindrical in shape. The cell wall is made of cellulose and pectic materials. Parenchyma makes the largest portion of a plant body. Parenchyma mainly works as packing material in plant parts. The main function of parenchyma is to provide support and to store food. In some plant parts, parenchyma has chlorophyll as well. In that case, parenchyma carries out photosynthesis and is then termed as chlorenchyma. In aquatic plants, large air cavities are present in parenchyma. This provides buoyancy to the plant, and then the parenchyma is known as aerenchyma.

COLLENCHYMA

The cells of collenchyma are polygonal in cross section and have unevenly thickened walls. These thickenings are due to the deposition of more cellulose, hemi-cellulose and pectin. The thickening is confined to the corners of the cells. They generally occur in the dicot stem in two or more layers below the epidermis. It is absent in the roots. It also occurs in petiole and pedicel. Like Parenchyma, Collenchyma is also a living tissue. The main function of Collenchyma is to provide strength and flexibility to the growing organs like young stem.



Section of a stem

SCLERENCHYMA


It is a dead tissue. The cells are thick with lignified walls. They give mechanical support to the organs. This has two types of cells – Sclereids and Fibres.

Sclereids

Sclereids are stone cells which are commonly found in shells of the nut, pulp of certain fruits such as Pear and Sapota.

Fibres

The fibres are elongated strands with simple pits throughout its length.



COMPLEX PERMANENT TISSUES

XYLEM

Xylem is mainly concerned with the transport of nutrients, water and minerals upwards in the plant body. It forms a continuous tube through the roots, stems, leaves, flowers and fruits by the fusion of elongated cells.

It is composed of different kinds of cells namely,

1. Tracheids
2. Xylem vessels.
3. Xylem fibres
4. Xylem parenchyma.

Tracheids

Tracheids are elongated, tapering cells with blunt ends. They have lignified secondary wall. They are the chief water conducting elements in Pteridophytes and Gymnosperms.

Xylem vessels


Xylem vessels have perforations at the end and are placed one above the other like a long pipe line. They are seen in the xylem of angiosperms. They conduct water, mineral nutrients and also provide mechanical strength to the plant body.

Xylem Fibres

The fibres of Sclerenchyma associated with the xylem are known as xylem fibres. They give additional mechanical strength to the plant. They are also called wood fibres.

Xylem Parenchyma

The parenchyma cells associated with xylem are known as xylem parenchyma. It is the only living tissue amongst xylem cells. They store food reserves in the form of starch and fat. They also help in conduction of water.



PHLOEM

Phloem conducts food materials from leaves to the other parts of the plant. It is made up of four types of cells.

1. Sieve elements
2. Companion cells
3. Phloem fibres
4. Phloem parenchyma

Sieve elements

Sieve elements are the conducting elements of the phloem. Sieve elements are of two types – sieve cells and sieve tubes.

Sieve cells are present in Pteridophytes and Gymnosperms whereas sieve tubes are present in Angiosperms.

Companion cells

Companion cells are thin walled elongated specialized Parenchyma cells. They are associated with sieve elements. They have a prominent nucleus and cytoplasm. They help the sieve tube in conduction of food materials in angiosperms.

Phloem fibres


The fibres of sclerenchyma associated with phloem are called phloem fibres. They are also called bast-fibres. They give mechanical support to the plant. Among the four types of phloem cells, phloem fibres are the only dead tissues.

Phloem parenchyma

The parenchyma cells associated with phloem are called phloem parenchyma. They store starch and fats.

ANIMAL TISSUES

Animal tissues are of four types, viz. epithelial tissue, connective tissue, muscular tissue and nervous tissue.



EPITHELIAL TISSUE:

The epithelial tissue forms the covering or lining of most of the organs. The cells of epithelial tissue are tightly packed and form a continuous sheet. There is small amount of cementing materials between the cells and no intercellular space is present. Permeability of the epithelial tissue plays a great role in exchange of materials among various organs it also plays an important role in osmoregulation. All epithelial tissues are separated by the underlying tissue by an extracellular fibrous basement membrane.

Epithelial tissues are of following types:


1. Simple Epithelium
2. Cuboidal Epithelium
3. Columnar Epithelium
4. Stratified Epithelium

Simple Epithelium

The simple epithelium is composed of a single layer of cells. This type of epithelial tissue forms the lining of blood vessels and alveoli. Thin layer of cells facilitates exchange of substances; in such cases.

Cuboidal Epithelium

The cells are cube-shaped in cuboidal epithelium. Linings of kidney tubules and ducts of salivary glands are composed of cuboidal epithelium. Cuboidal cells provide mechanical support. Cells of epithelium may play the role of secretion and then they are called glandular epithelium.



Columnar Epithelium

Cells are column-shaped in columnar epithelium. Columnar epithelium facilitates secretion and absorption. For example; the lining of intestine is composed of columnar epithelium. In some organs, columnar epithelium has cilia present on the outer surface. Cilia facilitate movements of certain substances. The ciliated epithelium in the respiratory tract pushes the mucus forward.

Stratified Epithelium

Cells of the stratified epithelium are in many layers. Skin is an example of stratified epithelium. Stratification of layers prevents wear and tear.

CONNECTIVE TISSUE:

The cells of a connective tissue are loosely scattered in a matrix. The matrix can be a fluid, jelly like, dense or rigid. The nature of matrix depends on the function a connective tissue serves. Following are the various connective tissues:

Areolar (Loose) Connective Tissue

Areolar tissue is found between skin and muscles, around blood vessels and nerves and in bone marrow. Areolar tissue fills the gap between tissues and provides support. It also helps in repair of tissues.

Dense connective tissue(Fibrous connective tissue)

It has thicker, denser fibers and fewer cells. The matrix is made up mostly of collagen fibers, with fibroblasts arranged in rows. This type of connective tissue forms tendons and ligaments, which attach muscle to bone and bone to bone, respectively.

Adipose Tissue

Adipose tissue is composed of fat globules. This tissue is found below the skin and beneath the organs. Adipose tissue provides insulation and works as a cushion.

Bone


Bone is mainly composed of osteoblasts. Bone makes the skeletal system. Skeletal system is responsible for providing structural framework to the body. It provides protection to important organs and facilitates movements.

Cartilage

Cartilage is mainly composed of chondrioblasts. Cartilage is present at the ends of articular bones. Cartilage is also present in external ear, bronchii, etc.


Blood

Blood is composed of blood cells, platelets and plasma. Blood plays an important role in transportation of various substances in the body. It also helps in osmoregulation and temperature control.



MUSCULAR TISSUE

Muscular tissue is composed of muscle cells. Muscle cells are specialized cells which have the capability to contract and expand. Due to contraction and expansion, muscles facilitate various kinds of movements in the body. Muscular tissues are of three types:



Striated Muscles


The cells of striated muscles are in the form of long, unbranched fibres. Cells are multinucleate. Light and dark bands (striations) are present on muscle fibres, which gives the name striated muscles. Striated muscles are found in those organs where voluntary movement is possible, e.g. hands, legs, back, neck, etc.

Smooth Muscles

The cells of smooth muscles are spindle shaped and each has one nucleus. Smooth muscle is found in those organs where involuntary movement is possible, e.g. alimentary canal.

Cardiac Muscles

The cells of cardiac muscles are in the form of branched fibres. Striations are present and cells are uninucleate. These are found in the heart. Cardiac muscles are capable continuous contraction and relaxation throughout the life.



Smooth muscle

- has spindle-shaped, nonstriated uninucleated fibers.
- occurs in walls of internal organs.
- is involuntary.

Cardiac muscle

- has striated, branched, uninucleated fibers.
- occurs in walls of heart.
- is involuntary.


Skeletal muscle

- has striated, tubular, multinucleated fibers.
- is usually attached to skeleton.
- is voluntary.

NERVOUS TISSUE

All cells possess the ability to respond to stimuli. However, cells of the nervous tissue are highly specialised for being stimulated and then transmitting the stimulus very rapidly from one place to another within the body. The brain, spinal cord and nerves are all composed of the nervous tissue. The cells of this tissue are called nerve cells or neurons.


A neuron consists of a cell body with a nucleus and cytoplasm, from which long thin hair-like parts arise.



Usually each neuron has a single long part, called the axon, and many short, branched parts called dendrites. An individual nerve cell may be up to a metre long. Many nerve fibres bound together by connective tissue make up a nerve.





SUMMARY

CLASSIFICATION OF PLANT TISSUE




TYPES OF EPITHELIAL TISSUE

Different epithelia show different structures as they perform different functions

Type of Epithelium	Structure	Location in the body	Function
Squamous epithelium 	Cells are thin, flat, irregular cells which fit like floor tiles to form delicate lining called PAVEMENT EPITHELIUM . Nuclei in centre	Oesophagus, lining of mouth, alveoli of the lungs, blood vessels	Protects the underlying tissue from injury, grems Exchange of gases in lungs and materials between cells and blood
Cuboidal epithelium 	Cells are cuboidal with round nucleus in centre Nuclei in centre	Kidney tubules, duct of salivary glands	Gives mechanical support At times the epithelial tissue folds, forms a gland that secretes substances. Such epithelium is called GLANDULAR EPITHELIUM
Columnar epithelium 	Cells are more tall and less wide (PILLAR LIKE), placed side by side. Nucleus is situated near the base. Nuclei near base	Inner lining of intestine, In respiratory tract, cells have cilia (hair like) that move and push the mucous to clear it. Such epithelium is called CILIATED COLUMNAR EPITHELIUM	Helps in absorption excretion and secretion
Striated squamous epithelium 	Squamous flat cells arranged in many layers to prevent wear and tear of parts.	Skin (to prevent wear and tear) tongue, oesophagus lining of mouth.	Protection, prevent wear and tear

CLASSIFICATION OF ANIMAL TISSUE



INTEXT QUESTIONS PAGE NO. 74

Q1. Name types of simple tissues.

Answer: The three main types of simple tissues are: (i) Parenchyma (ii) Collenchyma (iii) Sclerenchyma

Q2. Where is apical meristem found?

Answer: Apical meristem is present in growing tips of stems and roots of plants. It helps in increasing the length of the stem and the root.

Q3. Which tissue makes up the husk of coconut?

Answer: The husk of coconut is made up of sclerenchymatous tissue.

Q4. What are the constituents of phloem?

Answer: The constituents of phloem tissue are:

- (i) Sieve tubes (tubular living cells with perforated end walls)
- (ii) Companion cell (living cells)
- (iii) Phloem parenchyma (living cells)
- (iv) Phloem fibres (non-living and sclerenchyma cells)


INTEXT QUESTIONS PAGE NO. 78

Q1. Name the tissue responsible for movement in our body.

Answer: The movement of our body depends on muscular tissue. It consists of elongated cells (muscle fibres).

Q2. What does a neuron look like?

Answer: A neuron consists of a cell body with a nucleus and cytoplasm. It has two important extensions known as the axon and dendrites. An axon is a long thread-like extension of nerve cells that transmits impulses away from the cell body. Dendrites, on the other hand, are thread-like extensions of cell body that receive nerve impulses. Thus, the axon transmits impulses away from the cell body, whereas the dendrite receives nerve impulses. This coordinated function helps in transmitting impulses very quickly.



Q3. Give three features of cardiac muscles.

Answer: Three features of cardiac muscles are:

- (i) Cardiac muscles are involuntary muscles that contract rapidly, but do not get fatigued.
- (ii) The cells of cardiac muscles are cylindrical, branched, and uninucleate.
- (iii) They control the contraction and relaxation of the heart.

Q4. What are the functions of areolar tissue?

Answer: Functions of areolar tissue:

- (i) It helps in supporting internal organs.
- (ii) It helps in repairing the tissues of the skin and muscles.

EXERCISE QUESTIONS PAGE NO. 66 and 67

Q1. Define the term “tissue”.

Answer: A group of cells that are similar in structure and/or work together to achieve a particular function is called tissue.

Q2. How many types of elements together make up the xylem tissue? Name them.

Answer: The following four types of elements make up xylem tissue:

- (i) Xylem tracheids (tubular unicellular).
- (ii) Xylem vessels (multicellular).
- (iii) Xylem parenchyma (stores food and helps in sideways conduction of water).
- (iv) Xylem fibres (provide mechanical support).

Q3. How are simple tissues different from complex tissues in plants?

Answer:

Simple tissue	Complex tissue
These tissues consist of only one type of cells.	These tissues are made up of more than one type of cells.
The cells are more or less similar in structure and perform similar functions.	Different types of cells perform different functions. For example, in the xylem tissue, tracheids help in water transport, whereas parenchyma stores food.
Three types of simple tissues in plants are parenchyma, collenchyma, and sclerenchyma.	Two types of complex permanent tissues in plants are xylem and phloem.

Q4. Differentiate between parenchyma, collenchyma and sclerenchyma on the basis of their cell wall.

Answer:

Parenchyma	Collenchyma	Sclerenchyma
Cell walls are relatively thin, and the cells in parenchyma tissues are loosely packed.	The cell wall is irregularly thickened at the corners, and there is very little space between the cells.	The cell walls are uniformly thickened, and there are no intercellular spaces.
The cell wall in this tissue is made up of cellulose.	Pectin and hemicellulose are the major constituents of the cell wall.	An additional layer of the cell wall composed mainly of lignin is found.

Q5. What are the functions of the stomata?

Answer: Functions of the stomata:


- (i) They allow the exchange of gases (CO₂ and O₂) with the atmosphere.
- (ii) Evaporation of water from the leaf surface occurs through the stomata. Thus, the stomata help in the process of transpiration.

Q6. Diagrammatically show the difference between the three types of muscle fibres.

Answer:

The three types of muscle fibres are:

Striated muscles, smooth muscles (unstriated muscle fibre), and cardiac muscles.



Smooth muscle

- has spindle-shaped, nonstriated uninucleated fibers.
- occurs in walls of internal organs.
- is involuntary.

Cardiac muscle

- has striated, branched, uninucleated fibers.
- occurs in walls of heart.
- is involuntary.

Skeletal muscle

- has striated, tubular, multinucleated fibers.
- is usually attached to skeleton.
- is voluntary.

Q7. What is the specific function of the cardiac muscle?

Answer: The specific function of the cardiac muscle is to control the contraction and relaxation of the heart.


Q8. Differentiate between striated, unstriated and cardiac muscles on the basis of their structure and site/location in the body.

Answer:

Striated muscle	Unstriated muscle	Cardiac muscle
On the basis of structure:		
Cells are cylindrical	Cells are long	Cells are cylindrical
Cells are not branched	Cells are not branched	Cells are branched
Cells are multinucleate	Cells are uninucleate	Cells are uninucleate
Alternate light and dark bands are present	There are no bands present	Faint bands are present
Its ends are blunt	Its ends are tapering	Its ends are flat and wavy
On the basis of location:		
These muscles are present in body parts such as hands, legs, tongue, etc.	These muscles control the movement of food in the alimentary canal, the contraction and relaxation of blood vessels, etc.	These muscles control the contraction and relaxation of the heart

Q9. Draw a labelled diagram of a neuron.

Answer:



Q10. Name the following.

- (a) Tissue that forms the inner lining of our mouth.
- (b) Tissue that connects muscle to bone in humans.
- (c) Tissue that transports food in plants.
- (d) Tissue that stores fat in our body.
- (e) Connective tissue with a fluid matrix.
- (f) Tissue present in the brain.

Answer:

- (a) Tissue that forms the inner lining of our mouth → Epithelial tissue
- (b) Tissue that connects muscle to bone in humans → Dense regular connective tissue (tendons)
- (c) Tissue that transports food in plants → Phloem
- (d) Tissue that stores fat in our body → Adipose tissue
- (e) Connective tissue with a fluid matrix → Blood
- (f) Tissue present in the brain → Nervous tissue

Q11. Identify the type of tissue in the following: skin, bark of tree, bone, lining of kidney tubule, vascular bundle.

Answer:

Skin: Stratified squamous epithelial tissue

Bark of tree: Simple permanent tissue

Bone: Connective tissue

Lining of kidney tubule: Cuboidal epithelial tissue

Vascular bundle: Complex permanent tissue

Q12. Name the regions in which parenchyma tissue is present.

Answer:

Leaves, fruits, and flowers are the regions where the parenchyma tissue is present.

Q13. What is the role of epidermis in plants?

Answer:

Epidermis is present on the outer surface of the entire plant body. The cells of the epidermal tissue form a continuous layer without any intercellular space. It performs the following important functions:


- (i) It is a protective tissue of the plant body
- (ii) It protects the plant against mechanical injury
- (iii) It allows exchange of gases through the stomata

Q14. How does the cork act as a protective tissue?

Answer:

The outer protective layer or bark of a tree is known as the cork. It is made up of dead cells. Therefore, it protects the plant against mechanical injury, temperature extremes, etc. It also prevents the loss of water by evaporation.

Q15. Complete the table:



Answer:

